

Discrete Mathematics for Computer Science

Lecture 8: Number Theory

Dr. Ming Tang

Department of Computer Science and Engineering
Southern University of Science and Technology (SUSTech)
Email: tangm3@sustech.edu.cn

Congruence Relation

If a and b are integers and m is a positive integer, then a is congruent to b modulo m if m divides $a - b$, denoted by $a \equiv b \pmod{m}$.

The integers a and b are congruent modulo m if and only if a and b have the same remainder when divided by m , i.e.,

$$a \bmod m = b \bmod m$$

Computing the mod Function

$$(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$$

$$ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$$


Proof: By the definitions of $\bmod m$ and of congruence modulo m , we know that $a \equiv (a \bmod m)(\bmod m)$ and $b \equiv (b \bmod m)(\bmod m)$. Hence,

$$a + b \equiv (a \bmod m) + (b \bmod m)(\bmod m)$$

$$ab \equiv (a \bmod m)(b \bmod m)(\bmod m).$$

According to the theorem that two integers are congruent modulo m if and only if they have the **same remainder**, we complete our proof.

This Lecture



Number Theory: divisibility and modular arithmetic, integer representations, primes, greatest common divisors, linear congruences, ...



SUSTech

Southern University
of Science and
Technology

Base-b Expansions

From decimal expansion to the base-b expansion:

Base-b Expansions

From decimal expansion to the base-b expansion:

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a}_0 \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a}_1) + \textcolor{blue}{a}_0 \\&= \cdots\end{aligned}$$

Base-b Expansions

From decimal expansion to the base-b expansion:

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a}_0 \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a}_1) + \textcolor{blue}{a}_0 \\&= \cdots\end{aligned}$$

- Divide $\textcolor{red}{n}$ by b to obtain $n = bq_0 + \textcolor{blue}{a}_0$, with $0 \leq \textcolor{blue}{a}_0 < b$

Base-b Expansions

From decimal expansion to the base-b expansion:

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a}_0 \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a}_1) + \textcolor{blue}{a}_0 \\&= \cdots\end{aligned}$$

- Divide $\textcolor{red}{n}$ by b to obtain $n = bq_0 + \textcolor{blue}{a}_0$, with $0 \leq a_0 < b$
- The $\textcolor{blue}{remainder}$ $\textcolor{blue}{a}_0$ is the rightmost digit in the base- b expansion of n .
Then divide $\textcolor{red}{q}_0$ by b to get $q_0 = bq_1 + \textcolor{blue}{a}_1$ with $0 \leq a_1 < b$;

Base-b Expansions

From decimal expansion to the base-b expansion:

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + a_0 \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + a_1) + a_0 \\&= \cdots\end{aligned}$$

- Divide n by b to obtain $n = bq_0 + a_0$, with $0 \leq a_0 < b$
- The remainder a_0 is the rightmost digit in the base- b expansion of n . Then divide q_0 by b to get $q_0 = bq_1 + a_1$ with $0 \leq a_1 < b$;
- a_1 is the second digit from the right; continue by successively dividing the quotients by b until the quotient is 0

Base- b Expansions

```
procedure base  $b$  expansion( $n, b$ : positive integers with  $b > 1$ )  
   $q := n$   
   $k := 0$   
  while ( $q \neq 0$ )  
     $a_k := q \bmod b$   
     $q := q \operatorname{div} b$   
     $k := k + 1$   
  return( $a_{k-1}, \dots, a_1, a_0$ )  $\{(a_{k-1} \dots a_1 a_0)_b$  is base  $b$  expansion of  $n\}$ 
```

Base-b Expansions

Example: Find the hexadecimal expansion of $(177130)_{10}$.

Base-b Expansions

Example: Find the hexadecimal expansion of $(177130)_{10}$.

Solution: First divide 177130 by 16 to obtain

$$177130 = 16 \cdot 11070 + 10.$$

Base-b Expansions

Example: Find the hexadecimal expansion of $(177130)_{10}$.

Solution: First divide 177130 by 16 to obtain

$$177130 = 16 \cdot 11070 + 10.$$

Successively dividing quotients by 16 gives

$$11070 = 16 \cdot 691 + 14,$$

$$691 = 16 \cdot 43 + 3,$$

$$43 = 16 \cdot 2 + 11,$$

$$2 = 16 \cdot 0 + 2.$$

The successive remainders that we have found, 10, 14, 3, 11, 2. It follows that $(177130)_{10} = (2B3EA)_{16}$.

Algorithms for Integer Operations

- Binary addition
- Binary multiplication
- div and mod
- Modular exponentiation

Binary Addition of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0), b = (b_{n-1}b_{n-2}\dots b_1b_0)$$

$$\begin{array}{r} 1\ 1\ 1 \\ 1\ 1\ 1\ 0 \\ + 1\ 0\ 1\ 1 \\ \hline 1\ 1\ 0\ 0\ 1 \end{array}$$

Binary Addition of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0), b = (b_{n-1}b_{n-2}\dots b_1b_0)$$

$$\begin{array}{r} 1\ 1\ 1 \\ 1\ 1\ 1\ 0 \\ + 1\ 0\ 1\ 1 \\ \hline 1\ 1\ 0\ 0\ 1 \end{array}$$

Let c be the **carry** and $s = (s_ns_{n-1}\dots s_0)_2$ be the **sum**:

```
procedure add( $a, b$ : positive integers)
{the binary expansions of  $a$  and  $b$  are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
 $c := 0$ 
for  $j := 0$  to  $n - 1$ 
     $d := \lfloor (a_j + b_j + c) / 2 \rfloor$ 
     $s_j := a_j + b_j + c - 2d$ 
     $c := d$ 
 $s_n := c$ 
return  $(s_0, s_1, \dots, s_n)$  {the binary expansion of the sum is  $(s_n, s_{n-1}, \dots, s_0)_2$ }
```



SUSTech

Southern University
of Science and
Technology

Binary Addition of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0), b = (b_{n-1}b_{n-2}\dots b_1b_0)$$

$$\begin{array}{r} 1\ 1\ 1 \\ 1\ 1\ 1\ 0 \\ + 1\ 0\ 1\ 1 \\ \hline 1\ 1\ 0\ 0\ 1 \end{array}$$

Let c be the **carry** and $s = (s_ns_{n-1}\dots s_0)_2$ be the **sum**:

```
procedure add( $a, b$ : positive integers)
{the binary expansions of  $a$  and  $b$  are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
 $c := 0$ 
for  $j := 0$  to  $n - 1$ 
     $d := \lfloor (a_j + b_j + c) / 2 \rfloor$ 
     $s_j := a_j + b_j + c - 2d$ 
     $c := d$ 
 $s_n := c$ 
return  $(s_0, s_1, \dots, s_n)$  {the binary expansion of the sum is  $(s_n, s_{n-1}, \dots, s_0)_2$ }
```

$O(n)$ bit additions.

Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2}\dots b_1b_0)_2$$

$$ab = a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) = a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1})$$

Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2}\dots b_1b_0)_2$$

$$ab = a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) = a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1})$$

$$\begin{array}{r} 110 \\ \times 101 \\ \hline 110 \\ 000 \\ 110 \\ \hline 11110 \end{array}$$

Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2}\dots b_1b_0)_2$$

$$ab = a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) = a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1})$$

Let $c_j = ab_j2^j$ and p be the **product**:

```
procedure multiply( $a, b$ : positive integers)
{the binary expansions of  $a$  and  $b$  are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
for  $j := 0$  to  $n - 1$ 
    if  $b_j = 1$  then  $c_j = a$  shifted  $j$  places
    else  $c_j := 0$ 
{ $c_0, c_1, \dots, c_{n-1}$  are the partial products}
 $p := 0$ 
for  $j := 0$  to  $n - 1$ 
     $p := \text{add}(p, c_j)$ 
return  $p$  { $p$  is the value of  $ab$ }
```

$$\begin{array}{r} 110 \\ \times 101 \\ \hline 110 \\ 000 \\ 110 \\ \hline 11110 \end{array}$$

Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2}\dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2}\dots b_1b_0)_2$$

$$ab = a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) = a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1})$$

Let $c_j = ab_j2^j$ and p be the **product**:

```
procedure multiply( $a, b$ : positive integers)
{the binary expansions of  $a$  and  $b$  are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
for  $j := 0$  to  $n - 1$ 
    if  $b_j = 1$  then  $c_j = a$  shifted  $j$  places
    else  $c_j := 0$ 
{ $c_0, c_1, \dots, c_{n-1}$  are the partial products}
 $p := 0$ 
for  $j := 0$  to  $n - 1$ 
     $p := \text{add}(p, c_j)$ 
return  $p$  { $p$  is the value of  $ab$ }
```

$$\begin{array}{r} 110 \\ \times 101 \\ \hline 110 \\ 000 \\ 110 \\ \hline 11110 \end{array}$$

$O(n^2)$ shifts and $O(n^2)$ bit additions.

Computing div and mod

Consider integers a and d . Compute $q = a \operatorname{div} d$ and $r = a \operatorname{mod} d$:

Computing div and mod

Consider integers a and d . Compute $q = a \text{ div } d$ and $r = a \text{ mod } d$:

Let q be the quotient, and r be the remainder:

```
procedure division algorithm ( $a$ : integer,  $d$ : positive integer)
   $q := 0$ 
   $r := |a|$ 
  while  $r \geq d$ 
     $r := r - d$ 
     $q := q + 1$ 
  if  $a < 0$  and  $r > 0$  then
     $r := d - r$ 
     $q := -(q+1)$ 
  return ( $q, r$ ) { $q = a \text{ div } d$  is the quotient,  $r = a \text{ mod } d$  is the remainder }
```

Computing div and mod

Consider integers a and d . Compute $q = a \text{ div } d$ and $r = a \text{ mod } d$:

Let q be the quotient, and r be the remainder:

```
procedure division algorithm ( $a$ : integer,  $d$ : positive integer)
   $q := 0$ 
   $r := |a|$ 
  while  $r \geq d$ 
     $r := r - d$ 
     $q := q + 1$ 
  if  $a < 0$  and  $r > 0$  then
     $r := d - r$ 
     $q := -(q+1)$ 
  return ( $q, r$ ) { $q = a \text{ div } d$  is the quotient,  $r = a \text{ mod } d$  is the remainder }
```

$O(q \log a)$ bit operations.

Binary Modular Exponentiation

Consider large integers b , n , and m . Compute $b^n \bmod m$:

Binary Modular Exponentiation

Consider large integers b , n , and m . Compute $b^n \bmod m$:

Let $n = (a_{k-1} \dots a_1 a_0)_2$.

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Binary Modular Exponentiation

Consider large integers b , n , and m . Compute $b^n \bmod m$:

Let $n = (a_{k-1} \dots a_1 a_0)_2$.

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Successively finds $b \bmod m$, $b^2 \bmod m$, $b^4 \bmod m$, . . . , $b^{2^{k-1}} \bmod m$, and multiplies together the terms b^{2^j} , where $a_j = 1$.

Binary Modular Exponentiation

Consider large integers b , n , and m . Compute $b^n \bmod m$:

Let $n = (a_{k-1} \dots a_1 a_0)_2$.

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Successively finds $b \bmod m$, $b^2 \bmod m$, $b^4 \bmod m$, . . . , $b^{2^{k-1}} \bmod m$, and multiplies together the terms b^{2^j} , where $a_j = 1$.

```
procedure modular_exponentiation( $b$ : integer,  $n = (a_{k-1} a_{k-2} \dots a_1 a_0)_2$ ,  $m$ : positive integers)
   $x := 1$ 
   $power := b \bmod m$ 
  for  $i := 0$  to  $k - 1$ 
    if  $a_i = 1$  then  $x := (x \cdot power) \bmod m$ 
     $power := (power \cdot power) \bmod m$ 
  return  $x$  { $x$  equals  $b^n \bmod m$ }
```

Recall that $ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$.

Binary Modular Exponentiation

Use the algorithm to find $3^{644} \bmod 645$:

```
procedure modular_exponentiation(b: integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ , m: positive integers)
  x := 1
  power := b mod m
  for i := 0 to k - 1
    if  $a_i = 1$  then x := (x · power) mod m
    power := (power · power) mod m
  return x {x equals  $b^n \bmod m$ }
```

Binary Modular Exponentiation

Use the algorithm to find $3^{644} \bmod 645$:

```
procedure modular_exponentiation(b: integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ , m: positive integers)
  x := 1
  power := b mod m
  for i := 0 to k - 1
    if  $a_i = 1$  then  $x := (x \cdot \text{power}) \bmod m$ 
     $\text{power} := (\text{power} \cdot \text{power}) \bmod m$ 
  return x {x equals  $b^n \bmod m$ }
```

The algorithm initially sets $x = 1$ and $\text{power} = 3 \bmod 645 = 3$. The binary expansion of 644 is $(1010000100)_2$. Here are the steps used:

$i = 0$: Because $a_0 = 0$, we have $x = 1$ and $\text{power} = 3^2 \bmod 645 = 9 \bmod 645 = 9$;

$i = 1$: Because $a_1 = 0$, we have $x = 1$ and $\text{power} = 9^2 \bmod 645 = 81 \bmod 645 = 81$;

$i = 2$: Because $a_2 = 1$, we have $x = 1 \cdot 81 \bmod 645 = 81$ and $\text{power} = 81^2 \bmod 645 = 6561 \bmod 645 = 111$;

$i = 3$: Because $a_3 = 0$, we have $x = 81$ and $\text{power} = 111^2 \bmod 645 = 12,321 \bmod 645 = 66$;

$i = 4$: Because $a_4 = 0$, we have $x = 81$ and $\text{power} = 66^2 \bmod 645 = 4356 \bmod 645 = 486$;

Binary Modular Exponentiation

Use the algorithm to find $3^{644} \bmod 645$:

```
procedure modular_exponentiation(b: integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ , m: positive integers)
  x := 1
  power := b mod m
  for i := 0 to k - 1
    if  $a_i = 1$  then x := (x · power) mod m
    power := (power · power) mod m
  return x {x equals  $b^n \bmod m$ }
```

The algorithm initially sets $x = 1$ and $power = 3 \bmod 645 = 3$. The binary expansion of 644 is $(1010000100)_2$. Here are the steps used:

$i = 0$: Because $a_0 = 0$, we have $x = 1$ and $power = 3^2 \bmod 645 = 9 \bmod 645 = 9$;

$i = 1$: Because $a_1 = 0$, we have $x = 1$ and $power = 9^2 \bmod 645 = 81 \bmod 645 = 81$;


$i = 2$: Because $a_2 = 1$, we have $x = 1 \cdot 81 \bmod 645 = 81$ and $power = 81^2 \bmod 645 = 6561 \bmod 645 = 111$;

$i = 3$: Because $a_3 = 0$, we have $x = 81$ and $power = 111^2 \bmod 645 = 12,321 \bmod 645 = 66$;

$i = 4$: Because $a_4 = 0$, we have $x = 81$ and $power = 66^2 \bmod 645 = 4356 \bmod 645 = 486$;

$O((\log m)^2 \log n)$ bit operations.

Next Lecture



Number Theory: divisibility and modular arithmetic, integer representations, **primes and greatest common divisors**, linear congruences,

...

Primes

A integer p that is greater than 1 is called a **prime** if the **only** positive factors of p are 1 and p .

Primes

A integer p that is greater than 1 is called a **prime** if the **only** positive factors of p are 1 and p .

A positive integer that is greater than 1 and is **not prime** is called **composite**.

Primes

A integer p that is greater than 1 is called a **prime** if the **only** positive factors of p are 1 and p .

A positive integer that is greater than 1 and is **not prime** is called **composite**.

Fundamental Theorem of Arithmetic: Every integer greater than 1 can be written **uniquely** as a **prime** or as **the product of two or more primes** where the prime factors are written in order of **nondecreasing** size.

Primes and Composites

How to determine whether a number is a prime or a composite?

Primes and Composites

How to determine whether a number is a prime or a composite?

Approach 1: test if each number $x < n$ divides n .

Primes and Composites

How to determine whether a number is a prime or a composite?

Approach 1: test if each number $x < n$ divides n .

Approach 2: test if each **prime** number $x < n$ divides n .

Primes and Composites

How to determine whether a number is a prime or a composite?

Approach 1: test if each number $x < n$ divides n .

Approach 2: test if each **prime** number $x < n$ divides n .

Approach 3: test if each **prime** number $x \leq \sqrt{n}$ divides n .

Primes and Composites

If n is composite, then n has a prime divisor less than or equal to \sqrt{n} .

Primes and Composites

If n is composite, then n has a prime divisor less than or equal to \sqrt{n} .

Proof: If n is composite, then it has a positive integer factor a such that $1 < a < n$ by definition. This means that $n = ab$, where b is an integer greater than 1.

Primes and Composites

If n is composite, then n has a prime divisor less than or equal to \sqrt{n} .

Proof: If n is composite, then it has a positive integer factor a such that $1 < a < n$ by definition. This means that $n = ab$, where b is an integer greater than 1.

Assume that $a > \sqrt{n}$ and $b > \sqrt{n}$. Then, $ab > n$, which leads to a contradiction. So either $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.

Primes and Composites

If n is composite, then n has a prime divisor less than or equal to \sqrt{n} .

Proof: If n is composite, then it has a positive integer factor a such that $1 < a < n$ by definition. This means that $n = ab$, where b is an integer greater than 1.

Assume that $a > \sqrt{n}$ and $b > \sqrt{n}$. Then, $ab > n$, which leads to a contradiction. So either $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.

Thus, n has a divisor less than \sqrt{n} .

Primes and Composites

If n is composite, then n has a prime divisor less than or equal to \sqrt{n} .

Proof: If n is composite, then it has a positive integer factor a such that $1 < a < n$ by definition. This means that $n = ab$, where b is an integer greater than 1.

Assume that $a > \sqrt{n}$ and $b > \sqrt{n}$. Then, $ab > n$, which leads to a contradiction. So either $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.

Thus, n has a divisor less than \sqrt{n} .

By the Fundamental Theorem of Arithmetic, this divisor is either prime, or is a product of primes. In either case, n has a prime divisor less than \sqrt{n} .

Primes

There are infinitely many primes.

Primes

There are infinitely many primes.

Proof: We will prove this theorem using a proof by contradiction. We assume that there are only finitely many primes, p_1, p_2, \dots, p_n . Let

$$Q = p_1 p_2 \dots p_n + 1.$$

Primes

There are infinitely many primes.

Proof: We will prove this theorem using a proof by contradiction. We assume that there are only finitely many primes, p_1, p_2, \dots, p_n . Let

$$Q = p_1 p_2 \dots p_n + 1.$$

By the fundamental theorem of arithmetic, Q is prime or else it can be written as the product of two or more primes.

Primes

There are infinitely many primes.

Proof: We will prove this theorem using a proof by contradiction. We assume that there are only finitely many primes, p_1, p_2, \dots, p_n . Let

$$Q = p_1 p_2 \dots p_n + 1.$$

By the fundamental theorem of arithmetic, Q is prime or else it can be written as the product of two or more primes.

However, none of the primes p_j divides Q , for if $p_j | Q$, then p_j divides $Q - p_1 p_2 \dots p_n = 1$.

Primes

There are infinitely many primes.

Proof: We will prove this theorem using a proof by contradiction. We assume that there are only finitely many primes, p_1, p_2, \dots, p_n . Let

$$Q = p_1 p_2 \dots p_n + 1.$$

By the fundamental theorem of arithmetic, Q is prime or else it can be written as the product of two or more primes.

However, none of the primes p_j divides Q , for if $p_j | Q$, then p_j divides $Q - p_1 p_2 \dots p_n = 1$.

Hence, there is a prime not in the list p_1, p_2, \dots, p_n . This prime is either Q , if it is prime, or a prime factor of Q .

Primes

There are infinitely many primes.

Proof: We will prove this theorem using a proof by contradiction. We assume that there are only finitely many primes, p_1, p_2, \dots, p_n . Let

$$Q = p_1 p_2 \dots p_n + 1.$$

By the fundamental theorem of arithmetic, Q is prime or else it can be written as the product of two or more primes.

However, none of the primes p_j divides Q , for if $p_j | Q$, then p_j divides $Q - p_1 p_2 \dots p_n = 1$.

Hence, there is a prime not in the list p_1, p_2, \dots, p_n . This prime is either Q , if it is prime, or a prime factor of Q .

This is a contradiction because we assumed that we have listed all the primes. Consequently, there are infinitely many primes.



SUSTech

Southern University
of Science and
Technology

Greatest Common Divisor (GCD)

Let a and b be integers, not both 0. The **largest** integer d such that $d|a$ and $d|b$ is called the **greatest common divisor** of a and b , denoted by $\gcd(a, b)$.

Greatest Common Divisor (GCD)

Let a and b be integers, not both 0. The **largest** integer d such that $d|a$ and $d|b$ is called the **greatest common divisor** of a and b , denoted by $\gcd(a, b)$.

Example: What is the greatest common divisor of 24 and 36?

Greatest Common Divisor (GCD)

Let a and b be integers, not both 0. The **largest** integer d such that $d|a$ and $d|b$ is called the **greatest common divisor** of a and b , denoted by $\gcd(a, b)$.

Example: What is the greatest common divisor of 24 and 36?
 $\gcd(24, 36) = 12$.

Greatest Common Divisor (GCD)

Let a and b be integers, not both 0. The **largest** integer d such that $d|a$ and $d|b$ is called the **greatest common divisor** of a and b , denoted by $\gcd(a, b)$.

Example: What is the greatest common divisor of 24 and 36?
 $\gcd(24, 36) = 12$.

Integers a and b are **relatively prime** if their greatest common divisor is 1.

Greatest Common Divisor (GCD)

Let a and b be integers, not both 0. The **largest** integer d such that $d|a$ and $d|b$ is called the **greatest common divisor** of a and b , denoted by $\gcd(a, b)$.

Example: What is the greatest common divisor of 24 and 36?
 $\gcd(24, 36) = 12$.

Integers a and b are **relatively prime** if their greatest common divisor is 1.

Example: Are integers 17 and 22 relatively prime?

Greatest Common Divisor (GCD)

Let a and b be integers, not both 0. The **largest** integer d such that $d|a$ and $d|b$ is called the **greatest common divisor** of a and b , denoted by $\gcd(a, b)$.

Example: What is the greatest common divisor of 24 and 36?
 $\gcd(24, 36) = 12$.

Integers a and b are **relatively prime** if their greatest common divisor is 1.

Example: Are integers 17 and 22 relatively prime? Yes, because $\gcd(17, 22) = 1$.

Greatest Common Divisor (GCD)

A systematic way to find the gcd is **factorization**.

Let $a = p_1^{a_1} p_2^{a_2} \dots p_n^{a_n}$ and $b = p_1^{b_1} p_2^{b_2} \dots p_n^{b_n}$. Then,

$$\gcd(a, b) = p^{\min(a_1, b_1)} p^{\min(a_2, b_2)} \dots p^{\min(a_n, b_n)}$$

Least Common Multiple (LCM)

Let a and b be positive integers. The **least common multiple** of a and b is the **smallest positive integer** that is divisible by both a and b , denoted by $\text{lcm}(a, b)$.

Least Common Multiple (LCM)

Let a and b be positive integers. The **least common multiple** of a and b is the **smallest positive integer** that is divisible by both a and b , denoted by $\text{lcm}(a, b)$.

We can also use **factorization** to find the lcm.

Let $a = p_1^{a_1} p_2^{a_2} \dots p_n^{a_n}$ and $b = p_1^{b_1} p_2^{b_2} \dots p_n^{b_n}$. Then,

$$\text{lcm}(a, b) = p^{\max(a_1, b_1)} p^{\max(a_2, b_2)} \dots p^{\max(a_n, b_n)}.$$

Euclidean Algorithm

Computing the **greatest common divisor** of two integers directly from the prime factorizations can be **time consuming** since we need to find all factors of the two integers.

Luckily, we have an efficient algorithm, called **Euclidean algorithm**. This algorithm has been known since ancient times and named after the ancient Greek mathematician Euclid.

Euclidean Algorithm

For two integers 287 and 91, we want to find $\gcd(287, 91)$.

Euclidean Algorithm

For two integers 287 and 91, we want to find $\gcd(287, 91)$.

$$\text{Step 1: } 287 = 91 \cdot 3 + 14$$

Euclidean Algorithm

For two integers 287 and 91, we want to find $\gcd(287, 91)$.

$$\text{Step 1: } 287 = 91 \cdot 3 + 14$$

$$\text{Step 2: } 91 = 14 \cdot 6 + 7$$

Euclidean Algorithm

For two integers 287 and 91, we want to find $\gcd(287, 91)$.

$$\text{Step 1: } 287 = 91 \cdot 3 + 14$$

$$\text{Step 2: } 91 = 14 \cdot 6 + 7$$

$$\text{Step 3: } 14 = 7 \cdot 2 + 0$$

Euclidean Algorithm

For two integers 287 and 91, we want to find $\gcd(287, 91)$.

$$\text{Step 1: } 287 = 91 \cdot 3 + 14$$

$$\text{Step 2: } 91 = 14 \cdot 6 + 7$$

$$\text{Step 3: } 14 = 7 \cdot 2 + 0$$

$$\gcd(287, 91) = \gcd(91, 14) = \gcd(14, 7) = 7$$

The Euclidean Algorithm in Pseudocode

ALGORITHM 1 The Euclidean Algorithm.

procedure $\text{gcd}(a, b$: positive integers)

$x := a$

$y := b$

while $y \neq 0$

$r := x \bmod y$

$x := y$

$y := r$

return $x \{ \text{gcd}(a, b) \text{ is } x \}$

Step 1: $287 = 91 \cdot 3 + 14$

Step 2: $91 = 14 \cdot 6 + 7$

Step 3: $14 = 7 \cdot 2 + 0$

The Euclidean Algorithm in Pseudocode

ALGORITHM 1 The Euclidean Algorithm.

procedure $\text{gcd}(a, b)$: positive integers)

$x := a$

$y := b$

while $y \neq 0$

$r := x \bmod y$

$x := y$

$y := r$

return x { $\text{gcd}(a, b)$ is x }

$$\text{Step 1: } 287 = 91 \cdot 3 + 14$$

$$\text{Step 2: } 91 = 14 \cdot 6 + 7$$

$$\text{Step 3: } 14 = 7 \cdot 2 + 0$$

The number of divisions required to find $\text{gcd}(a, b)$ is $O(\log b)$, where $a \geq b$.

(This will be proven in later sections. Mathematical induction.)

Validity of Euclidean Algorithm

Lemma: Let $a = bq + r$, where a , b , q and r are integers. Then $\gcd(a, b) = \gcd(b, r)$.

Validity of Euclidean Algorithm

Lemma: Let $a = bq + r$, where a , b , q and r are integers. Then $\gcd(a, b) = \gcd(b, r)$.

Proof: We will show that the common divisors of a and b are the same as the common divisors of b and r , which leads to $\gcd(a, b) = \gcd(b, r)$.

Validity of Euclidean Algorithm

Lemma: Let $a = bq + r$, where a , b , q and r are integers. Then $\gcd(a, b) = \gcd(b, r)$.

Proof: We will show that the common divisors of a and b are the same as the common divisors of b and r , which leads to $\gcd(a, b) = \gcd(b, r)$.

- So suppose that d divides both a and b . Then it follows that d also divides $a - bq = r$. Hence, any common divisor of a and b is also a common divisor of b and r .

Validity of Euclidean Algorithm

Lemma: Let $a = bq + r$, where a , b , q and r are integers. Then $\gcd(a, b) = \gcd(b, r)$.

Proof: We will show that the common divisors of a and b are the same as the common divisors of b and r , which leads to $\gcd(a, b) = \gcd(b, r)$.

- So suppose that d divides both a and b . Then it follows that d also divides $a - bq = r$. Hence, any common divisor of a and b is also a common divisor of b and r .
- Suppose that d divides both b and r . Then d also divides $bq + r = a$. Hence, any common divisor of b and r is also a common divisor of a and b .

Validity of Euclidean Algorithm

Lemma: Let $a = bq + r$, where a , b , q and r are integers. Then $\gcd(a, b) = \gcd(b, r)$.

Proof: We will show that the common divisors of a and b are the same as the common divisors of b and r , which leads to $\gcd(a, b) = \gcd(b, r)$.

- So suppose that d divides both a and b . Then it follows that d also divides $a - bq = r$. Hence, any common divisor of a and b is also a common divisor of b and r .
- Suppose that d divides both b and r . Then d also divides $bq + r = a$. Hence, any common divisor of b and r is also a common divisor of a and b .

Hence, $\gcd(a, b) = \gcd(b, r)$.

Validity of Euclidean Algorithm

Suppose that a and b are positive integers with $a \geq b$. Let $r_0 = a$ and $r_1 = b$.

$$\begin{aligned} r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\ r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\ &\vdots \\ &\vdots \\ &\vdots \\ r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\ r_{n-1} &= r_n q_n. \end{aligned}$$

$$\gcd(a, b) = \gcd(r_0, r_1) = \dots = \gcd(r_{n-1}, r_n) = \gcd(r_n, 0) = r_n$$

GCD as Linear Combinations

$\gcd(a, b)$ can be expressed as a linear combination with integer coefficients of a and b .

Example: $\gcd(6, 14) = 2$, and $2 = (-2) \cdot 6 + 1 \cdot 14$.

GCD as Linear Combinations

$\gcd(a, b)$ can be expressed as a linear combination with integer coefficients of a and b .

Example: $\gcd(6, 14) = 2$, and $2 = (-2) \cdot 6 + 1 \cdot 14$.

Bezout'S Theorem: If a and b are positive integers, then there exist integers s and t such that

$$\gcd(a, b) = sa + tb.$$

This equation is called Bezout's identity.

GCD as Linear Combinations

We can use **extended Euclidean algorithm** to find Bezout's identity.

GCD as Linear Combinations

We can use **extended Euclidean algorithm** to find Bezout's identity.

Example: Express $\gcd(252, 198) = 18$ as a linear combination of 252 and 198.

GCD as Linear Combinations

We can use **extended Euclidean algorithm** to find Bezout's identity.

Example: Express $\gcd(252, 198) = 18$ as a linear combination of 252 and 198.

Solution: To show that $\gcd(252, 198) = 18$, the Euclidean algorithm uses these divisions:

$$252 = 1 \cdot 198 + 54$$

$$198 = 3 \cdot 54 + 36$$

$$54 = 1 \cdot 36 + 18$$

$$36 = 2 \cdot 18.$$

GCD as Linear Combinations

We can use **extended Euclidean algorithm** to find Bezout's identity.

Example: Express $\gcd(252, 198) = 18$ as a linear combination of 252 and 198.

Solution: To show that $\gcd(252, 198) = 18$, the Euclidean algorithm uses these divisions:

$$252 = 1 \cdot 198 + 54$$

$$198 = 3 \cdot 54 + 36$$

$$54 = 1 \cdot 36 + 18$$

$$36 = 2 \cdot 18.$$

Substituting the above expressions:

$$18 = 54 - 1 \cdot 36 = 54 - 1 \cdot (198 - 3 \cdot 54) = 4 \cdot 54 - 1 \cdot 198.$$

$$18 = 4 \cdot (252 - 1 \cdot 198) - 1 \cdot 198 = 4 \cdot 252 - 5 \cdot 198.$$



SUSTech

Southern University
of Science and
Technology

Corollaries of Bezout's Theorem

Lemma: If a , b , c are positive integers such that $\gcd(a, b) = 1$ and $a|bc$, then $a|c$.

Corollaries of Bezout's Theorem

Lemma: If a , b , c are positive integers such that $\gcd(a, b) = 1$ and $a|bc$, then $a|c$.

Proof: Since $\gcd(a, b) = 1$, by Bezout's Theorem there exist s and t such that $1 = sa + tb$. This yields $c = sac + tbc$.

Corollaries of Bezout's Theorem

Lemma: If a, b, c are positive integers such that $\gcd(a, b) = 1$ and $a|bc$, then $a|c$.

Proof: Since $\gcd(a, b) = 1$, by Bezout's Theorem there exist s and t such that $1 = sa + tb$. This yields $c = sac + tbc$.

Since $a|bc$, we have $a|tbc$. Then, since $a|sac$, we have $a|(sac + tbc)$, i.e., $a|c$.

Corollaries of Bezout's Theorem

Lemma: If a, b, c are positive integers such that $\gcd(a, b) = 1$ and $a|bc$, then $a|c$.

Proof: Since $\gcd(a, b) = 1$, by Bezout's Theorem there exist s and t such that $1 = sa + tb$. This yields $c = sac + tbc$.

Since $a|bc$, we have $a|tbc$. Then, since $a|sac$, we have $a|(sac + tbc)$, i.e., $a|c$.

Lemma: If p is prime and $p|a_1a_2\dots a_n$, then $p|a_i$ for some i .

Corollaries of Bezout's Theorem

Lemma: If a, b, c are positive integers such that $\gcd(a, b) = 1$ and $a|bc$, then $a|c$.

Proof: Since $\gcd(a, b) = 1$, by Bezout's Theorem there exist s and t such that $1 = sa + tb$. This yields $c = sac + tbc$.

Since $a|bc$, we have $a|tbc$. Then, since $a|sac$, we have $a|(sac + tbc)$, i.e., $a|c$.

Lemma: If p is prime and $p|a_1a_2\dots a_n$, then $p|a_i$ for some i .

(This will be proven in later sections. Mathematical induction.)

Uniqueness of Prime Factorization

Theorem: A **prime factorization** of a positive integer, where the primes are in nondecreasing order, is **unique**.

Uniqueness of Prime Factorization

Theorem: A **prime factorization** of a positive integer, where the primes are in nondecreasing order, is **unique**.

Proof (by contradiction): Suppose that the positive integer n can be written as a product of primes in two distinct ways:

$$n = p_1 p_2 \dots p_s \text{ and } n = q_1 q_2 \dots q_t$$

Uniqueness of Prime Factorization

Theorem: A **prime factorization** of a positive integer, where the primes are in nondecreasing order, is **unique**.

Proof (by contradiction): Suppose that the positive integer n can be written as a product of primes in two distinct ways:

$$n = p_1 p_2 \dots p_s \text{ and } n = q_1 q_2 \dots q_t$$

Remove all common primes from the factorizations to get

$$p_{i_1} p_{i_2} \dots p_{i_u} = q_{j_1} q_{j_2} \dots q_{j_v}$$

Uniqueness of Prime Factorization

Theorem: A prime factorization of a positive integer, where the primes are in nondecreasing order, is **unique**.

Proof (by contradiction): Suppose that the positive integer n can be written as a product of primes in two distinct ways:

$$n = p_1 p_2 \dots p_s \text{ and } n = q_1 q_2 \dots q_t$$

Remove all common primes from the factorizations to get

$$p_{i_1} p_{i_2} \dots p_{i_u} = q_{j_1} q_{j_2} \dots q_{j_v}$$

Thus, $p_{i_1} \mid q_{j_1} q_{j_2} \dots q_{j_v}$. It then follows that p_{i_1} divides q_{j_k} for some k , contradicting the assumption that p_{i_1} and q_{j_k} are distinct primes.

Dividing Congruences by an Integer

Theorem: Let m be a positive integer. Let a, b, c be integers. If $ac \equiv bc \pmod{m}$ and $\gcd(c, m) = 1$, then $a \equiv b \pmod{m}$.

Dividing Congruences by an Integer

Theorem: Let m be a positive integer. Let a, b, c be integers. If $ac \equiv bc \pmod{m}$ and $\gcd(c, m) = 1$, then $a \equiv b \pmod{m}$.

Example:

- $14 \equiv 8 \pmod{6}$, but $7 \not\equiv 4 \pmod{6}$
- $14 \equiv 8 \pmod{3}$, but $7 \not\equiv 4 \pmod{3}$

Dividing Congruences by an Integer

Theorem: Let m be a positive integer. Let a, b, c be integers. If $ac \equiv bc \pmod{m}$ and $\gcd(c, m) = 1$, then $a \equiv b \pmod{m}$.

Example:

- $14 \equiv 8 \pmod{6}$, but $7 \not\equiv 4 \pmod{6}$
- $14 \equiv 8 \pmod{3}$, but $7 \not\equiv 4 \pmod{3}$

Proof: Since $ac \equiv bc \pmod{m}$, we have $m \mid ac - bc$, i.e., $m \mid c(a - b)$. Because $\gcd(c, m) = 1$, it follows that $m \mid a - b$.

Mersenne Primes

Prime numbers of the form $2^p - 1$, where p is a prime.

Mersenne Primes

Prime numbers of the form $2^p - 1$, where p is a prime.

- $2^2 - 1 = 3$, $2^3 - 1 = 7$, $2^5 - 1 = 31$, $2^7 - 1 = 127$ are Mersenne primes.
- $2^{11} - 1 = 2047 = 23 \cdot 89$ is not a Mersenne prime.
- The largest known prime numbers are Mersenne primes.

Largest Known Prime, 49th Known Mersenne Prime Found!

January 7, 2016 — GIMPS celebrated its 20th anniversary with the discovery of the largest known prime number, $2^{74,207,281}-1$.

50th Known Mersenne Prime Found!

January 3, 2018 — Persistence pays off. Jonathan Pace, a GIMPS volunteer for over 14 years, discovered the 50th known Mersenne prime, $2^{77,232,917}-1$ on December 26, 2017. The prime number is calculated by multiplying together 77,232,917 twos, and then subtracting one. It weighs in at [23,249,425 digits](#), becoming the largest prime number known to mankind. It bests the [previous record prime](#), also discovered by GIMPS, by 910,807 digits.

51st Known Mersenne Prime Found!


December 21, 2018 — The [Great Internet Mersenne Prime Search \(GIMPS\)](#) has discovered the largest known prime number, $2^{82,589,933}-1$, having [24,862,048 digits](#). A computer volunteered by Patrick Laroche from Ocala, Florida made the find on December 7, 2018. The new prime number, also known as [M82589933](#), is calculated by multiplying together 82,589,933 twos and then subtracting one. It is more than one and a half million digits larger than the [previous record prime number](#).

Conjectures about Primes

Goldbach's Conjecture ($1 + 1$): Every even integer $n > 2$, is the sum of two primes.

Twin-prime Conjecture: There are infinitely many twin primes (i.e., pairs of primes that differ by 2).

This Lecture



Number Theory: divisibility and modular arithmetic, integer representations, primes, greatest common divisors, **linear congruences**



SUSTech

Southern University
of Science and
Technology

Linear Congruences

A congruence of the form $ax \equiv b \pmod{m}$, where m is a positive integer, a and b are integers, and x is a variable, is called a **linear congruence**.

Linear Congruences

A congruence of the form $ax \equiv b \pmod{m}$, where m is a positive integer, a and b are integers, and x is a variable, is called a **linear congruence**.

The solutions to a linear congruence $ax \equiv b \pmod{m}$ are **all integers x** that satisfy the congruence.

Linear Congruences

A congruence of the form $ax \equiv b \pmod{m}$, where m is a positive integer, a and b are integers, and x is a variable, is called a **linear congruence**.

The solutions to a linear congruence $ax \equiv b \pmod{m}$ are **all integers x** that satisfy the congruence.

Systems of linear congruences have been studied since ancient times.

今有物不知其数 三三数之剩二 五五数之剩三 七七数之剩二 问物几何

About 1500 years ago, the Chinese mathematician Sun-Tsu asked: “**There are certain things whose number is unknown. When divided by 3, the remainder is 2; when divided by 5, the remainder is 3; when divided by 7, the remainder is 2. What will be the number of things?**”

Modular Inverse

An integer \bar{a} such that $\bar{a}a \equiv 1 \pmod{m}$ is said to be an **inverse** of a modulo m .

Modular Inverse

An integer \bar{a} such that $\bar{a}a \equiv 1 \pmod{m}$ is said to be an **inverse** of a modulo m .

One method of solving linear congruences makes use of an inverse a modulo m (if it exists):

Modular Inverse

An integer \bar{a} such that $\bar{a}a \equiv 1 \pmod{m}$ is said to be an **inverse** of a modulo m .

One method of solving linear congruences makes use of an inverse a modulo m (if it exists):

From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Modular Inverse

An integer \bar{a} such that $\bar{a}a \equiv 1 \pmod{m}$ is said to be an **inverse** of a modulo m .

One method of solving linear congruences makes use of an inverse a modulo m (if it exists):

From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Note that $\bar{a}ax \bmod m = ((\bar{a}a \bmod m)(x \bmod m)) \bmod m = x \bmod m$.

Modular Inverse

An integer \bar{a} such that $\bar{a}a \equiv 1 \pmod{m}$ is said to be an **inverse** of a modulo m .

One method of solving linear congruences makes use of an inverse a modulo m (if it exists):

From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Note that $\bar{a}ax \pmod{m} = ((\bar{a}a \pmod{m})(x \pmod{m})) \pmod{m} = x \pmod{m}$.

Thus, $x \pmod{m} = \bar{a}ax \pmod{m} = \bar{a}b \pmod{m}$, which implies that

$$x \equiv \bar{a}b \pmod{m}.$$

Modular Inverse

An integer \bar{a} such that $\bar{a}a \equiv 1 \pmod{m}$ is said to be an **inverse** of a modulo m .

One method of solving linear congruences makes use of an inverse a modulo m (if it exists):

From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Note that $\bar{a}ax \pmod{m} = ((\bar{a}a \pmod{m})(x \pmod{m})) \pmod{m} = x \pmod{m}$.

Thus, $x \pmod{m} = \bar{a}ax \pmod{m} = \bar{a}b \pmod{m}$, which implies that

$$x \equiv \bar{a}b \pmod{m}.$$

When does an inverse of a modulo m exist?

Inverse of a modulo m

Theorem: If a and m are relatively prime integers and $m > 1$, then an inverse of a modulo m exists. The inverse is unique modulo m . That is,

- there is a unique positive integer \bar{a} less than m that is an inverse of a modulo m and
- every other inverse of a modulo m is congruent to \bar{a} modulo m .)

Inverse of a modulo m

Theorem: If a and m are relatively prime integers and $m > 1$, then an inverse of a modulo m exists. The inverse is unique modulo m .

Proof: Since $\gcd(a, m) = 1$, there are integers s and t such that

$$sa + tm = 1.$$

Hence $sa + tm \equiv 1 \pmod{m}$.

Inverse of a modulo m

Theorem: If a and m are relatively prime integers and $m > 1$, then an inverse of a modulo m exists. The inverse is unique modulo m .

Proof: Since $\gcd(a, m) = 1$, there are integers s and t such that

$$sa + tm = 1.$$

Hence $sa + tm \equiv 1 \pmod{m}$. Since $tm \equiv 0 \pmod{m}$, it follows that $sa \equiv 1 \pmod{m}$. This means that s is an inverse of a modulo m .

Inverse of a modulo m

Theorem: If a and m are relatively prime integers and $m > 1$, then an inverse of a modulo m exists. The inverse is unique modulo m .

Proof: Since $\gcd(a, m) = 1$, there are integers s and t such that

$$sa + tm = 1.$$

Hence $sa + tm \equiv 1 \pmod{m}$. Since $tm \equiv 0 \pmod{m}$, it follows that $sa \equiv 1 \pmod{m}$. This means that s is an inverse of a modulo m .

How to prove the uniqueness of the inverse?

Inverse of a modulo m

Theorem: If a and m are relatively prime integers and $m > 1$, then an inverse of a modulo m exists. The inverse is unique modulo m .

Proof: Since $\gcd(a, m) = 1$, there are integers s and t such that

$$sa + tm = 1.$$

Hence $sa + tm \equiv 1 \pmod{m}$. Since $tm \equiv 0 \pmod{m}$, it follows that $sa \equiv 1 \pmod{m}$. This means that s is an inverse of a modulo m .

How to prove the uniqueness of the inverse?

Suppose that b and c are both inverses of a modulo m . Then $ba \equiv 1 \pmod{m}$ and $ca \equiv 1 \pmod{m}$. Hence, $ba \equiv ca \pmod{m}$. Because $\gcd(a, m) = 1$ it follows that $b \equiv c \pmod{m}$.

How to find inverses?

Using extended Euclidean algorithm:

How to find inverses?

Using **extended Euclidean algorithm**:

Example: Find an inverse of 101 modulo 4620. That is, find \bar{a} such that $\bar{a} \cdot 101 \equiv 1 \pmod{4620}$.

How to find inverses?

Using **extended Euclidean algorithm**:

Example: Find an inverse of 101 modulo 4620. That is, find \bar{a} such that $\bar{a} \cdot 101 \equiv 1 \pmod{4620}$.

$$4620 = 45 \cdot 101 + 75$$

$$101 = 1 \cdot 75 + 26$$

$$75 = 2 \cdot 26 + 23$$

$$26 = 1 \cdot 23 + 3$$

$$23 = 7 \cdot 3 + 2$$

$$3 = 1 \cdot 2 + 1$$

$$2 = 2 \cdot 1$$

How to find inverses?

Using **extended Euclidean algorithm**:

Example: Find an inverse of 101 modulo 4620. That is, find \bar{a} such that $\bar{a} \cdot 101 \equiv 1 \pmod{4620}$.

	$1 = 3 - 1 \cdot 2$
$4620 = 45 \cdot 101 + 75$	$1 = 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$
$101 = 1 \cdot 75 + 26$	$1 = -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23$
$75 = 2 \cdot 26 + 23$	$1 = 8 \cdot 26 - 9 \cdot (75 - 2 \cdot 26) = 26 \cdot 26 - 9 \cdot 75$
$26 = 1 \cdot 23 + 3$	$1 = 26 \cdot (101 - 1 \cdot 75) - 9 \cdot 75$
$23 = 7 \cdot 3 + 2$	$\quad = 26 \cdot 101 - 35 \cdot 75$
$3 = 1 \cdot 2 + 1$	$1 = 26 \cdot 101 - 35 \cdot (4620 - 45 \cdot 101)$
$2 = 2 \cdot 1$	$\quad = -35 \cdot 4620 + 1601 \cdot 101$

How to find inverses?

Using **extended Euclidean algorithm**:

Example: Find an inverse of 101 modulo 4620. That is, find \bar{a} such that $\bar{a} \cdot 101 \equiv 1 \pmod{4620}$.

	$1 = 3 - 1 \cdot 2$
$4620 = 45 \cdot 101 + 75$	$1 = 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3$
$101 = 1 \cdot 75 + 26$	$1 = -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23$
$75 = 2 \cdot 26 + 23$	$1 = 8 \cdot 26 - 9 \cdot (75 - 2 \cdot 26) = 26 \cdot 26 - 9 \cdot 75$
$26 = 1 \cdot 23 + 3$	$1 = 26 \cdot (101 - 1 \cdot 75) - 9 \cdot 75$
$23 = 7 \cdot 3 + 2$	$\quad = 26 \cdot 101 - 35 \cdot 75$
$3 = 1 \cdot 2 + 1$	$1 = 26 \cdot 101 - 35 \cdot (4620 - 45 \cdot 101)$
$2 = 2 \cdot 1$	$\quad = -35 \cdot 4620 + 1601 \cdot 101$

That $-35 \cdot 4620 + 1601 \cdot 101 = 1$ tells us that -35 and 1601 are Bezout coefficients of 4620 and 101. We have

$$1 \bmod 4620 = 1601 \cdot 101 \bmod 4620.$$

Thus, 1601 is an inverse of 101 modulo 4620.

Using Inverses to Solve Congruences

Solve the congruence $ax \equiv b \pmod{m}$ by multiplying both sides by \bar{a} .

Using Inverses to Solve Congruences

Solve the congruence $ax \equiv b \pmod{m}$ by multiplying both sides by \bar{a} .

Recall: From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Note that $\bar{a}ax \bmod m = ((\bar{a}a \bmod m)(x \bmod m)) \bmod m = x \bmod m$.

Thus, $x \bmod m = \bar{a}ax \bmod m = \bar{a}b \bmod m$, which implies that

$$x \equiv \bar{a}b \pmod{m}.$$

Using Inverses to Solve Congruences

Solve the congruence $ax \equiv b \pmod{m}$ by multiplying both sides by \bar{a} .

Recall: From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Note that $\bar{a}ax \bmod m = ((\bar{a}a \bmod m)(x \bmod m)) \bmod m = x \bmod m$.

Thus, $x \bmod m = \bar{a}ax \bmod m = \bar{a}b \bmod m$, which implies that

$$x \equiv \bar{a}b \pmod{m}.$$

Example: What are the solutions of the congruence $3x \equiv 4 \pmod{7}$?

Using Inverses to Solve Congruences

Solve the congruence $ax \equiv b \pmod{m}$ by multiplying both sides by \bar{a} .

Recall: From $ax \equiv b \pmod{m}$, it follows that $\bar{a}ax \equiv \bar{a}b \pmod{m}$.

Note that $\bar{a}ax \bmod m = ((\bar{a}a \bmod m)(x \bmod m)) \bmod m = x \bmod m$.

Thus, $x \bmod m = \bar{a}ax \bmod m = \bar{a}b \bmod m$, which implies that

$$x \equiv \bar{a}b \pmod{m}.$$

Example: What are the solutions of the congruence $3x \equiv 4 \pmod{7}$?

Solution: We found that -2 is an inverse of 3 modulo 7. Multiply both sides of the congruence by -2 . Since $-8 \equiv 6 \pmod{7}$, we have $x \equiv 6 \pmod{7}$, namely, 6, 13, 20, . . . and -1, -8, ...

Number of Solutions to Congruences

The previous approach (based on the inverse of a modulo m) works for only the scenario with $\gcd(a, m) = 1$.

Number of Solutions to Congruences

Theorem*: Let $\gcd(a, m) = d$. Let $m' = m/d$ and $a' = a/d$. The congruence $ax \equiv b \pmod{m}$ has solutions **if and only if** $d \mid b$.

- If $d \mid b$, then there are exactly d solutions, where by “solution” we mean a congruence class mod m
- If x_0 is a solution, then the other solutions are given by $x_0 + m', x_0 + 2m', \dots, x_0 + (d - 1)m'$.

Number of Solutions to Congruences

Theorem*: Let $\gcd(a, m) = d$. Let $m' = m/d$ and $a' = a/d$. The congruence $ax \equiv b \pmod{m}$ has solutions **if and only if** $d \mid b$.

- If $d \mid b$, then there are exactly d solutions, where by “solution” we mean a congruence class mod m
- If x_0 is a solution, then the other solutions are given by $x_0 + m', x_0 + 2m', \dots, x_0 + (d - 1)m'$.

Proof:

“only if”: Let x_0 be a solution, then $ax_0 - b = km$. Thus, $ax_0 - km = b$. Since $d \mid ax_0 - km$, we must have $d \mid b$.

Number of Solutions to Congruences

Theorem*: Let $\gcd(a, m) = d$. Let $m' = m/d$ and $a' = a/d$. The congruence $ax \equiv b \pmod{m}$ has solutions **if and only if** $d \mid b$.

- If $d \mid b$, then there are exactly d solutions, where by “solution” we mean a congruence class mod m
- If x_0 is a solution, then the other solutions are given by $x_0 + m', x_0 + 2m', \dots, x_0 + (d - 1)m'$.

Proof:

“only if”: Let x_0 be a solution, then $ax_0 - b = km$. Thus, $ax_0 - km = b$. Since $d \mid ax_0 - km$, we must have $d \mid b$.

“if”: Suppose that $d \mid b$. Let $b = kd$. Since $\gcd(a, m) = d$, there exist integers s and t such that $d = as + mt$. Multiplying both sides by k . Then, $b = ask + mtk$. Let $x_0 = sk$. Then $ax_0 \equiv b \pmod{m}$.

Number of Solutions to Congruences

Theorem*: Let $\gcd(a, m) = d$. Let $m' = m/d$ and $a' = a/d$. The congruence $ax \equiv b \pmod{m}$ has solutions **if and only if** $d \mid b$.

- If $d \mid b$, then there are exactly d “solutions”, where by “solution” we mean a congruence class mod m .
- If x_0 is a solution, then the other solutions are given by $x_0 + m', x_0 + 2m', \dots, x_0 + (d - 1)m'$.

Proof:

“The number of solutions is d ”: Consider two solutions x_0 and x_1 . $ax_0 \equiv b \pmod{m}$ and $ax_1 \equiv b \pmod{m}$ imply that $m \mid a(x_1 - x_0)$ and $m' \mid a'(x_1 - x_0)$. This implies further that $x_1 = x_0 + km'$.

Number of Solutions to Congruences

Theorem*: Let $\gcd(a, m) = d$. Let $m' = m/d$ and $a' = a/d$. The congruence $ax \equiv b \pmod{m}$ has solutions **if and only if** $d \mid b$.

- If $d \mid b$, then there are exactly d “solutions”, where by “solution” we mean a congruence class mod m .
- If x_0 is a solution, then the other solutions are given by $x_0 + m', x_0 + 2m', \dots, x_0 + (d - 1)m'$.

Proof:

“The number of solutions is d ”: Consider two solutions x_0 and x_1 . $ax_0 \equiv b \pmod{m}$ and $ax_1 \equiv b \pmod{m}$ imply that $m \mid a(x_1 - x_0)$ and $m' \mid a'(x_1 - x_0)$. This implies further that $x_1 = x_0 + km'$.

To finish the proof, observe that as k runs through the values $0, 1, \dots, d - 1$ (the residues mod d), the congruence classes $[x_0 + (m/d)k]_m$ run through all the solutions.

The Chinese Remainder Theorem

Systems of linear congruences have been studied since ancient times.

今有物不知其数 三三数之剩二 五五数之剩三 七七数之剩二 问物几何

About 1500 years ago, the Chinese mathematician Sun-Tsu asked: “There are certain things whose number is unknown. When divided by 3, the remainder is 2; when divided by 5, the remainder is 3; when divided by 7, the remainder is 2. What will be the number of things?”

- $x \equiv 2 \pmod{3}$
- $x \equiv 3 \pmod{5}$
- $x \equiv 2 \pmod{7}$

The Chinese Remainder Theorem

Theorem (The Chinese Remainder Theorem): Let m_1, m_2, \dots, m_n be pairwise relatively prime positive integers greater than 1 and a_1, a_2, \dots, a_n arbitrary integers. Then, the system

$$x \equiv a_1 \pmod{m_1}$$

$$x \equiv a_2 \pmod{m_2}$$

...

$$x \equiv a_n \pmod{m_n}$$

has a **unique solution** modulo $m = m_1 m_2 \dots m_n$.

(That is, there is a solution x with $0 \leq x < m$, and all other solutions are congruent modulo m to this solution.)

The Chinese Remainder Theorem

Proof: To show such a solution exists: Let $M_k = m/m_k$ for $k = 1, 2, \dots, n$ and $m = m_1 m_2 \dots m_n$. Thus, $M_k = m_1 \dots m_{k-1} m_{k+1} \dots m_n$.

The Chinese Remainder Theorem

Proof: To show such a solution exists: Let $M_k = m/m_k$ for $k = 1, 2, \dots, n$ and $m = m_1 m_2 \dots m_n$. Thus, $M_k = m_1 \dots m_{k-1} m_{k+1} \dots m_n$.

Since $\gcd(m_k, M_k) = 1$, there is an integer y_k , an inverse of M_k modulo m_k , such that $M_k y_k \equiv 1 \pmod{m_k}$. Let

$$x = a_1 M_1 y_1 + a_2 M_2 y_2 + \dots + a_n M_n y_n.$$

It is checked that x is a solution to the n congruences:

The Chinese Remainder Theorem

Proof: To show such a solution exists: Let $M_k = m/m_k$ for $k = 1, 2, \dots, n$ and $m = m_1 m_2 \dots m_n$. Thus, $M_k = m_1 \dots m_{k-1} m_{k+1} \dots m_n$.

Since $\gcd(m_k, M_k) = 1$, there is an integer y_k , an inverse of M_k modulo m_k , such that $M_k y_k \equiv 1 \pmod{m_k}$. Let

$$x = a_1 M_1 y_1 + a_2 M_2 y_2 + \dots + a_n M_n y_n.$$

It is checked that x is a solution to the n congruences:

$$x \bmod m_k = (a_1 M_1 y_1 + a_2 M_2 y_2 + \dots + a_n M_n y_n) \bmod m_k$$

Since $M_k = m/m_k$, we have $x \bmod m_k = a_k M_k y_k \bmod m_k$. Since $M_k y_k \equiv 1 \pmod{m_k}$, we have $a_k M_k y_k \bmod m_k = a_k \bmod m_k$. Thus,

$$x \equiv a_k \pmod{m_k}.$$

The Chinese Remainder Theorem

How to prove the **uniqueness** of the solution modulo m ?

Proof: Suppose that x and x' are both solutions to all the congruences. As x and x' give the same remainder, when divided by m_k , their difference $x - x'$ is a multiple of each m_k for all $k = 1, 2, \dots, n$.

The Chinese Remainder Theorem

How to prove the **uniqueness** of the solution modulo m ?

Proof: Suppose that x and x' are both solutions to all the congruences. As x and x' give the same remainder, when divided by m_k , their difference $x - x'$ is a multiple of each m_k for all $k = 1, 2, \dots, n$.

As m_1, m_2, \dots, m_n be pairwise relatively prime positive integers, their product m divides $x - x'$, and thus x and x' are congruent modulo m , i.e., $x \equiv x' \pmod{m}$.

The Chinese Remainder Theorem

How to prove the **uniqueness** of the solution modulo m ?

Proof: Suppose that x and x' are both solutions to all the congruences. As x and x' give the same remainder, when divided by m_k , their difference $x - x'$ is a multiple of each m_k for all $k = 1, 2, \dots, n$.

As m_1, m_2, \dots, m_n be pairwise relatively prime positive integers, their product m divides $x - x'$, and thus x and x' are congruent modulo m , i.e., $x \equiv x' \pmod{m}$.

This implies that given a solution x with $0 \leq x < m$, all other solutions are congruent modulo m to this solution.

The Chinese Remainder Theorem: Example

$$x \equiv 2 \pmod{3}$$

$$x \equiv 3 \pmod{5}$$

$$x \equiv 2 \pmod{7}$$

The Chinese Remainder Theorem: Example

$$x \equiv 2 \pmod{3}$$

$$x \equiv 3 \pmod{5}$$

$$x \equiv 2 \pmod{7}$$

- ① Let $m = 3 \cdot 5 \cdot 7 = 105$, $M_1 = m/3 = 35$, $M_2 = m/5 = 21$, and $M_3 = m/7 = 15$.

The Chinese Remainder Theorem: Example

$$x \equiv 2 \pmod{3}$$

$$x \equiv 3 \pmod{5}$$

$$x \equiv 2 \pmod{7}$$

- ① Let $m = 3 \cdot 5 \cdot 7 = 105$, $M_1 = m/3 = 35$, $M_2 = m/5 = 21$, and $M_3 = m/7 = 15$.
- ② Compute the inverse of M_k modulo m_k :
 - ▶ $35 \cdot 2 \equiv 1 \pmod{3}$ $y_1 = 2$
 - ▶ $21 \equiv 1 \pmod{5}$ $y_2 = 1$
 - ▶ $15 \equiv 1 \pmod{7}$ $y_3 = 1$

The Chinese Remainder Theorem: Example

$$x \equiv 2 \pmod{3}$$

$$x \equiv 3 \pmod{5}$$

$$x \equiv 2 \pmod{7}$$

- ① Let $m = 3 \cdot 5 \cdot 7 = 105$, $M_1 = m/3 = 35$, $M_2 = m/5 = 21$, and $M_3 = m/7 = 15$.
- ② Compute the inverse of M_k modulo m_k :
 - ▶ $35 \cdot 2 \equiv 1 \pmod{3}$ $y_1 = 2$
 - ▶ $21 \equiv 1 \pmod{5}$ $y_2 = 1$
 - ▶ $15 \equiv 1 \pmod{7}$ $y_3 = 1$
- ③ Compute a solution x :
$$x = 2 \cdot 35 \cdot 2 + 3 \cdot 21 \cdot 1 + 2 \cdot 15 \cdot 1 \equiv 233 \equiv 23 \pmod{105}$$

The Chinese Remainder Theorem: Example

$$x \equiv 2 \pmod{3}$$

$$x \equiv 3 \pmod{5}$$

$$x \equiv 2 \pmod{7}$$

- ① Let $m = 3 \cdot 5 \cdot 7 = 105$, $M_1 = m/3 = 35$, $M_2 = m/5 = 21$, and $M_3 = m/7 = 15$.
- ② Compute the inverse of M_k modulo m_k :
 - ▶ $35 \cdot 2 \equiv 1 \pmod{3}$ $y_1 = 2$
 - ▶ $21 \equiv 1 \pmod{5}$ $y_2 = 1$
 - ▶ $15 \equiv 1 \pmod{7}$ $y_3 = 1$
- ③ Compute a solution x :
$$x = 2 \cdot 35 \cdot 2 + 3 \cdot 21 \cdot 1 + 2 \cdot 15 \cdot 1 \equiv 233 \equiv 23 \pmod{105}$$
- ④ The solutions are all integers x that satisfy $x \equiv 23 \pmod{105}$.



SUSTech

Southern University
of Science and
Technology

Back Substitution

We may also solve systems of linear congruences with pairwise relatively prime moduli m_1, m_2, \dots, m_n by back substitution.

Back Substitution

We may also solve systems of linear congruences with pairwise relatively prime moduli m_1, m_2, \dots, m_n by back substitution.

Example:

$$(1) \quad x \equiv 1 \pmod{5}$$

$$(2) \quad x \equiv 2 \pmod{6}$$

$$(3) \quad x \equiv 3 \pmod{7}$$

Back Substitution

We may also solve systems of linear congruences with pairwise relatively prime moduli m_1, m_2, \dots, m_n by back substitution.

Example:

$$(1) \ x \equiv 1 \pmod{5}$$

$$(2) \ x \equiv 2 \pmod{6}$$

$$(3) \ x \equiv 3 \pmod{7}$$

According to (1), $x = 5t + 1$, where t is an integer.

Back Substitution

We may also solve systems of linear congruences with pairwise relatively prime moduli m_1, m_2, \dots, m_n by back substitution.

Example:

$$(1) \quad x \equiv 1 \pmod{5}$$

$$(2) \quad x \equiv 2 \pmod{6}$$

$$(3) \quad x \equiv 3 \pmod{7}$$

According to (1), $x = 5t + 1$, where t is an integer.

Substituting this expression into (2), we have $5t + 1 \equiv 2 \pmod{6}$, which means that $t \equiv 5 \pmod{6}$. Thus, $t = 6u + 5$, where u is an integer.

Back Substitution

We may also solve systems of linear congruences with pairwise relatively prime moduli m_1, m_2, \dots, m_n by back substitution.

Example:

$$(1) \ x \equiv 1 \pmod{5}$$

$$(2) \ x \equiv 2 \pmod{6}$$

$$(3) \ x \equiv 3 \pmod{7}$$

According to (1), $x = 5t + 1$, where t is an integer.

Substituting this expression into (2), we have $5t + 1 \equiv 2 \pmod{6}$, which means that $t \equiv 5 \pmod{6}$. Thus, $t = 6u + 5$, where u is an integer.

Substituting $x = 5t + 1$ and $t = 6u + 5$ into (3), we have $30u + 26 \equiv 3 \pmod{7}$, which implies that $u \equiv 6 \pmod{7}$. Thus, $u = 7v + 6$, where v is an integer.

Back Substitution

We may also solve systems of linear congruences with pairwise relatively prime moduli m_1, m_2, \dots, m_n by back substitution.

Example:

$$(1) \ x \equiv 1 \pmod{5}$$

$$(2) \ x \equiv 2 \pmod{6}$$

$$(3) \ x \equiv 3 \pmod{7}$$

According to (1), $x = 5t + 1$, where t is an integer.

Substituting this expression into (2), we have $5t + 1 \equiv 2 \pmod{6}$, which means that $t \equiv 5 \pmod{6}$. Thus, $t = 6u + 5$, where u is an integer.

Substituting $x = 5t + 1$ and $t = 6u + 5$ into (3), we have $30u + 26 \equiv 3 \pmod{7}$, which implies that $u \equiv 6 \pmod{7}$. Thus, $u = 7v + 6$, where v is an integer.

Thus, we must have $x = 210v + 206$. Translating this back into a congruence,

$$x \equiv 206 \pmod{210}.$$

Modular Arithmetic in CS

Modular arithmetic and congruencies are used in CS:

- Pseudorandom number generators
- Hash functions
- Cryptography

Next Lecture

