



# CSE5014 CRYPTOGRAPHY AND NETWORK SECURITY

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# Private-key schemes

- We have seen how to construct schemes based on various lower-level primitives
  - Stream ciphers / PRGs
  - Block ciphers / PRFs
  - Hash functions
- How do we construct these primitives?



# Hash functions

- Main goal is *collision resistance*
  - Want *optimal* birthday security

# Hash functions

- Main goal is *collision resistance*
  - Want *optimal* birthday security
- Also want *preimage resistance*, *2nd-preimage resistance*
  - Want *optimal* security here as well
- “*Optimal*” measured relative to a *random* function
  - Why not design  $H$  to be a “*random function*”?



# The random-oracle (RO) model

- Treat  $H$  as a public, *random* function
- Then  $H(x)$  is *uniform* for any  $x$ 
  - Unless the attacker computes  $H(x)$  explicitly

# The random-oracle (RO) model

- Treat  $H$  as a public, *random* function
- Then  $H(x)$  is *uniform* for any  $x$ 
  - Unless the attacker computes  $H(x)$  explicitly
- Intuitively
  - Assume the hash function “*is random*”
  - Models attacks that are *agnostic* to the specific hash function being used
  - Security in the real world as long as “*no weaknesses found*” in the hash function



# The random-oracle (RO) model

- Formally
  - Choose a *uniform* hash function as part of the security experiment
  - Attacker can *only* evaluate  $H$  via *explicit* queries to an oracle
  - Simulate  $H$  for the attacker as part of the security proof/reduction



# The random-oracle (RO) model

- Formally
  - Choose a *uniform* hash function as part of the security experiment
  - Attacker can *only* evaluate  $H$  via *explicit* queries to an oracle
  - Simulate  $H$  for the attacker as part of the security proof/reduction
- In practice
  - Prove security in the RO model
  - Instantiate the RO with a “*good*” hash function
  - Hope for the best





# Pros and cons of the RO model

## ■ Cons

- There is **no** such a thing as a public hash function that “*is random*”
  - Not even clear what this means formally
- Known counterexamples
  - There are (contrived) schemes secure in the RO model, but insecure when using any real-world hash function
- Sometimes ***over-abused*** (arguably)

# Pros and cons of the RO model

## ■ Pros

- No known example of “*natural*” scheme secure in the RO model being attacked in the real world
- If an attack is found, just replace the hash
- Proof in the RO model better than no proof at all
  - Evidence that the basic design principles are sound

# Groups

- A *group* is a set  $G$  and a binary operation  $\circ$  defined on  $G$  such that:
  - (*Closure*) For all  $g, h \in G$ ,  $g \circ h$  is in  $G$
  - (*Identity*) There is a **unique** element  $e \in G$  such that  $e \circ g = g$  for all  $g \in G$
  - (*Inverse*) Every element  $g \in G$  has an *inverse*  $h \in G$  such that  $g \circ h = e$
  - (*Associativity*) For all  $f, g, h \in G$ ,  $f \circ (g \circ h) = (f \circ g) \circ h$



# Groups

- A *group* is a set  $G$  and a binary operation  $\circ$  defined on  $G$  such that:
  - (*Closure*) For all  $g, h \in G$ ,  $g \circ h$  is in  $G$
  - (*Identity*) There is a **unique** element  $e \in G$  such that  $e \circ g = g$  for all  $g \in G$
  - (*Inverse*) Every element  $g \in G$  has an *inverse*  $h \in G$  such that  $g \circ h = e$
  - (*Associativity*) For all  $f, g, h \in G$ ,  $f \circ (g \circ h) = (f \circ g) \circ h$
- A *group* is called *abelian* if the further property holds.
  - (*Commutativity*) For all  $g, h \in G$ ,  $g \circ h = h \circ g$



# Groups

- A *group* is a set  $G$  and a binary operation  $\circ$  defined on  $G$  such that:
  - (*Closure*) For all  $g, h \in G$ ,  $g \circ h$  is in  $G$
  - (*Identity*) There is a **unique** element  $e \in G$  such that  $e \circ g = g$  for all  $g \in G$
  - (*Inverse*) Every element  $g \in G$  has an *inverse*  $h \in G$  such that  $g \circ h = e$
  - (*Associativity*) For all  $f, g, h \in G$ ,  $f \circ (g \circ h) = (f \circ g) \circ h$
- A *group* is called *abelian* if the further property holds.
  - (*Commutativity*) For all  $g, h \in G$ ,  $g \circ h = h \circ g$
- The *order* of a **finite** group  $G$  is  $\#$  of elements in  $G$ .



# Examples

- $\mathbb{Z}$  under addition
- $\mathbb{Z} \setminus \{0\}$  under multiplication
- $\mathbb{Q}$  under addition
- $\mathbb{Q} \setminus \{0\}$  under multiplication
- $\mathbb{R}$  under addition
- $\mathbb{R} \setminus \{0\}$  under multiplication
- $\{0, 1\}^*$  under concatenation
- $\{0, 1\}^n$  under bitwise XOR
- $2 \times 2$  real matrices under addition
- $2 \times 2$  invertible, real matrices under multiplication

# Groups

- The group operation can be written *additively* or *multiplicatively*
  - I.e., instead of  $g \circ h$ , write  $g + h$  or  $gh$
  - Does *not* mean that the group operation corresponds to (integer) addition or multiplication



# Groups

- The group operation can be written *additively* or *multiplicatively*
  - I.e., instead of  $g \circ h$ , write  $g + h$  or  $gh$
  - Does *not* mean that the group operation corresponds to (integer) addition or multiplication
- Identity denoted by 0 or 1, respectively
- Inverse of  $g$  denoted by  $-g$  or  $g^{-1}$ , respectively
- Group *exponentiation*:  $m \cdot a$  or  $a^m$ , respectively





# Useful example

- $\mathbb{Z}_N = \{0, 1, \dots, N - 1\}$  under addition modulo  $n$ 
  - *Identity* is 0
  - *Inverse* of  $a$  is  $-a \bmod N$
  - *Associativity*, *commutativity* obvious
  - *Order*  $N$



# Useful example

- $\mathbb{Z}_N = \{0, 1, \dots, N - 1\}$  under addition modulo  $n$ 
  - *Identity* is 0
  - *Inverse* of  $a$  is  $-a \bmod N$
  - *Associativity*, *commutativity* obvious
  - *Order*  $N$
- What happens if we consider *multiplication* modulo  $N$ ?



# Useful example

- $\mathbb{Z}_N = \{0, 1, \dots, N - 1\}$  under addition modulo  $n$ 
  - *Identity* is 0
  - *Inverse* of  $a$  is  $-a \bmod N$
  - *Associativity*, *commutativity* obvious
  - *Order*  $N$
- What happens if we consider *multiplication* modulo  $N$ ?
- $\mathbb{Z}_N$  is *not* a group under this operation!
  - 0 has *no* inverse
  - Even if we exclude 0, there is, e.g., *no* inverse of 2 modulo 4



# Example

- What happens if we consider *multiplication* modulo  $N$ ?
- Consider instead the *invertible* elements modulo  $N$ , under multiplication modulo  $N$



# Example

- What happens if we consider *multiplication* modulo  $N$ ?
- Consider instead the *invertible* elements modulo  $N$ , under multiplication modulo  $N$
- $\mathbb{Z}_N^* = \{0 < x < N : \gcd(x, N) = 1\}$ 
  - *Closure*
  - *Identity* is 1
  - *Inverse* of  $a$  is  $a^{-1} \bmod N$
  - *Associativity*, *commutativity* obvious



# Example

- What happens if we consider *multiplication* modulo  $N$ ?
- Consider instead the *invertible* elements modulo  $N$ , under multiplication modulo  $N$
- $\mathbb{Z}_N^* = \{0 < x < N : \gcd(x, N) = 1\}$ 
  - *Closure*
  - *Identity* is 1
  - *Inverse* of  $a$  is  $a^{-1} \bmod N$
  - *Associativity*, *commutativity* obvious
- If  $p$  is prime, then  $\mathbb{Z}_p^* = \{1, 2, \dots, p-1\}$ 
  - $\mathbb{Z}_p$  is a (prime) *field*



# Permutation Group

- Let  $s_n = \langle 1, 2, \dots, n \rangle$  denote a *sequence* of integers 1 through  $n$ . Denote by  $P_n$  the set of all *permutations* of the sequence  $s_n$ .



# Permutation Group

- Let  $s_n = \langle 1, 2, \dots, n \rangle$  denote a *sequence* of integers 1 through  $n$ . Denote by  $P_n$  the set of all *permutations* of the sequence  $s_n$ .

For example,  $s_3 = \langle 1, 2, 3 \rangle$

$$P_3 = \{ \langle 1, 2, 3 \rangle, \langle 1, 3, 2 \rangle, \langle 2, 1, 3 \rangle, \langle 2, 3, 1 \rangle, \langle 3, 1, 2 \rangle, \langle 3, 2, 1 \rangle \}$$





# Permutation Group

- Let  $s_n = \langle 1, 2, \dots, n \rangle$  denote a *sequence* of integers 1 through  $n$ . Denote by  $P_n$  the set of all *permutations* of the sequence  $s_n$ .

For example,  $s_3 = \langle 1, 2, 3 \rangle$

$$P_3 = \{ \langle 1, 2, 3 \rangle, \langle 1, 3, 2 \rangle, \langle 2, 1, 3 \rangle, \langle 2, 3, 1 \rangle, \langle 3, 1, 2 \rangle, \langle 3, 2, 1 \rangle \}$$

- Define a binary operation  $\circ$  on the elements of  $P_n$ :  
for  $\rho, \pi \in P_n$ ,  $\pi \circ \rho$  denotes a *re-permutation* of the elements of  $\rho$  according to the elements of  $\pi$ .



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.  
$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$
$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each  $\rho \in P_3$ , there exists another unique  $\pi \in P_3$  such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.  
$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$
$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each  $\rho \in P_3$ , there exists another unique  $\pi \in P_3$  such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$

$(P_n, \circ)$  is called a *permutation group*.



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.  
$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$
$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each  $\rho \in P_3$ , there exists another unique  $\pi \in P_3$  such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$

$(P_n, \circ)$  is called a *permutation group*.  
 $(P_n, \circ)$  is **not abelian**.



# Ring

- If  $(R, +)$  is an *abelian group*, we define one more operation (denoted as *multiplication*  $\times$  for convenience) to have a *ring*  $(R, +, \times)$  satisfying the following properties.





# Ring

- If  $(R, +)$  is an *abelian group*, we define one more operation (denoted as *multiplication*  $\times$  for convenience) to have a *ring*  $(R, +, \times)$  satisfying the following properties.

**Closure:**  $R$  must be closed w.r.t.  $\times$

**Associativity:**  $(a \times b) \times c = a \times (b \times c)$

**Distributivity:**  $a \times (b + c) = a \times b + a \times c$   
 $(a + b) \times c = a \times c + b \times c$



# Ring

- If  $(R, +)$  is an *abelian group*, we define one more operation (denoted as *multiplication*  $\times$  for convenience) to have a *ring*  $(R, +, \times)$  satisfying the following properties.

**Closure:**  $R$  must be closed w.r.t.  $\times$

**Associativity:**  $(a \times b) \times c = a \times (b \times c)$

**Distributivity:**  $a \times (b + c) = a \times b + a \times c$   
 $(a + b) \times c = a \times c + b \times c$

**Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$ ,  $(\mathbb{M}_{n \times n}, +, \cdot)$  ?



# Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ( $ab = ba$ )



# Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ( $ab = ba$ )
- An *integral domain*  $(R, +, \times)$  is a *commutative ring* that satisfies the following two additional properties.

**Identity element** for multiplication:  $a1 = 1a = a$

**Nonzero product** for any two nonzero elements:  
if  $ab = 0$ , then either  $a$  or  $b$  **must be** 0.



# Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ( $ab = ba$ )
- An *integral domain*  $(R, +, \times)$  is a *commutative ring* that satisfies the following two additional properties.

**Identity element** for multiplication:  $a1 = 1a = a$

**Nonzero product** for any two nonzero elements:

if  $ab = 0$ , then either  $a$  or  $b$  **must be** 0.

**Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$  ?

$(\mathbb{Z}_m, +, \times)$ ,  $(M_{n \times n}, +, \cdot)$  ?



# Field

- A *field*, denoted by  $(F, +, \times)$ , is an *integral domain* whose elements satisfy the following additional property.

**Inverse** for multiplication: For every  $a \in F$ , there exists an element  $b$ , denoted by  $a^{-1}$ , such that  $ab = ba = 1$ .



# Field

- A *field*, denoted by  $(F, +, \times)$ , is an *integral domain* whose elements satisfy the following additional property.

**Inverse** for multiplication: For every  $a \in F$ , there exists an element  $b$ , denoted by  $a^{-1}$ , such that  $ab = ba = 1$ .

- **Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$  ?  
 $(\mathbb{Z}_p, +, \times)$  ?



# Field

- A *field*, denoted by  $(F, +, \times)$ , is an *integral domain* whose elements satisfy the following additional property.

**Inverse** for multiplication: For every  $a \in F$ , there exists an element  $b$ , denoted by  $a^{-1}$ , such that  $ab = ba = 1$ .

- **Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$  ?  
 $(\mathbb{Z}_p, +, \times)$  ?

- If  $\mathbb{F}$  is finite,  $\mathbb{F}$  is called a *finite field*.





# Field

- A *field*, denoted by  $(F, +, \times)$ , is an *integral domain* whose elements satisfy the following additional property.

**Inverse** for multiplication: For every  $a \in F$ , there exists an element  $b$ , denoted by  $a^{-1}$ , such that  $ab = ba = 1$ .

- **Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$  ?  
 $(\mathbb{Z}_p, +, \times)$  ?

- If  $\mathbb{F}$  is finite,  $\mathbb{F}$  is called a *finite field*.
- $\mathbb{F}_q = \mathbb{Z}_p = \{0, 1, \dots, p-1\}$  with the operations *addition*, *multiplication* of integers modulo  $p$ , is called a *prime field*
  - The properties can be verified

# Prime subfield and characteristic

- Consider a *finite field*  $\mathbb{F}$ , define  $S_r = 1 + 1 + \cdots + 1$  as sum of  $r$  1's for a positive integer  $r$ 
  - Let  $p$  be the smallest positive number with  $S_p = 0$ .  
If such a  $p$  exists, it must be *prime*



# Prime subfield and characteristic

- Consider a *finite field*  $\mathbb{F}$ , define  $S_r = 1 + 1 + \cdots + 1$  as sum of  $r$  1's for a positive integer  $r$ 
  - Let  $p$  be the smallest positive number with  $S_p = 0$ .  
If such a  $p$  exists, it must be *prime*
  - If  $p = a \cdot b$  with  $0 < a, b < p$ , then by *distributivity*,  $0 = S_p = S_a \cdot S_b$ . Then one of  $S_a, S_b$  must be 0, *contradicting* the minimality of  $p$ .



# Prime subfield and characteristic

- Consider a *finite field*  $\mathbb{F}$ , define  $S_r = 1 + 1 + \cdots + 1$  as sum of  $r$  1's for a positive integer  $r$ 
  - Let  $p$  be the smallest positive number with  $S_p = 0$ .  
If such a  $p$  exists, it must be *prime*
  - If  $p = a \cdot b$  with  $0 < a, b < p$ , then by *distributivity*,  $0 = S_p = S_a \cdot S_b$ . Then one of  $S_a, S_b$  must be 0, *contradicting* the minimality of  $p$ .
- This  $p$  is called the *characteristic* of the field  $\mathbb{F}$ .



# Prime subfield and characteristic

- Consider a *finite field*  $\mathbb{F}$ , define  $S_r = 1 + 1 + \cdots + 1$  as sum of  $r$  1's for a positive integer  $r$ 
  - Let  $p$  be the smallest positive number with  $S_p = 0$ .  
If such a  $p$  exists, it must be *prime*
  - If  $p = a \cdot b$  with  $0 < a, b < p$ , then by *distributivity*,  $0 = S_p = S_a \cdot S_b$ . Then one of  $S_a, S_b$  must be 0, *contradicting* the minimality of  $p$ .
- This  $p$  is called the *characteristic* of the field  $\mathbb{F}$ .
- The subset  $\{0, S_1, S_2, \dots, S_{p-1}\} \subseteq \mathbb{F}$  is *isomorphic* to  $\mathbb{F}$  (*prime field*)



# Prime subfield and characteristic

- Consider a *finite field*  $\mathbb{F}$ , define  $S_r = 1 + 1 + \cdots + 1$  as sum of  $r$  1's for a positive integer  $r$ 
  - Let  $p$  be the smallest positive number with  $S_p = 0$ .  
If such a  $p$  exists, it must be *prime*
  - If  $p = a \cdot b$  with  $0 < a, b < p$ , then by *distributivity*,  $0 = S_p = S_a \cdot S_b$ . Then one of  $S_a, S_b$  must be 0, *contradicting* the minimality of  $p$ .
- This  $p$  is called the *characteristic* of the field  $\mathbb{F}$ .
- The subset  $\{0, S_1, S_2, \dots, S_{p-1}\} \subseteq \mathbb{F}$  is *isomorphic* to  $\mathbb{F}$  (*prime field*)
- *Any* finite field  $\mathbb{F}$  is a *finite dimensional vector space* over  $\mathbb{F}_p$ , with  $n = \dim_{\mathbb{F}_p}(\mathbb{F})$ ,  $|\mathbb{F}| = p^n$ , i.e., the cardinality of  $\mathbb{F}$  must be a prime power



# Finite fields

- For **any** prime power  $q$ , there is essentially **only one** finite field of order  $q$ . Any two finite fields of order  $q$  are the **same** except that the labelling used to represent the field elements may be different



# Finite fields

- For **any** prime power  $q$ , there is essentially **only one** finite field of order  $q$ . Any two finite fields of order  $q$  are the **same** except that the labelling used to represent the field elements may be different
- *Binary field – characteristic-2* finite fields  $\mathbb{F}_{2^m}$ 
  - Elements are polynomials over  $\mathbb{F}_2$  of degree  $\leq m - 1$
  - $\mathbb{F}_{2^m} := \{a_{m-1}x^{m-1} + a_{m-2}x^{m-2} + \cdots + a_2x^2 + a_1x + a_0 : a_i \in \mathbb{F}_2\}$





# Finite fields

- For **any** prime power  $q$ , there is essentially **only one** finite field of order  $q$ . Any two finite fields of order  $q$  are the **same** except that the labelling used to represent the field elements may be different
- *Binary field – characteristic-2* finite fields  $\mathbb{F}_{2^m}$ 
  - Elements are polynomials over  $\mathbb{F}_2$  of degree  $\leq m - 1$
  - $\mathbb{F}_{2^m} := \{a_{m-1}x^{m-1} + a_{m-2}x^{m-2} + \dots + a_2x^2 + a_1x + a_0 : a_i \in \mathbb{F}_2\}$
- An *irreducible polynomial*  $f(x)$  of degree  $m$  is chosen:  
 $f(x)$  **cannot** be factored as a product of binary polynomials each of degree less than  $m$ 
  - *Addition*: usual
  - *Multiplication*: modulo  $f(x)$



# Hard problems

- So far, we have only discussed number-theoretic problem that are *easy*
  - E.g., addition, multiplication, modular arithmetic, exponentiation



# Hard problems

- So far, we have only discussed number-theoretic problem that are *easy*
  - E.g., addition, multiplication, modular arithmetic, exponentiation
- Some problems are (*conjectured* to be) *hard*



# Factoring

- Multiplying two numbers is *easy*; factoring a number *hard*
  - Given  $x, y$ , *easy* to compute  $x \cdot y$
  - Given  $N$ , *hard* (in general) to find  $x, y > 1$  such that  $x \cdot y = N$



# Factoring

- Multiplying two numbers is *easy*; factoring a number *hard*
  - Given  $x, y$ , *easy* to compute  $x \cdot y$
  - Given  $N$ , *hard* (in general) to find  $x, y > 1$  such that  $x \cdot y = N$
- Compare:
  - Multiply 10101023 and 29100257
  - Find the factors of 293942365262911



# Factoring

- It is **not** hard to factor most numbers
  - 50% of the time, random number is even
  - $1/3$  of the time, random number is divisible by 3



# Factoring

- It is **not** hard to factor most numbers
  - 50% of the time, random number is even
  - 1/3 of the time, random number is divisible by 3
- The hardest numbers to factor are those that are the *product* of two, **equal-length** primes



# Factoring

- It is **not** hard to factor most numbers
  - 50% of the time, random number is even
  - 1/3 of the time, random number is divisible by 3
- The hardest numbers to factor are those that are the *product* of two, **equal-length** primes
- The *RSA problem* is related to *factoring*





# The RSA problem

- Let  $N = pq$  with  $p$  and  $q$  distinct, odd primes
- $\mathbb{Z}_N^*$  = *invertible* elements under multiplication modulo  $N$ 
  - The order of  $\mathbb{Z}_N^*$  is  $\phi(N) = (p - 1) \cdot (q - 1)$
  - $\phi(N)$  is *easy* to compute if  $p, q$  are known
  - $\phi(N)$  is *hard* to compute if  $p, q$  are *not* known
    - Equivalent (believed) to factoring  $N$



# The RSA problem

- Let  $N = pq$  with  $p$  and  $q$  distinct, odd primes
- $\mathbb{Z}_N^*$  = *invertible* elements under multiplication modulo  $N$ 
  - The order of  $\mathbb{Z}_N^*$  is  $\phi(N) = (p - 1) \cdot (q - 1)$
  - $\phi(N)$  is *easy* to compute if  $p, q$  are known
  - $\phi(N)$  is *hard* to compute if  $p, q$  are *not* known
    - Equivalent (believed) to factoring  $N$
- Fix  $e$  with  $\gcd(e, \phi(N)) = 1$ 
  - Raising to the  $e$ -th power is a permutation of  $\mathbb{Z}_N^*$



# The RSA problem

- Let  $N = pq$  with  $p$  and  $q$  distinct, odd primes
- $\mathbb{Z}_N^*$  = *invertible* elements under multiplication modulo  $N$ 
  - The order of  $\mathbb{Z}_N^*$  is  $\phi(N) = (p - 1) \cdot (q - 1)$
  - $\phi(N)$  is *easy* to compute if  $p, q$  are known
  - $\phi(N)$  is *hard* to compute if  $p, q$  are *not* known
    - Equivalent (believed) to factoring  $N$
- Fix  $e$  with  $\gcd(e, \phi(N)) = 1$ 
  - Raising to the  $e$ -th power is a permutation of  $\mathbb{Z}_N^*$
- If  $ed \equiv 1 \pmod{\phi(N)}$ , raising to the  $d$ -th power is the *inverse* of raising to the  $e$ -th power
  - I.e.,  $(x^e)^d \equiv x \pmod{N}$
  - $x^d$  is the  $e$ -th root of  $x$  modulo  $N$



# The RSA problem

- If  $p, q$  are known:
  - $\Rightarrow \phi(N)$  can be computed
  - $\Rightarrow d = e^{-1} \bmod \phi(N)$  can be computed
  - $\Rightarrow$  possible to compute  $e$ -th roots modulo  $N$



# The RSA problem

- If  $p, q$  are known:
  - $\Rightarrow \phi(N)$  can be computed
  - $\Rightarrow d = e^{-1} \bmod \phi(N)$  can be computed
  - $\Rightarrow$  possible to compute  $e$ -th roots modulo  $N$
- If  $p, q$  are *not* known:
  - $\Rightarrow$  computing  $\phi(N)$  is as hard as factoring  $N$
  - $\Rightarrow$  computing  $d$  is as hard as factoring  $N$



# The RSA problem

- If  $p, q$  are known:
    - $\Rightarrow \phi(N)$  can be computed
    - $\Rightarrow d = e^{-1} \bmod \phi(N)$  can be computed
    - $\Rightarrow$  possible to compute  $e$ -th roots modulo  $N$
  - If  $p, q$  are *not* known:
    - $\Rightarrow$  computing  $\phi(N)$  is as hard as factoring  $N$
    - $\Rightarrow$  computing  $d$  is as hard as factoring  $N$
- Q*: Given  $d$  and  $e$ , can we factor  $N$ ?



# The RSA problem

- If  $p, q$  are known:
    - $\Rightarrow \phi(N)$  can be computed
    - $\Rightarrow d = e^{-1} \bmod \phi(N)$  can be computed
    - $\Rightarrow$  possible to compute  $e$ -th roots modulo  $N$
  - If  $p, q$  are *not* known:
    - $\Rightarrow$  computing  $\phi(N)$  is as hard as factoring  $N$
    - $\Rightarrow$  computing  $d$  is as hard as factoring  $N$
- Q*: Given  $d$  and  $e$ , can we factor  $N$ ?
- Very useful for *public-key* cryptography



# The RSA problem

- **Informally**: given  $N$ ,  $e$ , and uniform element  $y \in \mathbb{Z}_N^*$ , compute the  $e$ -th root of  $y$





# The RSA problem

- *Informally*: given  $N$ ,  $e$ , and uniform element  $y \in \mathbb{Z}_N^*$ , compute the  $e$ -th root of  $y$
- *RSA assumption*: this is a **hard** problem!



# The RSA problem

- **Informally**: given  $N$ ,  $e$ , and uniform element  $y \in \mathbb{Z}_N^*$ , compute the  $e$ -th root of  $y$
- ***RSA assumption***: this is a **hard** problem!
- **Formally**:
- **GenRSA**: on input  $1^n$ , outputs  $(N, e, d)$  with  $N = pq$  a product of two distinct  $n$ -bit primes, with  $ed = 1 \bmod \phi(N)$



# The RSA problem

- **Informally**: given  $N$ ,  $e$ , and uniform element  $y \in \mathbb{Z}_N^*$ , compute the  $e$ -th root of  $y$
- **RSA assumption**: this is a **hard** problem!
- **Formally**:
- **GenRSA**: on input  $1^n$ , outputs  $(N, e, d)$  with  $N = pq$  a product of two distinct  $n$ -bit primes, with  $ed = 1 \bmod \phi(N)$
- **Experiment**  $\text{RSA-inv}_{A, \text{GenRSA}}(n)$ :
  - Compute  $(N, e, d) \leftarrow \text{GenRSA}(1^n)$
  - Choose uniform  $y \in \mathbb{Z}_N^*$
  - Run  $A(N, e, y)$  to get  $x$
  - Experiment evaluates to 1 if  $x^e = y \bmod N$



# The RSA assumption (formal)

- **GenRSA**: on input  $1^n$ , outputs  $(N, e, d)$  with  $N = pq$  a product of two distinct  $n$ -bit primes, with  $ed = 1 \bmod \phi(N)$
- **Experiment**  $\text{RSA-inv}_{A, \text{GenRSA}}(n)$ :
  - Compute  $(N, e, d) \leftarrow \text{GenRSA}(1^n)$
  - Choose uniform  $y \in \mathbb{Z}_N^*$
  - Run  $A(N, e, y)$  to get  $x$
  - Experiment evaluates to 1 if  $x^e = y \bmod N$

# The RSA assumption (formal)

- **GenRSA**: on input  $1^n$ , outputs  $(N, e, d)$  with  $N = pq$  a product of two distinct  $n$ -bit primes, with  $ed = 1 \bmod \phi(N)$
- **Experiment**  $\text{RSA-inv}_{A, \text{GenRSA}}(n)$ :
  - Compute  $(N, e, d) \leftarrow \text{GenRSA}(1^n)$
  - Choose uniform  $y \in \mathbb{Z}_N^*$
  - Run  $A(N, e, y)$  to get  $x$
  - Experiment evaluates to 1 if  $x^e = y \bmod N$
- The **RSA problem** is **hard** relative to **GenRSA** if for all PPT algorithms  $A$ ,

$$\Pr[\text{RSA-inv}_{A, \text{GenRSA}}(n) = 1] < \text{negl}(n)$$



# Implementing GenRSA

- One way to implement GenRSA:
  - Generate uniform  $n$ -bit primes  $p, q$
  - Set  $N := pq$
  - Choose arbitrary  $e$  with  $\gcd(e, \phi(N)) = 1$
  - Compute  $d := e^{-1} \bmod \phi(N)$
  - Output  $(N, e, d)$



# Implementing GenRSA

- One way to implement GenRSA:
  - Generate uniform  $n$ -bit primes  $p, q$
  - Set  $N := pq$
  - Choose arbitrary  $e$  with  $\gcd(e, \phi(N)) = 1$
  - Compute  $d := e^{-1} \bmod \phi(N)$
  - Output  $(N, e, d)$
- Choice of  $e$ ?
  - Does *not* seem to affect hardness of the *RSA problem*
  - $e = 3$  or  $e = 2^{16} + 1$  for *efficient* exponentiation



# RSA and factoring

- If factoring moduli output by **GenRSA** is easy, then the ***RSA problem*** is easy relative to **GenRSA**
  - Factoring is easy  $\Rightarrow$  RSA problem is easy





# RSA and factoring

- If factoring moduli output by *GenRSA* is easy, then the *RSA problem* is easy relative to *GenRSA*
  - Factoring is easy  $\Rightarrow$  RSA problem is easy
- Hardness of the *RSA problem* is **not** known to be implied by hardness of factoring
  - Possible factoring is hard but *RSA problem* is easy
  - Possible both are hard but *RSA problem* is “easier”
  - Currently, RSA is **believed** to be *as hard as factoring*



# Trapdoor functions

- **Definition 10.1** (*Trapdoor functions*) A *trapdoor function collection* is a collection  $\mathcal{F}$  of finite functions such that every  $f \in \mathcal{F}$  is a **one-to-one** function from some set  $S_f$  to a set  $T_f$ . The following properties are required.

- **Efficient generation, computation and inversion**

There is a PPT algorithm  $G$  that on input  $1^n$  outputs a pair  $(f, f^{-1})$ , where these are two  $\text{poly}(n)$  size strings that describe the functions  $f, f^{-1}$

- **Efficient sampling** There is a PPT algorithm that given  $f$  can output a **random** element of  $S_f$

- **One-wayness** The function  $f$  is **hard to invert** without knowing the *inversion key*. For **all** PPT  $A$  there is a negligible function  $\epsilon$  s.t.

$$\Pr_{(f, f^{-1}) \leftarrow_R G(1^n), x \leftarrow_R S_f} [A(1^n, f, f(x)) = x] < \epsilon(n)$$

# RSA trapdoor function

- **Keys:** choose  $P, Q$  as random primes of length  $n$ ,  $N = P \cdot Q$ . Choose  $e$  at random from  $\{1, \dots, \phi(N) - 1\}$  with  $\gcd(e, \phi(N)) = 1$

**Forward Key:**  $N, e$

**Backward Key:**  $d$  with  $ed \equiv 1 \pmod{\phi(N)}$

**Function:**  $RSA_{N,e}(X) = X^e \pmod{N}$

**Inverse:** If  $Y = RSA_{N,e}(X) = X^e \pmod{N}$ , then  $Y^d \pmod{N} = X$ .

# RSA trapdoor function

- **Keys:** choose  $P, Q$  as random primes of length  $n$ ,  $N = P \cdot Q$ . Choose  $e$  at random from  $\{1, \dots, \phi(N) - 1\}$  with  $\gcd(e, \phi(N)) = 1$

Forward **Key:**  $N, e$

Backward **Key:**  $d$  with  $ed \equiv 1 \pmod{\phi(N)}$

**Function:**  $RSA_{N,e}(X) = X^e \pmod{N}$

**Inverse:** If  $Y = RSA_{N,e}(X) = X^e \pmod{N}$ , then  $Y^d \pmod{N} = X$ .

- **RSA Assumption:** the RSA function is indeed a *trapdoor function*
  - This is **stronger** than the assumption that **factoring** is **hard**



# Rabin's trapdoor function

- Assume that *factoring* random *Blum integers* is hard. A *Blum integer* is a number  $n = pq$  where  $p, q \equiv 3 \pmod{4}$ .



# Rabin's trapdoor function

- Assume that *factoring* random *Blum integers* is hard. A *Blum integer* is a number  $n = pq$  where  $p, q \equiv 3 \pmod{4}$ .
- Define  $\mathcal{B}_n := \{P \in [1 \dots 2^n] : P \text{ prime and } P \equiv 3 \pmod{4}\}$

**The Factoring Axiom** For *every* PPT algorithm  $A$  there is a negligible function  $\epsilon$  s.t.

$$\Pr_{P, Q \leftarrow_R \mathcal{B}_n} [A(P \cdot Q) = \{P, Q\}] < \epsilon(n)$$



# Rabin's trapdoor function

- **Keys**: choose  $P, Q$  as random primes of length  $n$  with  
 $P, Q \equiv 3 \pmod{4}, N = P \cdot Q$ .

Forward **Key**:  $N$

Backward **Key**:  $P, Q$

**Function**:  $Y = \text{RABIN}_N(X) = X^2 \pmod{N}$ , which is a permutation on  $QR_N$ , where  $QR_N$  denotes the set of quadratic residues modulo  $N$



# Rabin's trapdoor function

- **Keys:** choose  $P, Q$  as random primes of length  $n$  with  
 $P, Q \equiv 3 \pmod{4}, N = P \cdot Q$ .

Forward **Key:**  $N$

Backward **Key:**  $P, Q$

**Function:**  $Y = \text{RABIN}_N(X) = X^2 \pmod{N}$ , which is a permutation on  $QR_N$ , where  $QR_N$  denotes the set of quadratic residues modulo  $N$

**Inverse:** Compute  $A = Y \pmod{P}$  and  $B = Y \pmod{Q}$ . Since  $P, Q \equiv 3 \pmod{4}$ , let  $P = 4t + 3$  and  $Q = 4t' + 3$ .





# Rabin's trapdoor function

- **Keys:** choose  $P, Q$  as random primes of length  $n$  with  
 $P, Q \equiv 3 \pmod{4}, N = P \cdot Q$ .

Forward **Key:**  $N$

Backward **Key:**  $P, Q$

**Function:**  $Y = \text{RABIN}_N(X) = X^2 \pmod{N}$ , which is a permutation on  $QR_N$ , where  $QR_N$  denotes the set of quadratic residues modulo  $N$

**Inverse:** Compute  $A = Y \pmod{P}$  and  $B = Y \pmod{Q}$ . Since  $P, Q \equiv 3 \pmod{4}$ , let  $P = 4t + 3$  and  $Q = 4t' + 3$ .

Compute  $X_1 = A^{t+1} \pmod{P}$  and  $X_2 = B^{t'+1} \pmod{Q}$ . Using CRT, we find  $X$ .

# Rabin's trapdoor function

- **Keys:** choose  $P, Q$  as random primes of length  $n$  with  
 $P, Q \equiv 3 \pmod{4}, N = P \cdot Q$ .

Forward **Key:**  $N$

Backward **Key:**  $P, Q$

**Function:**  $Y = \text{RABIN}_N(X) = X^2 \pmod{N}$ , which is a permutation on  $QR_N$ , where  $QR_N$  denotes the set of quadratic residues modulo  $N$

**Inverse:** Compute  $A = Y \pmod{P}$  and  $B = Y \pmod{Q}$ . Since  $P, Q \equiv 3 \pmod{4}$ , let  $P = 4t + 3$  and  $Q = 4t' + 3$ .

Compute  $X_1 = A^{t+1} \pmod{P}$  and  $X_2 = B^{t'+1} \pmod{Q}$ . Using CRT, we find  $X$ .

We know that  $X = S^2 \pmod{P}$ , then

$$X_1 = (X^2)^{t+1} = S^{4(t+1)} = S^{P-1+2} = S^2 = X \pmod{P}.$$

Similarly,  $X_2 = S^2 = X \pmod{Q}$ .



# Rabin's trapdoor function

- **Lemma 10.2** Let  $X, Y$  be such that  $X \not\equiv \pm Y \pmod{N}$  but  $X^2 \equiv Y^2 \pmod{N}$ . Then  $\gcd(X - Y, N) \notin \{1, N\}$ .

**Proof.** easy.



# Rabin's trapdoor function

- **Lemma 10.2** Let  $X, Y$  be such that  $X \not\equiv \pm Y \pmod{N}$  but  $X^2 \equiv Y^2 \pmod{N}$ . Then  $\gcd(X - Y, N) \notin \{1, N\}$ .

**Proof.** easy.

**Theorem 10.3** (*One-wayness of Rabin's function*)

Rabin's function is a *trapdoor function* under the factoring axiom.



# Rabin's trapdoor function

- **Lemma 10.2** Let  $X, Y$  be such that  $X \not\equiv \pm Y \pmod{N}$  but  $X^2 \equiv Y^2 \pmod{N}$ . Then  $\gcd(X - Y, N) \notin \{1, N\}$ .

**Proof.** easy.

**Theorem 10.3** (*One-wayness of Rabin's function*)

Rabin's function is a *trapdoor function* under the factoring axiom.

**Proof.** By contradiction. (see blackboard)



# Next Lecture

- public key encryption ...

