

## 2 Solving LPs: The Simplex Algorithm of George Dantzig

### 2.1 Simplex Pivoting: Dictionary Format

We illustrate a general solution procedure, called the *simplex algorithm*, by implementing it on a very simple example. Consider the LP

$$(2.1) \quad \begin{aligned} \max \quad & 5x_1 + 4x_2 + 3x_3 \\ \text{s.t.} \quad & 2x_1 + 3x_2 + x_3 \leq 5 \\ & 4x_1 + x_2 + 2x_3 \leq 11 \\ & 3x_1 + 4x_2 + 2x_3 \leq 8 \\ & 0 \leq x_1, x_2, x_3 \end{aligned}$$

In devising our solution procedure we take a standard mathematical approach; reduce the problem to one that we already know how to solve. Since the structure of this problem is essentially linear, we will try to reduce it to a problem of solving a system of linear equations, or perhaps a series of such systems. By encoding the problem as a system of linear equations we bring into play our knowledge and experience with such systems in the new context of linear programming.

In order to encode the LP (2.1) as a system of linear equations we must first transform linear inequalities into linear equation. This is done by introducing a new non-negative variable, called a *slack variable*, for each inequality:

$$\begin{aligned} x_4 &= 5 - [2x_1 + 3x_2 + x_3] \geq 0, \\ x_5 &= 11 - [4x_1 + x_2 + 2x_3] \geq 0, \\ x_6 &= 8 - [3x_1 + 4x_2 + 2x_3] \geq 0. \end{aligned}$$

To handle the objective, we introduce a new variable  $z$ :

$$z = 5x_1 + 4x_2 + 3x_3.$$

Then all of the information associated with the LP (2.1) can be coded as follows:

$$(2.2) \quad \begin{aligned} 2x_1 + 3x_2 + x_3 + x_4 &= 5 \\ 4x_1 + x_2 + 2x_3 + x_5 &= 11 \\ 3x_1 + 4x_2 + 2x_3 + x_6 &= 8 \\ -z + 5x_1 + 4x_2 + 3x_3 &= 0 \\ 0 \leq x_1, x_2, x_3, x_4, x_5, x_6. \end{aligned}$$

The new variables  $x_4$ ,  $x_5$ , and  $x_6$  are called slack variables since they take up the *slack* in the linear inequalities. This system can also be written using block structured matrix notation as

$$\begin{bmatrix} 0 & A & I \\ -1 & c^T & 0 \end{bmatrix} \begin{bmatrix} z \\ x \end{bmatrix} = \begin{bmatrix} b \\ 0 \end{bmatrix},$$

where

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 4 & 1 & 2 \\ 3 & 4 & 2 \end{bmatrix}, \quad I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 11 \\ 8 \end{bmatrix}, \quad \text{and } c = \begin{bmatrix} 5 \\ 4 \\ 3 \end{bmatrix}.$$

The augmented matrix associated with the system (2.2) is

$$(2.3) \quad \left[ \begin{array}{ccc|c} 0 & A & I & b \\ -1 & c & 0 & 0 \end{array} \right]$$

and is referred to as the *initial simplex tableau* for the LP (2.1).

Again consider the system

$$(2.4) \quad \begin{aligned} x_4 &= 5 - 2x_1 - 3x_2 - x_3 \\ x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\ z &= 5x_1 + 4x_2 + 3x_3. \end{aligned}$$

This system defines the variables  $x_4$ ,  $x_5$ ,  $x_6$  and  $z$  as linear combinations of the variables  $x_1$ ,  $x_2$ , and  $x_3$ . We call this system a *dictionary* for the LP (2.1). More specifically, it is the *initial dictionary* for the the LP (2.1). This initial dictionary defines the objective value  $z$  and the slack variables as a linear combination of the initial decision variables. The variables that are “defined” in this way are called the *basic variables*, while the remaining variables are called *nonbasic*. The set of all basic variables is called the *basis*. A particular solution to this system is easily obtained by setting the non-basic variables equal to zero. In this case, we get

$$\begin{aligned} x_4 &= 5 \\ x_5 &= 11 \\ x_6 &= 8 \\ z &= 0. \end{aligned}$$

Note that the solution

$$(2.5) \quad \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 5 \\ 11 \\ 8 \end{pmatrix}$$

is feasible for the extended system (2.2) since all components are non-negative. For this reason, we call the dictionary (2.4) a *feasible dictionary* for the LP (2.1), and we say that this LP has *feasible origin*.

In general, a dictionary for the LP (2.1) is any system of 4 linear equations that defines three of the variables  $x_1, \dots, x_6$  and  $z$  in terms of the remaining 3 variables and has the same solution set as the initial dictionary. The variables other than  $z$  that are being defined in the dictionary are called the basis for the dictionary, and the remaining variables are said to be non-basic in the dictionary. Every dictionary identifies a particular solution to the linear system obtained by setting the non-basic variables equal to zero. Such a solution is said to be a *basic feasible solution* (BFS) for the LP (2.1) if it componentwise non-negative, that is, all of the numbers in the vector are non-negative so that the point lies in the feasible region for the LP.

The grand strategy of the simplex algorithm is to move from one feasible dictionary representation of the system (2.2) to another (and hence from one BFS to another) while simultaneously increasing the value of the objective variable  $z$ . In the current setting, beginning with the dictionary (2.4), what strategy might one employ in order to determine a new dictionary whose associated BFS gives a greater value for the objective variable  $z$ ?

Each feasible dictionary is associated with one and only one feasible point. This is the associated BFS obtained by setting all of the non-basic variables equal to zero. This is how we obtain (2.5). To change the feasible point identified in this way, we need to increase the value of one of the non-basic variables from its current value of zero. Note that we cannot decrease the value of a non-basic variable since we wish to remain feasible, that is, we wish to keep all variables non-negative.

Note that the coefficient of each of the non-basic variables in the representation of the objective value  $z$  in (2.4) is positive. Hence, if we pick any one of these variables and increase its value from zero while leaving remaining two at zero, we automatically increase the value of the objective variable  $z$ . Since the coefficient on  $x_1$  in the representation of  $z$  is the greatest, we can increase  $z$  the fastest by increasing  $x_1$ .

By how much can we increase  $x_1$  and still remain feasible? For example, if we increase  $x_1$  to 3 then (2.4) says that  $x_4 = -1$ ,  $x_5 = -1$ ,  $x_6 = -1$  which is not feasible. Let us consider this question by examining the equations in (2.4) one by one. Note that the first equation in the dictionary (2.4),

$$x_4 = 5 - 2x_1 - 3x_2 - x_3,$$

shows that  $x_4$  remains non-negative as long as we do not increase the value of  $x_1$  beyond  $5/2$  (remember,  $x_2$  and  $x_3$  remain at the value zero). Similarly, using the second equation in the dictionary (2.4),

$$x_5 = 11 - 4x_1 - x_2 - 2x_3,$$

$x_5$  remains non-negative if  $x_1 \leq 11/4$ . Finally, the third equation in (2.4),

$$x_6 = 8 - 3x_1 - 4x_2 - 2x_3,$$

implies that  $x_6$  remains non-negative if  $x_1 \leq 8/3$ . Therefore, we remain feasible, i.e. keep **all** variables non-negative, if our increase to the variable  $x_1$  remains less than

$$\frac{5}{2} = \min \left\{ \frac{5}{2}, \frac{11}{4}, \frac{8}{3} \right\}.$$

If we now increase the value of  $x_1$  to  $\frac{5}{2}$ , then the value of  $x_4$  is driven to zero. One way to think of this is that  $x_1$  *enters the basis while*  $x_4$  *leaves the basis*. Mechanically, we obtain the new dictionary having  $x_1$  basic and  $x_4$  non-basic by using the defining equation for  $x_4$  in the current dictionary:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3.$$

By moving  $x_1$  to the left hand side of this equation and  $x_4$  to the right, we get the new equation

$$2x_1 = 5 - x_4 - 3x_2 - x_3$$

or equivalently

$$x_1 = \frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}x_3.$$

The variable  $x_1$  can now be *eliminated* from the remaining two equations in the dictionary by substituting in this equation for  $x_1$  where it appears in these equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}x_3 \\ x_5 &= 11 - 4\left(\frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}x_3\right) - x_2 - 2x_3 \\ &= 1 + 2x_4 + 5x_2 \\ x_6 &= 8 - 3\left(\frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}x_3\right) - 4x_2 - 2x_3 \\ &= \frac{1}{2} + \frac{3}{2}x_4 + \frac{1}{2}x_2 - \frac{1}{2}x_3 \\ z &= 5\left(\frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}x_3\right) + 4x_2 + 3x_3 \\ &= \frac{25}{2} - \frac{5}{2}x_4 - \frac{7}{2}x_2 + \frac{1}{2}x_3. \end{aligned}$$

When this substitution is complete, we have the new dictionary and the new BFS:

$$\begin{aligned} (2.6) \quad x_1 &= \frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}x_3 \\ x_5 &= 1 + 2x_4 + 5x_2 \\ x_6 &= \frac{1}{2} + \frac{3}{2}x_4 + \frac{1}{2}x_2 - \frac{1}{2}x_3 \\ z &= \frac{25}{2} - \frac{5}{2}x_4 - \frac{7}{2}x_2 + \frac{1}{2}x_3, \end{aligned}$$

and the associated BFS is

$$(2.7) \quad \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 5/2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1/2 \end{pmatrix} \quad \text{with} \quad z = \frac{25}{2}.$$

This process might seem very familiar to you. It is simply Gaussian elimination. As we know from our knowledge of linear systems of equations, Gaussian elimination can be performed in a matrix context with the aid of the augmented matrix (or, simplex tableau) (2.3). We return to this observation later to obtain a more efficient computational technique.

We now have a new dictionary (2.6) which identifies the basic feasible solution (BFS) (2.7) with associated objective value  $z = \frac{25}{2}$ . Can we improve on this BFS and obtain a higher objective value? Let's try the same trick again, and repeat the process we followed in going from the initial dictionary (2.4) to the new dictionary (2.6). Note that the coefficient of  $x_3$  in the representation of  $z$  in the new dictionary (2.6) is positive. Hence if we increase the value of  $x_3$  from zero, we will increase the value of  $z$ . By how much can we increase the value of  $x_3$  and yet keep all the remaining variables non-negative? As before, we see that the first equation in the dictionary (2.6) combined with the need to keep  $x_1$  non-negative implies that we cannot increase  $x_3$  by more than  $(5/2)/(1/2) = 5$ . However, the second equation in (2.6) places no restriction on increasing  $x_3$  since  $x_3$  does not appear in this equation. Finally, the third equation in (2.6) combined with the need to keep  $x_6$  non-negative implies that we cannot increase  $x_3$  by more than  $(1/2)/(1/2) = 1$ . Therefore, in order to preserve the non-negativity of all variables, we can increase  $x_3$  by at most

$$1 = \min\{5, 1\}.$$

When we do this  $x_6$  is driven to zero, so  $x_3$  enters the basis and  $x_6$  leaves. More precisely, first move  $x_3$  to the left hand side of the defining equation for  $x_6$  in (2.6),

$$\frac{1}{2}x_3 = \frac{1}{2} + \frac{3}{2}x_4 + \frac{1}{2}x_2 - x_6,$$

or, equivalently,

$$x_3 = 1 + 3x_4 + x_2 - 2x_6,$$

then substitute this expression for  $x_3$  into the remaining equations,

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{1}{2}x_4 - \frac{3}{2}x_2 - \frac{1}{2}[1 + 3x_4 + x_2 - 2x_6] \\ &= 2 - 2x_4 - 2x_2 + x_6 \\ x_5 &= 1 + 2x_4 + 5x_2 \\ z &= \frac{25}{2} - \frac{5}{2}x_4 - \frac{7}{2}x_2 + \frac{1}{2}[1 + 3x_4 + x_2 - 2x_6] \\ &= 13 - x_4 - 3x_2 - x_6, \end{aligned}$$

yielding the dictionary

$$\begin{aligned} x_3 &= 1 + 3x_4 + x_2 - 2x_6 \\ x_1 &= 2 - 2x_4 + 2x_2 + x_6 \\ x_5 &= 1 + 2x_4 + 2x_2 \\ z &= 13 - x_4 - 3x_2 - x_6 \end{aligned}$$

which identifies the feasible solution

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

having objective value  $z = 13$ .

Can we do better? NO! This solution is optimal! The coefficient on the variables in the cost row of the dictionary,

$$z = 13 - x_4 - 3x_2 - x_6,$$

are all non-positive, so increasing any one of their values will not increase the value of the objective. (Why does this prove optimality?) The process of moving from one feasible dictionary to the next is called a *simplex pivot*. The overall process of stringing a sequence of simplex pivots together in order to locate an optimal solution is called the *Simplex Algorithm*. The simplex algorithm is consistently ranked as one of the ten most important algorithmic discoveries of the 20th century (<http://www.uta.edu/faculty/rcli/TopTen/topten.pdf>). The algorithm is generally attributed to George Dantzig (1914-2005) who is known as the father of linear programming. In 1984 Narendra Karmarkar published a paper describing a new approach to solving linear programs that was both numerically efficient and had *polynomial complexity*. This new class of methods are called *interior point* methods. These new methods have revolutionized the optimization field over the last 30 years, and they have led to efficient numerical methods for a wide variety of optimization problems well beyond the confines of linear programming. However, the simplex algorithm continues as an important numerical method for solving LPs, and for many specially structured LPs it is still the most efficient algorithm.

## 2.2 Simplex Pivoting: Tableau Format (Augmented Matrix Format)

We now review the implementation of the simplex algorithm by applying Gaussian elimination to the augmented matrix (2.3), also known as the simplex tableau. For this problem, the initial simplex tableau is given by

$$(2.8) \quad \left[ \begin{array}{ccc|c} 0 & A & I & b \\ -1 & c & 0 & 0 \end{array} \right] = \left[ \begin{array}{cccccc|c} 0 & 2 & 3 & 1 & 1 & 0 & 5 \\ 0 & 4 & 1 & 2 & 0 & 1 & 11 \\ 0 & 3 & 4 & 2 & 0 & 0 & 8 \\ -1 & 5 & 4 & 3 & 0 & 0 & 0 \end{array} \right].$$

Each simplex pivot on a dictionary corresponds to one step of Gaussian elimination on the augmented matrix associated with the dictionary. For example, in the first simplex pivot,  $x_1$

enters the basis and  $x_4$  leaves the basis. That is, we use the first equation of the dictionary to rewrite  $x_1$  as a function of the remaining variables, and then use this representation to eliminate  $x_1$  from the remaining equations. In terms of the augmented matrix (2.8), this corresponds to first making the coefficient for  $x_1$  in the first equation the number 1 by dividing this first equation through by 2. Then use this entry to eliminate the column under  $x_1$ , that is, make all other entries in this column zero (Gaussian elimination):

Pivot column							ratios		← Pivot row
	↓ 2						↓ 5/2		
0	2	3	1	1	0	0	5	5/2	
0	4	1	2	0	1	0	11	11/4	
0	3	4	2	0	0	1	8	8/3	
-1	5	4	3	0	0	0	0		
<hr/>									
0	1	3/2	1/2	1/2	0	0	5/2		
0	0	-5	0	-2	1	0	1		
0	0	-1/2	1/2	-3/2	0	1	1/2		
<hr/>									
-1	0	-7/2	1/2	-5/2	0	0	-25/2		

In this illustration, we have placed a line above the cost row to delineate its special roll in the pivoting process. In addition, we have also added a column on the right hand side which contains the ratios that we computed in order to determine the pivot row. Recall that we must use the smallest ratio in order to keep all variables in the associated BFS non-negative. Note that we performed the exact same arithmetic operations but in the more efficient matrix format. The new augmented matrix,

$$(2.9) \quad \left[ \begin{array}{cccccc|c} 0 & 1 & 3/2 & 1/2 & 1/2 & 0 & 0 & 5/2 \\ 0 & 0 & -5 & 0 & -2 & 1 & 0 & 1 \\ 0 & 0 & -1/2 & 1/2 & -3/2 & 0 & 1 & 1/2 \\ -1 & 0 & -7/2 & 1/2 & -5/2 & 0 & 0 & -25/2 \end{array} \right],$$

is the augmented matrix for the dictionary (2.6).

The initial augmented matrix (2.8) has basis  $x_4$ ,  $x_5$ , and  $x_6$ . The columns associated with these variables in the initial tableau (2.8) are distinct columns of the identity matrix. Correspondingly, the basis for the second tableau is  $x_1$ ,  $x_5$ , and  $x_6$ , and again this implies that the columns for these variables in the tableau (2.9) are the corresponding distinct columns of the identity matrix. In tableau format, this will always be true of the basic variables, i.e., their associated columns are distinct columns of the identity matrix. To recover the BFS (basic feasible solution) associated with this tableau we first set the non-basic variables equal to zero (i.e. the variables not associated with columns of the identity matrix (except in very unusual circumstances)):  $x_2 = 0$ ,  $x_3 = 0$ , and  $x_4 = 0$ . To find the value of the basic variables go to the column associated with that variable (for example,  $x_1$  is in the second

column), in that column find the row with the number 1 in it, then in that row go to the number to the right of the vertical bar (for  $x_1$  this is the first row with the number to the right of the bar being  $5/2$ ). Then set this basic variable equal to that number ( $x_1 = 5/2$ ). Repeating this for  $x_5$  and  $x_6$  we get  $x_5 = 1$  and  $x_6 = 1/2$ . To get the corresponding value for  $z$ , look at the  $z$  row and observe that the corresponding linear equation is

$$-z - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 = -\frac{25}{2},$$

but  $x_2$ ,  $x_3$ , and  $x_4$  are non-basic and so take the value zero giving  $-z = -25/2$ , or  $z = 25/2$ .

Of course this is all exactly the same information we obtained from the dictionary approach. The simplex, or augmented matrix approach is simply a more efficient computational procedure. We will use the simplex procedure in class to solve LPs. In addition, on quizzes and exams you will be required to understand how to go back and forth between these two representations, i.e the dictionary representation and its corresponding simplex tableau (or, augmented matrix). Let us now continue with the second simplex pivot.

In every tableau we always reserve the bottom row for encoding the linear relationship between the objective variable  $z$  and the currently non-basic variables. For this reason we call this row the *cost row*, and to distinguish its special role, we place a line above it in the tableau (this is reminiscent of the way we place a vertical bar in an augmented matrix to distinguish the right hand side of a linear equation). In the cost row of the tableau (2.9),

$$[-1, 0, -7/2, 1/2, -5/2, 0, 0, | -25/2],$$

we see a positive coefficient,  $1/2$ , in the 4th column. Hence the cost row coefficient for the non-basic variable  $x_3$  in this tableau is  $1/2$ . This indicates that if we increase the value of  $x_3$ , we also increase the value of the objective  $z$ . This is not true for any of the other currently non-basic variables since their cost row coefficients are all non-positive. Thus, the only way we can increase the value of  $z$  is to bring  $x_3$  into the basis, or equivalently, pivot on the  $x_3$  column which is the 4th column of the tableau. For this reason, we call the  $x_3$  column the *pivot column*. Now if  $x_3$  is to enter the basis, then which variable leaves? Just as with the dictionary representation, the variable that leaves the basis is that currently basic variable whose non-negativity places the greatest restriction on increasing the value of  $x_3$ . This restriction is computed as the smallest ratio of the right hand sides and the positive coefficients in the  $x_3$  column:

$$1 = \min\{(5/2)/(1/2), (1/2)/(1/2)\}.$$

The ratios are only computed with the positive coefficients since a non-positive coefficient means that by increasing this variable we do not decrease the value of the corresponding basic variable and so it is not a restricting equation. Since the minimum ratio in this instance is 1 and it comes from the third row, we find that the *pivot row* is the third row. Looking across the third row, we see that this row identifies  $x_6$  as a basic variable since the  $x_6$  column is a column of the identity with a 1 in the third row. Hence  $x_6$  is the variable leaving the



basis when  $x_3$  enters. The intersection of the pivot column and the pivot row is called the *pivot*. In this instance it is the number  $1/2$  which is the  $(3, 4)$  entry of the simplex tableau. Pivoting on this entry requires us to first make it 1 by multiplying this row through by 2, and then to apply Gaussian elimination to force all other entries in this column to zero:

Pivot column ↓							r	
0	1	3/2	1/2	1/2	0	0	5/2	5
0	0	-5	0	-2	1	0	1	
0	0	-1/2	1/2	-3/2	0	1	1/2	① ← pivot row
-1	0	-7/2	1/2	-5/2	0	0	-25/2	
<hr/>								
0	1	2	0	2	0	-1	2	
0	0	-5	0	-2	1	0	1	
0	0	-1	1	-3	0	2	1	
-1	0	-3	0	-1	0	-1	-13	

This simplex tableau is said to be optimal since it is feasible (the associated BFS is non-negative) and the cost row coefficients for the variables are all non-positive. A BFS that is optimal is called an *optimal basic feasible solution*. The optimal BFS is obtained by setting the non-basic variables equal to zero and setting the basic variables equal to the value on the right hand side corresponding to the one in its column:  $x_1 = 2$ ,  $x_2 = 0$ ,  $x_3 = 1$ ,  $x_4 = 0$ ,  $x_5 = 1$ ,  $x_6 = 0$ . The optimal objective value is obtained by taking the negative of the number in the lower right hand corner of the optimal tableau:  $z = 13$ .

We now recap the complete sequence of pivots in order to make a final observation that will help streamline the pivoting process: pivots are circled,

0	②	3	1	1	0	0	5
0	4	1	2	0	1	0	11
0	3	4	2	0	0	1	8
-1	5	4	3	0	0	0	0
<hr/>							
0	1	3/2	1/2	1/2	0	0	5/2
0	0	-5	0	-2	1	0	1
0	0	-1/2	1/2	-3/2	0	1	1/2
-1	0	-7/2	1/2	-5/2	0	0	-25/2
<hr/>							
0	1	2	0	2	0	-1	2
0	0	-5	0	-2	1	0	1
0	0	-1	1	-3	0	2	1
-1	0	-3	0	-1	0	-1	-13

Observe from this sequence of pivots that the  $z$  column is never touched, that is, it remains the same in all tableaus. Essentially, it just serves as a place holder reminding us that in the linear equation for the cost row the coefficient of  $z$  is  $-1$ . Therefore, for the sake of expediency we will drop this column from our simplex computations in most settings, and simply re-insert it whenever instructive or convenient. However, *it is very important to always remember that it is there!* Indeed, we will make explicit and essential use of this column in order to arrive at a full understanding of the duality theory for linear programming. After removing this column, the above pivots take the following form:

$\textcircled{2}$	3	1	1	0	0	5
4	1	2	0	1	0	11
3	4	2	0	0	1	8
5	4	3	0	0	0	0
<hr/>						
1	3/2	1/2	1/2	0	0	5/2
0	-5	0	-2	1	0	1
0	-1/2	$\textcircled{1/2}$	-3/2	0	1	1/2
0	-7/2	1/2	-5/2	0	0	-25/2
<hr/>						
1	2	0	2	0	-1	2
0	-5	0	-2	1	0	1
0	-1	1	-3	0	2	1
0	-3	0	-1	0	-1	-13

We close this section with a final example of simplex pivoting on a tableau giving only the essential details.

### The LP

$$\begin{aligned}
 &\text{maximize} && 3x &+& 2y &-& 4z \\
 &\text{subject to} && x &+& 4y && \leq 5 \\
 &&& 2x &+& 4y &-& 2z \leq 6 \\
 &&& x &+& y &-& 2z \leq 2 \\
 &&& 0 &\leq x, y, z
 \end{aligned}$$

### Simplex Iterations

1	4	0	1	0	0	5	ratios
2	4	-2	0	1	0	6	5
①	1	-2	0	0	1	2	3
3	2	-4	0	0	0	0	2
0	3	2	1	0	-1	3	3/2
0	2	②	0	1	-2	2	1
1	1	-2	0	0	1	2	
0	-1	2	0	0	-3	-6	
0	1	0	1	-1	1	1	
0	1	1	0	1/2	-1	1	
1	3	0	0	1	-1	4	
0	-3	0	0	-1	-1	-8	

Optimal Solution

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix} \quad \text{optimal value} = 8$$

A final word of advise, when doing simplex pivoting by hand, it is helpful to keep the tableaux vertically aligned in order to keep track of the arithmetic operations. Lined paper helps to keep the rows straight. But the columns need to be straight as well. Many students find that it is easy to keep both the rows and columns straight if they do pivoting on graph paper having large boxes for the numbers.

## 2.3 Dictionaries: The General Case for LPs in Standard Form

Recall the following standard form for LPs:

$$\begin{aligned} \mathcal{P} : \quad & \text{maximize} \quad c^T x \\ & \text{subject to} \quad Ax \leq b \\ & \quad \quad \quad 0 \leq x, \end{aligned}$$

where  $c \in \mathbb{R}^n$ ,  $b \in \mathbb{R}^m$ ,  $A \in \mathbb{R}^{m \times n}$  and the inequalities  $Ax \leq b$  and  $0 \leq x$  are to be interpreted componentwise. We now provide a formal definition for a dictionary associated with an LP in standard form. Let

$$(D_I) \quad \begin{aligned} x_{n+i} &= b_i - \sum_{j=1}^n a_{ij} x_j \\ z &= \sum_{j=1}^n c_j x_j \end{aligned}$$

be the defining system for the slack variables  $x_{n+i}$ ,  $i = 1, \dots, n$  and the objective variable  $z$ . A dictionary for  $\mathcal{P}$  is any system of the form

$$(D_B) \quad \begin{aligned} x_i &= \widehat{b}_i - \sum_{j \in N} \widehat{a}_{ij} x_j & i \in B \\ z &= \widehat{z} + \sum_{j \in N} \widehat{c}_j x_j \end{aligned}$$

where  $B$  and  $N$  are index sets contained in the set of integers  $\{1, \dots, n+m\}$  satisfying

- (1)  $B$  contains  $m$  elements,
- (2)  $B \cap N = \emptyset$
- (3)  $B \cup N = \{1, 2, \dots, n+m\}$ ,

and such that the systems  $(D_I)$  and  $(D_B)$  have identical solution sets. The set  $\{x_j : j \in B\}$  is said to be the basis associated with the dictionary  $(D_B)$  (sometimes we will refer to the index set  $B$  as the basis for the sake of simplicity), and the variables  $x_i$ ,  $i \in N$  are said to be the non-basic variables associated with this dictionary. The point identified by this dictionary is

$$(2.10) \quad \begin{aligned} x_i &= \widehat{b}_i & i \in B \\ x_j &= 0 & j \in N. \end{aligned}$$

The dictionary is said to be feasible if  $0 \leq \widehat{b}_i$  for  $i \in B$ . If the dictionary  $D_B$  is feasible, then the point identified by the dictionary (2.10) is said to be a basic feasible solution (BFS) for the LP. A feasible dictionary and its associated BFS are said to be optimal if  $\widehat{c}_j \leq 0$   $j \in N$ .

### Simplex Pivoting by Matrix Multiplication

As we have seen simplex pivoting can either be performed on dictionaries or on the augmented matrices that encode the linear equations of a dictionary in matrix form. In matrix form, simplex pivoting reduces to our old friend, Gaussian elimination. In this section, we show that Gaussian elimination can be represented as a consequence of left multiplication by a specially designed matrix called a *Gaussian pivot matrix*.

Consider the vector  $v \in \mathbb{R}^m$  block decomposed as

$$v = \begin{bmatrix} a \\ \alpha \\ b \end{bmatrix}$$

where  $a \in \mathbb{R}^s$ ,  $\alpha \in \mathbb{R}$ , and  $b \in \mathbb{R}^t$  with  $m = s + 1 + t$ . Assume that  $\alpha \neq 0$ . We wish to determine a matrix  $G$  such that

$$Gv = e_{s+1}$$

where for  $j = 1, \dots, n$ ,  $e_j$  is the unit coordinate vector having a one in the  $j$ th position and zeros elsewhere. We claim that the matrix

$$G = \begin{bmatrix} I_{s \times s} & -\alpha^{-1}a & 0 \\ 0 & \alpha^{-1} & 0 \\ 0 & -\alpha^{-1}b & I_{t \times t} \end{bmatrix}$$

does the trick. Indeed,

$$Gv = \begin{bmatrix} I_{s \times s} & -\alpha^{-1}a & 0 \\ 0 & \alpha^{-1} & 0 \\ 0 & -\alpha^{-1}b & I_{t \times t} \end{bmatrix} \begin{pmatrix} a \\ \alpha \\ b \end{pmatrix} = \begin{bmatrix} a - a \\ \alpha^{-1}\alpha \\ -b + b \end{bmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = e_{s+1}.$$

The matrix  $G$  is called a *Gaussian Pivot Matrix*. Note that  $G$  is invertible since

$$G^{-1} = \begin{bmatrix} I & a & 0 \\ 0 & \alpha & 0 \\ 0 & b & I \end{bmatrix},$$

and that for any vector of the form  $w = \begin{pmatrix} x \\ 0 \\ y \end{pmatrix}$  where  $x \in \mathbb{R}^s$   $y \in \mathbb{R}^t$ , we have

$$Gw = w.$$

The Gaussian pivot matrices perform precisely the operations required in order to execute a simplex pivot. That is, each simplex pivot can be realized as left multiplication of the simplex tableau by the appropriate Gaussian pivot matrix.

For example, consider the following initial feasible tableau:

$$\left[ \begin{array}{cccccc|c} 1 & 4 & 2 & 1 & 0 & 0 & 11 \\ 3 & \textcircled{2} & 1 & 0 & 1 & 0 & 5 \\ 4 & 2 & 2 & 0 & 0 & 1 & 8 \\ \hline 4 & 5 & 3 & 0 & 0 & 0 & 0 \end{array} \right]$$

where the (2, 2) element is chosen as the pivot element. In this case,

$$s = 1, \quad t = 2, \quad a = 4, \quad \alpha = 2, \quad \text{and} \quad b = \begin{bmatrix} 2 \\ 5 \end{bmatrix},$$

and so the corresponding Gaussian pivot matrix is

$$G_1 = \begin{bmatrix} I_{1 \times 1} & -\alpha^{-1}a & 0 \\ 0 & \alpha^{-1} & 0 \\ 0 & -\alpha^{-1}b & I_{2 \times 2} \end{bmatrix} = \begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & -\frac{5}{2} & 0 & 1 \end{bmatrix}.$$

Multiplying the simplex on the left by  $G_1$  gives

$$\begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & \frac{-5}{2} & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 2 & 1 & 0 & 0 & | & 11 \\ 3 & 2 & 1 & 0 & 1 & 0 & | & 5 \\ 4 & 2 & 2 & 0 & 0 & 1 & | & 8 \\ \hline 4 & 5 & 3 & 0 & 0 & 0 & | & 0 \end{bmatrix} = \begin{bmatrix} -5 & 0 & 0 & 1 & -2 & 0 & | & 1 \\ \frac{3}{2} & 1 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & | & \frac{5}{2} \\ 1 & 0 & \textcircled{1} & 0 & -1 & 1 & | & 3 \\ \hline -\frac{7}{2} & 0 & \frac{1}{2} & 0 & \frac{-5}{2} & 0 & | & \frac{-25}{2} \end{bmatrix}.$$

Repeating this process with the new pivot element in the (3,3) position yields the Gaussian pivot matrix

$$G_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \frac{-1}{2} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & \frac{-1}{2} & 1 \end{bmatrix},$$

and left multiplication by  $G_2$  gives

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \frac{-1}{2} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & \frac{-1}{2} & 1 \end{bmatrix} \begin{bmatrix} -5 & 0 & 0 & 1 & -2 & 0 & | & 1 \\ \frac{3}{2} & 1 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & | & \frac{5}{2} \\ 1 & 0 & 1 & 0 & -1 & 1 & | & 3 \\ \hline -\frac{7}{2} & 0 & \frac{1}{2} & 0 & \frac{-5}{2} & 0 & | & \frac{-25}{2} \end{bmatrix} = \begin{bmatrix} -5 & 0 & 0 & 1 & -2 & 0 & | & 1 \\ 1 & 1 & 0 & 0 & 1 & \frac{-1}{2} & | & 1 \\ 1 & 0 & 1 & 0 & -1 & 1 & | & 3 \\ \hline -4 & 0 & 0 & 0 & \frac{-3}{2} & \frac{-1}{2} & | & -14 \end{bmatrix}$$

yielding the optimal tableau.

If

$$(2.4) \quad \begin{bmatrix} A & I & b \\ c^T & 0 & 0 \end{bmatrix}$$

is the initial tableau, then

$$G_2 G_1 \begin{bmatrix} A & I & b \\ c^T & 0 & 0 \end{bmatrix} = \begin{bmatrix} -5 & 0 & 0 & 1 & -2 & 0 & | & 1 \\ 1 & 1 & 0 & 0 & 1 & \frac{-1}{2} & | & 1 \\ 1 & 0 & 1 & 0 & -1 & 1 & | & 3 \\ \hline -4 & 0 & 0 & 0 & -2 & \frac{-1}{2} & | & -14 \end{bmatrix}$$

That is, we would be able to go directly from the initial tableau to the optimal tableau if we knew the matrix

$$G = G_2 G_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \frac{-1}{2} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & \frac{-1}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & \frac{-5}{2} & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & 1 & \frac{-1}{2} & 0 \\ 0 & -1 & 1 & 0 \\ 0 & -2 & \frac{-1}{2} & 1 \end{bmatrix}$$

beforehand. Moreover, the matrix  $G$  is invertible since both  $G_1$  and  $G_2$  are invertible:

$$G^{-1} = G_1^{-1}G_2^{-1} = \begin{bmatrix} 1 & 4 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 5 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \frac{1}{2} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & \frac{1}{2} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 4 & 2 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 2 & 3 & 0 \\ 0 & 5 & 3 & 1 \end{bmatrix}$$

(you should check that  $GG^{-1} = I$  by doing the multiplication by hand). In general, every sequence of simplex pivots has a representation as left multiplication by some invertible matrix since pivoting corresponds to left multiplication of the tableau by a Gaussian pivot matrix, and Gaussian pivot matrices are always invertible. We now examine the consequence of this observation more closely in the general case. In this discussion, it is essential that we include the column associated with the objective variable  $z$  which we have largely ignored up to this point.

Recall the initial simplex tableau, or augmented matrix associated with the system  $(D_I)$ :

$$T_0 = \begin{bmatrix} 0 & A & I & b \\ -1 & c^T & 0 & 0 \end{bmatrix}.$$

In this discussion, it is essential that we include the first column, i.e. the column associated with the objective variable  $z$  in the augmented matrix. Let the matrix

$$T_k = \begin{bmatrix} 0 & \hat{A} & R & \hat{b} \\ -1 & \hat{c}^T & -y^T & \hat{z} \end{bmatrix}$$

be another simplex tableau obtained from the initial tableau after a series of  $k$  simplex pivots. Note that the first column remains unchanged. Indeed, the fact that simplex pivots do not alter the first column is the reason why we drop it in our hand computations. But in the discussion that follows its presence and the fact that it remains unchanged by simplex pivoting is very important. Since  $T_k$  is another simplex tableau the  $m \times (n + m)$  matrix  $[\hat{A} \ R]$  must possess among its columns the  $m$  columns of the  $m \times m$  identity matrix. These columns of the identity matrix correspond precisely to the basic variables associated with this tableau.

Our prior discussion on Gaussian pivot matrices tells us that  $T_k$  can be obtained from  $T_0$  by multiplying  $T_0$  on the left by some nonsingular  $(m + 1) \times (m + 1)$  matrix  $G$  where  $G$  is the product of a sequence of Gaussian pivot matrices. In order to better understand the action of  $G$  on  $T_0$  we need to decompose  $G$  into a block structure that is conformal with that of  $T_0$ :

$$G = \begin{bmatrix} M & u \\ v^T & \beta \end{bmatrix},$$

where  $M \in \mathbb{R}^{m \times m}$ ,  $u, v \in \mathbb{R}^m$ , and  $\beta \in \mathbb{R}$ . Then

$$\begin{aligned} \begin{bmatrix} 0 & \hat{A} & R & \hat{b} \\ -1 & \hat{c}^T & -y^T & \hat{z} \end{bmatrix} &= T_k \\ &= GT_0 \\ &= \begin{bmatrix} M & u \\ v^T & \beta \end{bmatrix} \begin{bmatrix} 0 & A & I & b \\ -1 & c^T & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} -u & MA + uc^T & M & Mb \\ -\beta & v^T A + \beta c^T & v^T & v^T b \end{bmatrix} . \end{aligned}$$

By equating the blocks in the matrices on the far left and far right hand sides of this equation, we find from the first column that

$$u = 0 \quad \text{and} \quad \beta = 1 .$$

Here we see the key role played by our knowledge of the structure of the objective variable column (the first column). From the  $(1, 3)$  and the  $(2, 3)$  terms on the far left and right hand sides of (2.5), we also find that

$$M = R, \quad \text{and} \quad v = -y .$$

Putting all of this together gives the following representation of the  $k^{th}$  tableau  $T_k$ :

$$(2.5) \quad T_k = \begin{bmatrix} R & 0 \\ -y^T & 1 \end{bmatrix} \begin{bmatrix} 0 & A & I & b \\ -1 & c^T & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & RA & R & Rb \\ -1 & c^T - y^T A & -y^T & -y^T b \end{bmatrix} ,$$

where the matrix  $R$  is necessarily invertible since the matrix

$$G = \begin{bmatrix} R & 0 \\ -y^T & 1 \end{bmatrix}$$

is invertible (prove this!):

$$G^{-1} = \begin{bmatrix} R^{-1} & 0 \\ y^T R^{-1} & 1 \end{bmatrix} . \quad (\text{check by multiplying out } GG^{-1})$$

The matrix  $R$  is called the *record* matrix for the tableau as it keeps track of all of the transformations required to obtain the new tableau. Again, the variables associated with the columns of the identity correspond to the basic variables. The tableau  $T_k$  is said to be *primal feasible*, or just *feasible*, if  $\hat{b} = Rb \geq 0$ .

We will return to this beautiful algebraic structure for the simplex algorithm later in the notes as it is the key ingredient in our understanding of the duality theory of linear programming. But first we address the question of whether or not the simplex algorithm works. That is, is the simplex algorithm a reliable methods for solving LPs?