

DEVELOPMENT OF MOLECULAR TOOLS TO UNDERSTAND METABOLIC REGULATION OF FUNGAL MORPHOGENESIS



Submitted by: **M Adishree**
Guided by: **Dr. Sriram Varahan**
(Senior Scientist, CSIR-CCMB)

CONTENTS

1. Acknowledgements
2. Aim
3. Objectives
4. Introduction
5. Rationale
6. Methodology, Results and Conclusion
7. References
8. Experiences and Skills Learned

ACKNOWLEDGEMENTS

Firstly, I would like to express my sincere gratitude to RICH-STEM program for providing me the platform to resume my career in science. My sincere thanks to Mrs. Aboli Ingle for her continuous support, motivation, and guidance.

I would like to extend my gratitude to my supervisor and mentor, Dr. Sriram Varahan for giving me the opportunity to work on this exciting and innovative project and for encouraging me to do my best. I thank my fellow lab mates Dhrumi Shah, Avanti Vairagkar, Nivedhitha R, Riya Bhattacharjee and Nikita Rewatkar for the stimulating discussions, for always lending a helping hand, and for all the fun we have had in the past months. I thank CSIR-CCMB for providing me the infrastructure and resources needed for my project.

Lastly, I would like to thank my friends and family for their endless love and support.

AIM

To develop molecular tools to characterize metabolic processes orchestrating fungal morphogenesis.

OBJECTIVES

1. To determine the metabolic state of pseudohyphal cells.
2. To test whether spatiotemporal metabolic heterogeneity influences pseudohyphal differentiation.

INTRODUCTION


- Cells in an isogenic population are capable of showing visual, behavioral, and functional heterogeneity.
- Studying pseudohyphal differentiation in *Saccharomyces cerevisiae* can provide fundamental insights into understanding the general biological mechanisms that underlie the adaptation of microorganisms to different environments.







RATIONALE

Previous studies in the lab identified central carbon metabolic pathways like glycolysis and pentose phosphate pathway (PPP) to be critical for pseudohyphal differentiation. The aim of my project was to determine the spatio-temporal activity/expression of glycolysis when *S. cerevisiae* undergoes pseudohyphal differentiation. This in turn, will allow us to understand how these metabolic pathways drive these morphogenetic switching events.

METHODOLOGY, RESULTS & CONCLUSION


(I) **EPITOPE TAGGING:** To check the spatiotemporal expression of glycolytic enzymes using western blotting.



Strain: Cdc19-HA (Epitope tagged)		Hxk2-HA (Epitope tagged)		Pgi1-HA (Epitope tagged)	
(colony – 1)	(colony – 2)	(colony – 1)	(colony – 2)	(colony – 1)	(colony – 2)
					
[SLAD day 10 imaging]		[SLAD day 10 imaging]		[SLAD day 10 imaging]	


Pyruvate kinase (Cdc19), hexokinase (Hxk2) and phosphoglucose isomerase (Pgi1) were epitope tagged to determine the spatio-temporal expression of these glycolytic enzymes. Colonies that were spotted on synthetic low ammonium dextrose (SLAD) media were imaged and cells were isolated for western blotting.







(II) **PROMOTER-REPORTER ASSAY:** To check the spatiotemporal activity of glycolysis and related pathways using fluorescence microscopy.



(III) GENERATION OF KNOCKOUT STRAINS

Previous studies in the field have shown that two strains namely, $\Sigma 1278b$ and CEN.PK undergo pseudohyphal differentiation under nitrogen-limiting conditions (1,2). Work from Dr. Sriram Varahan's lab (CSIR-CCMB) has shown that glycolysis is critical for driving pseudohyphal differentiation since disruption of genes that encode for glycolytic enzymes, alcohol dehydrogenase (*adh1*) and phosphofructokinase (*pfk1*) resulted in significant decrease in pseudohyphal differentiation in the $\Sigma 1278b$ diploid strain. In order to determine if glycolysis-dependent regulation of pseudohyphal differentiation is conserved across yeast strains, we generated knockout strains that lack genes encoding for the glycolytic enzymes, *adh1* and *pfk1* in the CEN.PK strain background using the following protocol.



CEN.PK (WT)		$\Delta\Delta adh1$		$\Delta\Delta pfk1$	
(colony – 1)	(colony – 2)	(colony – 1)	(colony – 2)	(colony – 1)	(colony – 2)
					
[SLAD day 10 imaging]		[SLAD day 10 imaging]		[SLAD day 10 imaging]	

Genes encoding glycolytic enzymes, alcohol dehydrogenase (*adh1*) and phosphofructokinase (*pfk1*) were deleted using homologous recombination and then spotted on SLAD medium along with CEN.PK (WT). They were imaged after 10 days. Disruption of *adh1* and *pfk1* in CEN.PK background results in significant reduction in pseudohyphal differentiation.

REFERENCES

- 1) Gimeno CJ, Ljungdahl PO, Styles CA, Fink GR. Unipolar cell divisions in the yeast *S. cerevisiae* lead to filamentous growth: regulation by starvation and RAS. *Cell*. 1992 Mar 20;68(6):1077-90. doi: 10.1016/0092-8674(92)90079-r. PMID: 1547504.
- 2) Laxman S, Tu BP. Multiple TORC1-associated proteins regulate nitrogen starvation-dependent cellular differentiation in *Saccharomyces cerevisiae*. *PLoS One*. 2011;6(10):e26081. doi: 10.1371/journal.pone.0026081. Epub 2011 Oct 17. PMID: 22043304; PMCID: PMC3197150.