

Aaghosh RAG Protocols: Localized Pakistan Context

Structured Action-Response Pairs Extracted from UNICEF Pakistan ECD Manual

System Note: These protocols are specifically adapted for the cultural, familial, and infrastructural context of Pakistan. They integrate the roles of extended family (like mothers-in-law) and utilize locally available resources.

Category: First-Aid & Health Emergencies

[Tags: infant, toddler, choking, suffocation, emergency]

Child Action / Event: The child is choking on an object or food.

Approved Parent Response Protocol:

1. **Positioning:** Hold the baby face-down along your thigh with their head lower than their bottom.
2. **Intervention:** Give them five blows firmly on their back between the shoulder blades.
3. **Airflow:** Open windows and doors to allow an abundant supply of fresh air.
4. **Escalate:** Take the child to the hospital immediately.

[Tags: health, high fever, home treatment]

Child Action / Event: The child develops a high fever.

Approved Parent Response Protocol:

1. **Cooling:** Remove the child's clothes and cool the body using a piece of cloth soaked in lukewarm water.
2. **Hydration:** Give more fluids to the child, and breastfeed repeatedly in the case of babies.
3. **Triage/Escalation:** If the baby/child has convulsions/fits, is unable to drink, or is abnormally sleepy, take them to a health facility immediately.

[Tags: health, diarrhea, hydration, nutrition]

Child Action / Event: The child has diarrhea (passing watery stools 3-4 times in a day).

Approved Parent Response Protocol:

1. **Preparation:** Wash hands. Mix one ORS sachet in 4 cups of clean, boiled, cool water.
2. **Administer:** Give one cup of ORS to the child after every stool.
3. **Hygiene:** Wash hands before and after cooking, feeding the child, and using the latrine to prevent further spread.

Category: Age-Specific Developmental Play & Stimulation

[Tags: infant, 0-6 months, stimulation, play]

Child State: The infant is awake and alert.

Approved Parent Response Protocol:

1. **Physical Contact:** Maintain frequent skin-to-skin contact (hugging, carrying).
2. **Visual/Auditory:** Slowly move colorful things for the child to see and reach for (e.g., shaker rattle, big ring on a string).
3. **Interaction:** Talk to the child during breastfeeding and make eye contact; play Peek-a-Boo.

[Tags: toddler, 1-2 years, independence, cognitive development]

Child State: The toddler is showing increasing mobility and curiosity.

Approved Parent Response Protocol:

1. **Responsibility:** Send the child on little errands in the house to develop independence.
2. **Cognitive Challenge:** Provide problem-solving toys.
3. **Validation:** Hug the child and praise them for little actions they complete.

Category: Cultural Family Dynamics & Positive Discipline

[Tags: family dynamics, joint family, mother-in-law, conflict resolution]

Parent Challenge: A mother-in-law or extended family member opposes a modern parenting/health practice (e.g., antenatal checkups or exclusive breastfeeding).

Approved Response Protocol (ABC Method for FLWs/Parents):

1. **Ask & Listen:** Do not be instantly judgmental or critical. Listen to the family member's views to earn their confidence and understand their reasoning.
2. **Brainstorm:** Jointly look for a solution that is acceptable and doable in the context of the joint family.
3. **Coach & Praise:** Break the desired behavior into smaller steps and praise the family for any small positive changes observed.

[Tags: positive discipline, verbal abuse, communication]

Child Action: The child does something wrong or misbehaves.

Approved Parent Response Protocol:

1. **De-escalate:** Do not use physical punishment (spanking, hitting, slapping, pinching, shaking) or verbal abuse.
2. **Communicate:** Talk one-to-one with the child calmly so that they understand the consequences of their action.
3. **Avoid Humiliation:** Do not tell the child they are "dumb" or "bad," and strictly avoid comparing the child with other children.