

THE LINEAR MODEL

• Projections

Theorem: Let $\mathbf{a}: n \times 1$ be a non-zero vector and let $\mathbf{d}: n \times 1$ be any vector. Then \mathbf{d} can be uniquely expressed as:

$$\mathbf{d} = \mathbf{b}_1 + \mathbf{b}_2$$

where \mathbf{b}_1 depends on \mathbf{a} and \mathbf{b}_2 is orthogonal to \mathbf{a} . Furthermore, $\mathbf{b}_1 = c\mathbf{a}$ with $c = \frac{\mathbf{d}'\mathbf{a}}{\mathbf{a}'\mathbf{a}}$ and $\mathbf{b}_2 = \mathbf{d} - \mathbf{b}_1$.

Proof:

Let $\mathbf{d} = \mathbf{b}_1 + \mathbf{b}_2 = c\mathbf{a} + \mathbf{b}_2$ where $\mathbf{b}_2 \perp \mathbf{a}$. Then $\mathbf{a}'\mathbf{d} = c\mathbf{a}'\mathbf{a} + \mathbf{0}$ or $c = \frac{\mathbf{d}'\mathbf{a}}{\mathbf{a}'\mathbf{a}}$ since $\mathbf{a}'\mathbf{a} \neq \mathbf{0}$.

Conversely, if $c = \frac{\mathbf{d}'\mathbf{a}}{\mathbf{a}'\mathbf{a}}$ and $\mathbf{b}_2 = \mathbf{d} - \mathbf{b}_1$, then $\mathbf{b}_2 = \mathbf{d} - c\mathbf{a}$. Therefore

$\mathbf{b}_2 \perp \mathbf{a}$ and since c is unique, \mathbf{b}_2 is also unique.

The vector \mathbf{b}_1 is the projection of \mathbf{d} on \mathbf{a} and \mathbf{b}_2 is the component of \mathbf{d} orthogonal to \mathbf{a} . The square of the length of the projection of \mathbf{d} on \mathbf{a} is

$$\mathbf{b}_1'\mathbf{b}_1 = c^2\mathbf{a}'\mathbf{a} = \frac{(\mathbf{d}'\mathbf{a})^2}{\mathbf{a}'\mathbf{a}}.$$

Theorem: Let V be a vector space and let \mathbf{d} be any vector. Then \mathbf{d} can be uniquely expressed as:

$$\mathbf{d} = \mathbf{c} + \mathbf{e}$$

where $\mathbf{c} \in V$ and $\mathbf{e} \perp V$. If V is generated by the columns of $\mathbf{A}: n \times m$, then

$$\mathbf{c} = \mathbf{A}(\mathbf{A}'\mathbf{A})^*\mathbf{A}'\mathbf{d} \quad \text{and} \quad \mathbf{e} = (\mathbf{I} - \mathbf{A}(\mathbf{A}'\mathbf{A})^*\mathbf{A}')\mathbf{d}$$

Proof:

Let $\mathbf{d} = \mathbf{c} + \mathbf{e}$ where $\mathbf{c} \in V$ and $\mathbf{e} \perp V$. Then $\mathbf{d} = \mathbf{A}\mathbf{c}_1 + \mathbf{e}$ for some \mathbf{c}_1 . In other words

$$\mathbf{A}'\mathbf{A}\mathbf{c}_1 = \mathbf{A}'\mathbf{d} \quad \text{since} \quad \mathbf{A}'\mathbf{e} = \mathbf{0}.$$

These equations to determine \mathbf{c}_1 are consistent, because $\text{rank}(\mathbf{A}'\mathbf{A}) = \text{rank}(\mathbf{A}'\mathbf{A}, \mathbf{A}'\mathbf{d})$. A solution of $\mathbf{A}'\mathbf{A}\mathbf{c}_1 = \mathbf{A}'\mathbf{d}$ is

$$\mathbf{c}_1 = (\mathbf{A}'\mathbf{A})^* \mathbf{A}'\mathbf{d}$$

for any generalized inverse $(\mathbf{A}'\mathbf{A})^*$ of $\mathbf{A}'\mathbf{A}$. Therefore

$$\mathbf{c} = \mathbf{A}\mathbf{c}_1 = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}'\mathbf{d}$$

and

$$\mathbf{e} = \mathbf{d} - \mathbf{c} = (\mathbf{I} - \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}')\mathbf{d}.$$

Suppose now that a solution \mathbf{c}_1 exists to

$$\mathbf{A}'\mathbf{A}\mathbf{c}_1 = \mathbf{A}'\mathbf{d}.$$

Let $\mathbf{c} = \mathbf{A}\mathbf{c}_1$ and $\mathbf{e} = \mathbf{d} - \mathbf{c}$, then

$$\mathbf{d} = \mathbf{c} + \mathbf{e}$$

where $\mathbf{c} \in V$ and $\mathbf{e} \perp V$, since

$$\mathbf{A}'\mathbf{e} = \mathbf{A}'\mathbf{d} - \mathbf{A}'\mathbf{c} = \mathbf{A}'\mathbf{d} - \mathbf{A}'\mathbf{A}\mathbf{c}_1 = \mathbf{0}.$$

To prove the uniqueness of the expression, let

$$\mathbf{d} = \mathbf{A}\mathbf{c}_0 + \mathbf{e}_0$$

with $\mathbf{e}_0 \perp V$. Then

$$\mathbf{A}'\mathbf{d} = \mathbf{A}'\mathbf{A}\mathbf{c}_0 = \mathbf{A}'\mathbf{A}\mathbf{c}_1.$$

Therefore $\mathbf{A}'\mathbf{A}(\mathbf{c}_0 - \mathbf{c}_1) = \mathbf{0}$ and $(\mathbf{c}_0 - \mathbf{c}_1)'\mathbf{A}'\mathbf{A}(\mathbf{c}_0 - \mathbf{c}_1) = \mathbf{0}$. The last identity is the sum of squares of the elements of $\mathbf{A}(\mathbf{c}_0 - \mathbf{c}_1)$, which is zero. In other words, $\mathbf{A}(\mathbf{c}_0 - \mathbf{c}_1) = \mathbf{0}$ and $\mathbf{A}\mathbf{c}_0 = \mathbf{A}\mathbf{c}_1$. Furthermore,

$$\mathbf{e}_0 = \mathbf{d} - \mathbf{A}\mathbf{c}_0 = \mathbf{d} - \mathbf{A}\mathbf{c}_1 = \mathbf{e}$$

and the expression is unique.

The vector \mathbf{c} is the projection of \mathbf{d} on V and \mathbf{e} is the component of \mathbf{d} which is orthogonal to V . Thus, $\mathbf{c} = \mathbf{A}\mathbf{c}_1$ is the projection of \mathbf{d} ($\mathbf{A}'\mathbf{A}\mathbf{c}_1 = \mathbf{A}'\mathbf{d}$)

Conclusion: The equations

$$\mathbf{A}'\mathbf{A}\mathbf{x} = \mathbf{A}'\mathbf{y}$$

imply that $\mathbf{A}\mathbf{x}$ is the projection of \mathbf{y} on the vector space generated by the columns of \mathbf{A} . Thus, $\mathbf{A}\mathbf{x} = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}'\mathbf{y}$ is the projection of \mathbf{y} . And $\mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}'$ is the projection matrix.

Note: Let V^\perp represent the vector space orthogonal to V . Since V is generated by the columns of $\mathbf{A}: n \times m$ the $\dim(V) + \dim(V^\perp) = n$.

Theorem: The projection matrix $\mathbf{P} = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}'$ is unique, symmetrical and idempotent. (Unique with respect to the choice of the generalized inverse and unique with respect to the specific \mathbf{A} . The only condition is that the vector space V must be generated by the columns of \mathbf{A} .)

Proof:

1. Uniqueness: Suppose that \mathbf{P}_1 is such a projection matrix as well. Then

$$\mathbf{P}_1 \mathbf{d} = \mathbf{P} \mathbf{d} \quad \forall \mathbf{d} \quad \Rightarrow \quad \mathbf{P}_1 = \mathbf{P}$$

2. Symmetry: It follows from (1) and the fact that $(\mathbf{A}'\mathbf{A})^{*'} is also a generalized inverse of $\mathbf{A}'\mathbf{A}$. Therefore$

$$(\mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}')' = \mathbf{A}(\mathbf{A}'\mathbf{A})^{*'} \mathbf{A}' = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}'$$

3. Idempotent: $\mathbf{P}^2 \mathbf{d} = \mathbf{P}(\mathbf{P} \mathbf{d}) = \mathbf{P} \mathbf{d}$ since $\mathbf{P} \mathbf{d} \in V \quad \forall \mathbf{d}$. Therefore $\mathbf{P}^2 = \mathbf{P}$.

Corollary:

1. If $\mathbf{A} = (\mathbf{A}_1, \mathbf{A}_2)$ with $\mathbf{A}_1' \mathbf{A}_2 = \mathbf{0}$, it also follows that

$$\mathbf{P} = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}' = \mathbf{A}_1(\mathbf{A}_1' \mathbf{A}_1)^* \mathbf{A}_1' + \mathbf{A}_2(\mathbf{A}_2' \mathbf{A}_2)^* \mathbf{A}_2'.$$

2. If the columns of \mathbf{A} are for instance mutually orthogonal, it follows that:

(a)

$$\begin{aligned} \mathbf{P} = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}' &= (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m) \begin{pmatrix} (\mathbf{a}_1' \mathbf{a}_1)^{-1} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & (\mathbf{a}_2' \mathbf{a}_2)^{-1} & \dots & \mathbf{0} \\ \dots & \dots & \dots & \dots \\ \mathbf{0} & \mathbf{0} & \dots & (\mathbf{a}_m' \mathbf{a}_m)^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{a}_1' \\ \mathbf{a}_2' \\ \dots \\ \mathbf{a}_m' \end{pmatrix} \\ &= \sum_{i=1}^m \mathbf{a}_i (\mathbf{a}_i' \mathbf{a}_i)^{-1} \mathbf{a}_i'. \end{aligned}$$

(b) The projection of the vector \mathbf{y} on V is:

$$\mathbf{P} \mathbf{y} = \mathbf{A}(\mathbf{A}'\mathbf{A})^* \mathbf{A}' \mathbf{y} = \sum_{i=1}^m \mathbf{a}_i (\mathbf{a}_i' \mathbf{a}_i)^{-1} \mathbf{a}_i' \mathbf{y} = \sum_{i=1}^m \frac{\mathbf{a}_i' \mathbf{y}}{\mathbf{a}_i' \mathbf{a}_i} \mathbf{a}_i$$

which is the sum of the projections on the individual mutually orthogonal vectors.

(c) The square of the length of the projection of \mathbf{y} on V is:

$$\mathbf{y}'\mathbf{P}\mathbf{y} = \sum_{i=1}^m \frac{(\mathbf{a}'_i\mathbf{y})^2}{\mathbf{a}'_i\mathbf{a}_i}.$$

The projection matrix \mathbf{P} can be calculated easily by using a convenient basis for V . The following theorem can for instance be used.

Theorem: Suppose that V with $\text{rank}(V) = r$ is generated by the columns of $\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_r)$. It is always possible to select a mutually orthogonal basis for V , say the columns of $\mathbf{B} = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_r)$, in such a way that \mathbf{b}_s only depends on the first s columns of \mathbf{A} .

Proof:

Assume the vector \mathbf{a}_2 . It can be expressed uniquely as

$$\mathbf{a}_2 = \mathbf{b}_1 + \mathbf{b}_2$$

where \mathbf{b}_1 is dependent on \mathbf{a}_1 and \mathbf{b}_2 is orthogonal to \mathbf{a}_1 . The first two vectors \mathbf{a}_1 and \mathbf{a}_2 can be replaced by \mathbf{b}_1 and \mathbf{b}_2 in the basis, without changing V . If $\mathbf{b}_1 = \mathbf{0}$, let $\mathbf{b}_1 = \mathbf{a}_1$ and $\mathbf{b}_2 = \mathbf{a}_2$. The vector \mathbf{a}_3 can be uniquely expressed as

$$\mathbf{a}_3 = \mathbf{b} + \mathbf{b}_3$$

where \mathbf{b} depends on \mathbf{b}_1 and \mathbf{b}_2 and where \mathbf{b}_3 is orthogonal to \mathbf{b}_1 and \mathbf{b}_2 . It follows that

$$\mathbf{b} = \frac{\mathbf{a}'_3\mathbf{b}_1}{\mathbf{b}'_1\mathbf{b}_1}\mathbf{b}_1 + \frac{\mathbf{a}'_3\mathbf{b}_2}{\mathbf{b}'_2\mathbf{b}_2}\mathbf{b}_2$$

and

$$\mathbf{b}_3 = \mathbf{a}_3 - \mathbf{b}.$$

This process can be continued until all the \mathbf{a}_i 's are replaced by \mathbf{b}_i 's in the basis.

It follows that:

$$\begin{aligned} \mathbf{P} &= \mathbf{A}(\mathbf{A}'\mathbf{A})^*\mathbf{A}' = \mathbf{B}(\mathbf{B}'\mathbf{B})^*\mathbf{B}' = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_r) \begin{pmatrix} (\mathbf{b}'_1\mathbf{b}_1)^{-1} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & (\mathbf{b}'_2\mathbf{b}_2)^{-1} & \dots & \mathbf{0} \\ \dots & \dots & \dots & \dots \\ \mathbf{0} & \mathbf{0} & \dots & (\mathbf{b}'_r\mathbf{b}_r)^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{b}'_1 \\ \mathbf{b}'_2 \\ \dots \\ \mathbf{b}'_r \end{pmatrix} \\ &= \sum_{i=1}^r \mathbf{b}_i(\mathbf{b}'_i\mathbf{b}_i)^{-1}\mathbf{b}'_i. \end{aligned}$$

Examples:

1. Assume $\mathbf{A} = \begin{pmatrix} 2 & 0 \\ 1 & 1 \\ 3 & 1 \end{pmatrix}$. Then, it follows that:

$$\mathbf{b}_1 = \frac{2}{7} \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} \text{ and } \mathbf{b}_2 = \mathbf{a}_2 - \mathbf{b}_1 = \frac{1}{7} \begin{pmatrix} -4 \\ 5 \\ 1 \end{pmatrix}$$

The projection matrix is:

$$\begin{aligned} \mathbf{P} &= \frac{1}{14} \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} (2 \quad 1 \quad 3) + \frac{1}{42} \begin{pmatrix} -4 \\ 5 \\ 1 \end{pmatrix} (-4 \quad 5 \quad 1) \\ &= \frac{1}{14} \begin{pmatrix} 4 & 2 & 6 \\ 2 & 1 & 3 \\ 6 & 3 & 9 \end{pmatrix} + \frac{1}{42} \begin{pmatrix} 16 & -20 & -4 \\ -20 & 25 & 5 \\ -4 & 5 & 1 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 2 & -1 & 1 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix} \end{aligned}$$

A better choice for \mathbf{A} would be $\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$ since we can replace the first column with $\frac{1}{2}(\mathbf{a}_1 - \mathbf{a}_2)$

in $\mathbf{A} = \begin{pmatrix} 2 & 0 \\ 1 & 1 \\ 3 & 1 \end{pmatrix}$. Then, we have

$$\begin{aligned} \mathbf{b}_1 &= \frac{1}{2} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \\ \mathbf{b}_2 &= \mathbf{a}_2 - \mathbf{b}_1 = \frac{1}{2} \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix} \end{aligned}$$

The projection matrix is:

$$\begin{aligned} \mathbf{P} &= \frac{1}{2} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} (1 \quad 0 \quad 1) + \frac{1}{6} \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix} (-1 \quad 2 \quad 1) \\ &= \frac{1}{2} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} + \frac{1}{6} \begin{pmatrix} 1 & -2 & -1 \\ -2 & 4 & 2 \\ -1 & 2 & 1 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 2 & -1 & 1 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix} \end{aligned}$$

An alternative calculation of \mathbf{P} follows from the fact that

$\text{rank}(V^\perp) = 1$. V^\perp is for instance generated by

$$\mathbf{e} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$$

Therefore

$$\begin{aligned} \mathbf{I} - \mathbf{P} &= \mathbf{e}(\mathbf{e}'\mathbf{e})^{-1}\mathbf{e}' \\ &= \frac{1}{3} \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} (1 \quad -1) = \frac{1}{3} \begin{pmatrix} 1 & 1 & -1 \\ 1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix} \end{aligned}$$

and

$$\mathbf{P} = \frac{1}{3} \begin{pmatrix} 2 & -1 & 1 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix}$$

2. For the previous Example 2 it follows that:

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 3 & 1 & 1 \end{pmatrix}$$

An orthogonal basis for V is the columns of

$$\mathbf{B} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \\ 1 & 1 \end{pmatrix}$$

The projection matrix is:

$$\begin{aligned} \mathbf{P} &= \frac{1}{2} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} (1 \quad 0 \quad 1) + \frac{1}{3} \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} (-1 \quad 1 \quad 1) \\ &= \frac{1}{2} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{pmatrix} = \frac{1}{6} \begin{pmatrix} 5 & -2 & 1 \\ -2 & 2 & 2 \\ 1 & 2 & 5 \end{pmatrix} \end{aligned}$$

Again it follows that $\text{rank}(V^\perp) = 1$ and V^\perp is generated by

$$\mathbf{e} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$$

Therefore

$$\begin{aligned} \mathbf{I} - \mathbf{P} &= \mathbf{e}(\mathbf{e}'\mathbf{e})^{-1}\mathbf{e}' \\ &= \frac{1}{6} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} (1 \quad 2 \quad -1) = \frac{1}{6} \begin{pmatrix} 1 & 2 & -1 \\ 2 & 4 & -2 \\ -1 & -2 & 1 \end{pmatrix} \end{aligned}$$

and

$$\mathbf{P} = \frac{1}{6} \begin{pmatrix} 5 & -2 & 1 \\ -2 & 2 & 2 \\ 1 & 2 & 5 \end{pmatrix}$$

• Sums of Squares of Linear Sets

Let $\mathbf{y}: n \times 1 \sim (\boldsymbol{\mu}, \sigma^2 \mathbf{I})$. The sum of squares of the linear function $\mathbf{c}'\mathbf{y}$ is defined as the square of the length of the projection of \mathbf{y} on \mathbf{c} , namely:

$$\begin{aligned} s^2 &= \left(\frac{\mathbf{c}'\mathbf{y}}{\mathbf{c}'\mathbf{c}} \mathbf{c} \right)' \left(\frac{\mathbf{c}'\mathbf{y}}{\mathbf{c}'\mathbf{c}} \mathbf{c} \right) \\ &= \frac{(\mathbf{c}'\mathbf{y})^2}{\mathbf{c}'\mathbf{c}}. \end{aligned}$$

Note that this sum of squares is the same as the sum of squares of the linear function $d\mathbf{c}'\mathbf{y}$, for any non-zero constant d . This sum of squares is therefore only dependent on the vector space generated by \mathbf{c} .

It follows that:

$$\begin{aligned} E(s^2) &= E(\mathbf{c}'\mathbf{y})^2 / \mathbf{c}'\mathbf{c} \\ &= \{\text{var}(\mathbf{c}'\mathbf{y}) + [E(\mathbf{c}'\mathbf{y})]^2\} / \mathbf{c}'\mathbf{c} \\ &= \sigma^2 + [\mathbf{c}'E(\mathbf{y})]^2 / \mathbf{c}'\mathbf{c} \end{aligned}$$

The expected value of s^2 is therefore σ^2 plus a term which is obtained by replacing the random variables by their expected values in s^2 . Note that:

$$E(s^2) = \sigma^2 \Leftrightarrow E(\mathbf{c}'\mathbf{y}) = 0$$

The sum of squares of a linear set is defined similarly. Suppose that $\mathbf{C}: n \times m = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_m)$ and $\text{rank}(\mathbf{C}) = r$. Let V_C be the vector space generated by the columns of \mathbf{C} . Let L_C be the linear set of all linear functions with coefficient vectors in V_C :

$$L_C = \{\mathbf{c}'\mathbf{y} : \mathbf{c} \in V_C\}.$$

L_C is generated by the linear functions $\mathbf{c}'_1\mathbf{y}, \dots, \mathbf{c}'_m\mathbf{y}$ just as V_C is generated by $\mathbf{c}_1, \dots, \mathbf{c}_m$. The linear functions are dependent or independent corresponding to dependence or independence of the coefficient vectors. Any r independent linear functions in L_C will therefore generate the linear set L_C . We say that L_C has r degrees of freedom.

The sum of squares of the linear set L_C is defined as the square of the length of the projection of \mathbf{y} on V_C , namely:

$$S^2 = \{\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'\mathbf{y}\}'\{\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'\mathbf{y}\} = \mathbf{y}'\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'\mathbf{y}$$

since $\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'$ is unique, symmetrical and idempotent.

The expression for S^2 is independent of the specific functions $\mathbf{C}\mathbf{y}$ which generate L_C . Suppose that the columns of $\mathbf{B} = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_r)$ form a mutually orthogonal basis for V_C , so that $L_B = L_C$. It then follows that:

$$\begin{aligned} S^2 &= \mathbf{y}'\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'\mathbf{y} = \mathbf{y}'\mathbf{B}(\mathbf{B}'\mathbf{B})^*\mathbf{B}'\mathbf{y} \\ &= \mathbf{y}'(\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_r) \begin{pmatrix} (\mathbf{b}'_1\mathbf{b}_1)^{-1} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & (\mathbf{b}'_2\mathbf{b}_2)^{-1} & \dots & \mathbf{0} \\ \dots & \dots & \dots & \dots \\ \mathbf{0} & \mathbf{0} & \dots & (\mathbf{b}'_r\mathbf{b}_r)^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{b}'_1 \\ \mathbf{b}'_2 \\ \dots \\ \mathbf{b}'_r \end{pmatrix} \mathbf{y} \\ &= \sum_{i=1}^r \frac{(\mathbf{b}'_i\mathbf{y})^2}{\mathbf{b}'_i\mathbf{b}_i} = \sum_{i=1}^r s_i^2 \end{aligned}$$

where s_i^2 is the sum of squares of $\mathbf{b}'_i\mathbf{y}$. Also,

$$E(S^2) = \sum_{i=1}^r E(s_i^2) = r\sigma^2 + \sum_{i=1}^r (\mathbf{b}'_i E(\mathbf{y}))^2 / \mathbf{b}'_i\mathbf{b}_i.$$

It follows that:

$$E(S^2) = r\sigma^2 + S_\mu^2$$

where S_μ^2 is a term obtained by replacing the random variables in S^2 with their expected values.

Note that $E(S^2) = r\sigma^2 \Leftrightarrow$ if and only if $E(\mathbf{B}'\mathbf{y}) = \mathbf{0}$.

But $E(\mathbf{B}'\mathbf{y}) = \mathbf{0}$ if and only if $E(\mathbf{C}'\mathbf{y}) = \mathbf{0}$, since $\mathbf{B} = \mathbf{C}\mathbf{D}_1$ for some \mathbf{D}_1 and $\mathbf{C} = \mathbf{B}\mathbf{D}_2$ for some \mathbf{D}_2 .

Suppose that V_{C_1} and V_{C_2} with ranks r_1 and r_2 respectively, are mutually orthogonal vector spaces with corresponding linear sets L_{C_1} and L_{C_2} . Therefore $\mathbf{C}'_1\mathbf{C}_2 = \mathbf{0}$. The linear sets L_{C_1} and L_{C_2} are orthogonal. Let

$$V_C = V_{C_1} \oplus V_{C_2} \text{ and } L_C = L_{C_1} \oplus L_{C_2}.$$

The implication is that V_C is generated by the columns of $\mathbf{C} = (\mathbf{C}_1, \mathbf{C}_2)$. If S_i^2 is the sum of squares of L_{C_i} and S^2 is the sum of squares of L_C , it then follows that:

$$S^2 = \mathbf{y}'\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'\mathbf{y} = \mathbf{y}'\mathbf{C}_1(\mathbf{C}'_1\mathbf{C}_1)^*\mathbf{C}'_1\mathbf{y} + \mathbf{y}'\mathbf{C}_2(\mathbf{C}'_2\mathbf{C}_2)^*\mathbf{C}'_2\mathbf{y} = S_1^2 + S_2^2$$

The same line of reasoning can be used to show that if $V_{C_i}, i = 1, \dots, k$ are mutually orthogonal vector spaces with corresponding linear sets $L_{C_i}, i = 1, \dots, k$ and

$$V_C = \bigoplus V_{C_i} \text{ and } L_C = \bigoplus L_{C_i}$$

in other words V_C is generated by the columns of $\mathbf{C} = (\mathbf{C}_1, \mathbf{C}_2, \dots, \mathbf{C}_k)$, then

$$S^2 = \mathbf{y}'\mathbf{C}(\mathbf{C}'\mathbf{C})^*\mathbf{C}'\mathbf{y} = \sum_{i=1}^k \mathbf{y}'\mathbf{C}_i(\mathbf{C}'_i\mathbf{C}_i)^*\mathbf{C}'_i\mathbf{y} = \sum_{i=1}^k S_i^2$$

with S_i^2 the sum of squares of L_{C_i} .

Consider the linear set

$$L_C = \{\mathbf{c}'\mathbf{y} : \mathbf{c} \in V_C\}.$$

with $\mathbf{C}: n \times m = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_m)$, $\text{rank}(\mathbf{C}) = r$ and V_C the vector space generated by the columns of \mathbf{C} . The mean square for L_C is

$$S^2/r$$

with expected value

$$\sigma^2 + S_\mu^2/r.$$

Examples

1. Suppose that $\mathbf{y}: 4 \times 1 \sim (\boldsymbol{\mu}, \sigma^2 \mathbf{I})$ with

$$\boldsymbol{\mu} = \begin{pmatrix} \alpha \\ \alpha \\ \beta \\ \beta \end{pmatrix}$$

Let

$$\mathbf{C}_1 = \begin{pmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{pmatrix}, \mathbf{C}_2 = \begin{pmatrix} 1 & 0 \\ -1 & 0 \\ 0 & 1 \\ 0 & -1 \end{pmatrix} \text{ and } \mathbf{C} = (\mathbf{C}_1, \mathbf{C}_2).$$

The sums of squares of the linear sets L_{C_1} and L_{C_2} are

$$S_1^2 = \frac{1}{2}(y_1 + y_2)^2 + \frac{1}{2}(y_3 + y_4)^2$$

and

$$S_2^2 = \frac{1}{2}(y_1 - y_2)^2 + \frac{1}{2}(y_3 - y_4)^2.$$

respectively. The total sum of squares is

$$S^2 = y_1^2 + y_2^2 + y_3^2 + y_4^2 = S_1^2 + S_2^2.$$

The expected values are:

$$E(S_1^2) = 2\sigma^2 + 2\alpha^2 + 2\beta^2$$

$$E(S_2^2) = 2\sigma^2$$

$$E(S^2) = 4\sigma^2 + 2\alpha^2 + 2\beta^2.$$

2. Suppose that $\mathbf{y}: n \times 1 \sim (\boldsymbol{\mu}, \sigma^2 \mathbf{I})$. The set of all linear functions of \mathbf{y} is generated by $\mathbf{I}\mathbf{y}$, with sum of squares:

$$S^2 = \mathbf{y}'\mathbf{I}(\mathbf{I}'\mathbf{I})^*\mathbf{I}'\mathbf{y} = \mathbf{y}'\mathbf{y} = \sum_{i=1}^n y_i^2.$$

Note also that the linear functions y_1, y_2, \dots, y_n form a mutually orthogonal basis for this linear set. The sum of squares, S^2 , is therefore the same as the total of the individual sums of squares of the y_i 's, and the sum of squares of

$$y_i = (0, 0, \dots, 0, 1, 0, \dots, 0)\mathbf{y}$$

is

$$s_i^2 = y_i^2.$$

Let L_E be the set of linear functions orthogonal to \bar{y} . The sum of squares of L_E is

$$S_E^2 = \mathbf{y}'\mathbf{y} - n\bar{y}^2 = \sum_{i=1}^n (y_i - \bar{y})^2,$$

the mean square of L_E is

$$S_E^2/(n-1)$$

with expected value

$$\sigma^2 + \frac{1}{n-1} \sum_{i=1}^n (\mu_i - \bar{\mu})^2.$$

Also,

$$E(n\bar{y}^2) = \sigma^2 + n\bar{\mu}^2.$$