

# Contents

|           |                                                                                                                                          |           |
|-----------|------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>1</b>  | <b>Introduction</b>                                                                                                                      | <b>3</b>  |
| <b>2</b>  | <b>Literature</b>                                                                                                                        | <b>5</b>  |
| <b>3</b>  | <b>Preliminaries</b>                                                                                                                     | <b>8</b>  |
| 3.1       | $M/M/1$ queue                                                                                                                            | 8         |
| 3.2       | Non-preemptive priority and the PASTA property                                                                                           | 9         |
| 3.3       | $M/M/1+M$ queue (Erlang-A)                                                                                                               | 9         |
| 3.4       | Pollaczek–Khinchine formula                                                                                                              | 10        |
| 3.5       | Confluent hypergeometric function                                                                                                        | 10        |
| 3.6       | First-order linear ODEs and PDEs                                                                                                         | 11        |
| 3.6.1     | Integrating factor                                                                                                                       | 11        |
| 3.6.2     | Variation of parameters                                                                                                                  | 11        |
| 3.6.3     | Method of Characteristics                                                                                                                | 11        |
| <b>4</b>  | <b>Model description: Model-<math>X</math></b>                                                                                           | <b>12</b> |
| 4.1       | Balance equations                                                                                                                        | 14        |
| 4.1.1     | Balance equations for state space $S$                                                                                                    | 14        |
| 4.1.2     | Balance equations for state space $\tilde{S}$                                                                                            | 16        |
| 4.2       | Derivation of equations for (Partial) Probability Generating functions                                                                   | 17        |
| 4.2.1     | Equation for PGF in state space $S$                                                                                                      | 18        |
| 4.2.2     | Equation for PPGF in state space $\tilde{S}$                                                                                             | 27        |
| <b>5</b>  | <b>Analysis of Model-<math>A</math> (<math>\gamma_1 = \gamma_2 = 0, \theta_1 = \theta_2 = 0</math>)</b>                                  | <b>33</b> |
| 5.1       | Analysis of Model- $A$ : state space $S$                                                                                                 | 33        |
| 5.1.1     | Analytical approach for Model- $A$ in state space $S$                                                                                    | 34        |
| 5.1.2     | Probabilistic approach for Model- $A$ in state space $S$                                                                                 | 35        |
| 5.1.3     | Partial-PGF recurrence approach for Model- $A$ in state space $S$                                                                        | 36        |
| 5.2       | Analysis of Model- $A$ on $\tilde{S}$ : scope and limitations of the partial-PGF recursion                                               | 38        |
| 5.3       | Mean queue lengths and mean waiting times                                                                                                | 40        |
| 5.3.1     | Class-1 marginal PGF and mean queue length                                                                                               | 40        |
| 5.3.2     | Total queue length from the diagonal                                                                                                     | 40        |
| 5.3.3     | Class-2 mean queue length by subtraction                                                                                                 | 41        |
| 5.3.4     | Mean waiting times via Little’s Law                                                                                                      | 41        |
| <b>6</b>  | <b>Analysis of Model-<math>B</math> (<math>\gamma_1, \gamma_2 &gt; 0, \theta_1 = \theta_2 = 0</math>)</b>                                | <b>41</b> |
| <b>7</b>  | <b>Analysis of Model-<math>X</math> (<math>\gamma_1, \gamma_2 &gt; 0, \theta_1, \theta_2 &gt; 0</math>)</b>                              | <b>47</b> |
| <b>8</b>  | <b>Analysis of Model-<math>B_2</math> (<math>\gamma_1 &gt; 0, \gamma_2 = 0, \theta_1 = \theta_2 = 0</math>)</b>                          | <b>53</b> |
| 8.1       | Reduced fundamental equation                                                                                                             | 53        |
| 8.2       | Mean queue lengths and mean waiting times                                                                                                | 58        |
| <b>9</b>  | <b>Analysis of Model-<math>C_2</math> (<math>\theta_1 &gt; 0, \gamma_1 = \gamma_2 = \theta_2 = 0</math>)</b>                             | <b>59</b> |
| 9.1       | Setup: reduction to a first-order linear ODE in $x$                                                                                      | 61        |
| 9.2       | Probabilistic derivation of $P_y(y)$                                                                                                     | 61        |
| 9.3       | Analytical re-derivation of $P_y(y)$ via boundedness at $x = 1$                                                                          | 63        |
| 9.4       | Recovery of $P(x, y)$ from $P_y(y)$                                                                                                      | 64        |
| 9.5       | Mean queue lengths and mean waiting times                                                                                                | 66        |
| <b>10</b> | <b>Analysis of Model-<math>B_2^H</math> (<math>\gamma_1 &gt; 0</math> head-of-line, <math>\gamma_2 = \theta_1 = \theta_2 = 0</math>)</b> | <b>68</b> |
| 10.1      | Balance equations                                                                                                                        | 69        |
| 10.2      | Fundamental PGF equation                                                                                                                 | 69        |
| 10.3      | Closed-form solution                                                                                                                     | 70        |
| 10.3.1    | Analytical approach for Model- $B_2^H$                                                                                                   | 70        |
| 10.3.2    | Probabilistic approach for Model- $B_2^H$                                                                                                | 71        |
| 10.4      | Limits and sanity checks                                                                                                                 | 72        |
| 10.5      | Mean queue lengths and mean waiting times                                                                                                | 72        |
| 10.5.1    | Class-1 marginal PGF and mean queue length                                                                                               | 72        |
| 10.5.2    | Total queue length from the diagonal                                                                                                     | 72        |

|           |                                                                                                                                          |            |
|-----------|------------------------------------------------------------------------------------------------------------------------------------------|------------|
| 10.5.3    | Class-2 mean queue length by subtraction . . . . .                                                                                       | 72         |
| 10.5.4    | Mean waiting times via Little's Law . . . . .                                                                                            | 73         |
| <b>11</b> | <b>Analysis of Model-<math>C_2^H</math> (<math>\theta_1 &gt; 0</math> head-of-line, <math>\gamma_1 = \gamma_2 = \theta_2 = 0</math>)</b> | <b>73</b>  |
| 11.1      | Balance equations . . . . .                                                                                                              | 73         |
| 11.2      | Fundamental PGF equation . . . . .                                                                                                       | 74         |
| 11.3      | Closed-form solution . . . . .                                                                                                           | 74         |
| 11.3.1    | Analytical approach for Model- $C_2^H$ . . . . .                                                                                         | 75         |
| 11.3.2    | Probabilistic approach for Model- $C_2^H$ . . . . .                                                                                      | 76         |
| 11.4      | Limits and sanity checks . . . . .                                                                                                       | 76         |
| 11.5      | Mean queue lengths and mean waiting times . . . . .                                                                                      | 76         |
| 11.5.1    | Class-1 marginal PGF and mean queue length . . . . .                                                                                     | 76         |
| 11.5.2    | Class-2 mean queue length . . . . .                                                                                                      | 76         |
| 11.5.3    | Mean waiting times via Little's Law . . . . .                                                                                            | 77         |
| 11.6      | Comparison of the head-of-line models . . . . .                                                                                          | 77         |
| <b>12</b> | <b>Comparison of the Solved Models</b>                                                                                                   | <b>78</b>  |
| 12.1      | Traffic-intensity sweep . . . . .                                                                                                        | 80         |
| 12.2      | The jockeying mechanism (Model $B_2$ ) . . . . .                                                                                         | 83         |
| 12.3      | Class asymmetry . . . . .                                                                                                                | 85         |
| 12.4      | The abandonment mechanism and the throughput–quality tradeoff (Model $C_2$ ) . . . . .                                                   | 86         |
| 12.4.1    | The price in lost customers . . . . .                                                                                                    | 88         |
| 12.5      | Head-of-line versus full-rate mechanisms (Models $B_2^H$ and $C_2^H$ ) . . . . .                                                         | 90         |
| <b>13</b> | <b>Results</b>                                                                                                                           | <b>91</b>  |
| 13.1      | Validation . . . . .                                                                                                                     | 91         |
| 13.2      | Mean queue lengths and mean waiting times . . . . .                                                                                      | 94         |
| 13.3      | Head-of-line models $B_2^H$ and $C_2^H$ . . . . .                                                                                        | 95         |
| 13.4      | Cross-validation: simulation versus the CTMC . . . . .                                                                                   | 96         |
| 13.5      | Convergence to Model $A$ . . . . .                                                                                                       | 97         |
| 13.5.1    | Model $C_2$ as $\theta_1 \rightarrow 0^+$ . . . . .                                                                                      | 98         |
| 13.5.2    | Model $B_2$ as $\gamma_1 \rightarrow 0^+$ . . . . .                                                                                      | 99         |
| 13.5.3    | Instant-jockeying limit $\gamma_1 \rightarrow \infty$ . . . . .                                                                          | 99         |
| 13.5.4    | Head-of-line models $B_2^H$ and $C_2^H$ . . . . .                                                                                        | 100        |
| 13.6      | Comparative dashboard . . . . .                                                                                                          | 101        |
| <b>14</b> | <b>Conclusion and Future Work</b>                                                                                                        | <b>101</b> |
| 14.1      | The obstruction taxonomy . . . . .                                                                                                       | 101        |
| 14.2      | The mechanism dichotomy . . . . .                                                                                                        | 102        |
| 14.3      | Operational reading and the deterioration-direction caveat . . . . .                                                                     | 102        |
| 14.4      | Limitations . . . . .                                                                                                                    | 102        |
| 14.5      | Future work . . . . .                                                                                                                    | 103        |

# Exact Generating-Function Analysis of a Two-Class Non-Preemptive Priority $M/M/1$ Queue with Jockeying and Abandonment

Victor Dominguez Sainz

June 12, 2026

## Abstract

We study the stationary queue-length distribution of a two-class, non-preemptive priority  $M/M/1$  queue augmented with two waiting-room mechanisms: *jockeying*, in which a waiting customer switches queue, and *abandonment*, in which a waiting customer leaves the system. Jockeying conserves the total number of customers, whereas abandonment does not, and combining both yields a model (Model- $X$ ) that is analytically intractable. We therefore organise the analysis as a hierarchy that isolates one mechanism at a time: the baseline Model- $A$  (no mechanism); Model- $B$  (bidirectional jockeying); the full Model- $X$ ; and the tractable specialisations Model- $B_2$  (one-way jockeying, class 1 to class 2) and Model- $C_2$  (class-1 abandonment). We also treat *head-of-line* variants  $B_2^H$  and  $C_2^H$ , in which a mechanism acts only on the customer at the head of the priority queue.

The central object is the bivariate probability generating function (PGF)  $P(x, y)$  of the two queue lengths, obtained by analytical and, where the structure permits, probabilistic arguments. Model- $A$  is classical and follows from the kernel method; the contribution of this work is the unified functional-equation framework built around it, within which each solvable model is closed by a single pinning principle — a kernel root inside the unit disc for Model- $A$ , analyticity at an interior point for Model- $B_2$ , and boundedness on the unit-circle boundary for Model- $C_2$  — while the head-of-line restriction collapses the functional equation to an algebraic one and yields rational PGFs in closed form. Two models are left open, with their obstructions named: Model- $B$  reduces the class-2 boundary function to a Volterra-type integral equation, and Model- $X$  compounds this with transcendental characteristics, so that even the kernel root ceases to be available in closed form. For Model- $A$  we additionally develop the partial-PGF recursion on an alternative total-count state space and characterise it as a heuristic approximation, accurate in the priority-dominated, moderate-load regime and quantified there against the exact solution.

Every closed form is validated against a truncated continuous-time Markov chain and recovers the baseline as the mechanism rates vanish. The comparison yields the central applied finding: the two mechanisms are not interchangeable, for jockeying merely redistributes delay between the classes at fixed total work, whereas abandonment shortens every queue only by shedding a substantial fraction of priority demand — nearly one class-1 arrival in four at moderate load — so that its apparent waiting-time gains are in part attrition rather than faster service.

## 1 Introduction

### Motivation

Priority queues arise naturally in a wide range of contexts where a common server must satisfy heterogeneous demand. The primary motivation for this thesis comes from the context of healthcare operations. In particular, the management of waiting lists for access to elderly care facilities, where the medical severity dictates a user’s —hereafter referred to as *customer*— priority class. In this context, prolonged waiting times may alter the system’s dynamics, given that the lack of appropriate care may deteriorate the health condition of a waiting potential user, leading to a change in their priority status (jockeying) or leaving the system (abandonment), be it due to seeking alternative care or passing away. Despite the initial motivation, this work aims to address the problem from a general modelling, which may be applicable to a broad class of systems: characterising the long-run behaviour of queueing systems when phenomena such as jockeying and/or abandonments play a role.

In this thesis, an analysis of a 2-class  $M/M/1$  queueing system with non-preemptive priority is carried out. In short, this means that arrivals from each priority class are driven by a Poisson process with arrival rates  $\lambda_1$  and  $\lambda_2$ , exponentially distributed service times at a rate  $\mu$  —common to both priority classes—, and a single server that completes the current service despite the arrivals of higher-priority customers; with the addition of the mechanisms of jockeying and abandonments. The model and its mechanisms are described in more depth in §4.

The interplay of these two mechanisms within a priority discipline and how they affect the difficulty of obtaining a closed-form expression for their Probability Generating Function (PGF) are the pivotal objects of this thesis.

## The analytical challenge

The combination of jockeying and abandonment in full generality defines Model- $X$ . The bivariate PGF satisfies a functional equation that involves terms of three kinds: algebraic coefficients, a partial derivative term in  $x$  —from jockeying and abandonments in class-1— and a partial derivative term in  $y$  —from the same mechanisms in class-2. When derivative terms of the two kinds are present, the equation belongs to a family of integro-differential equations for which no close-form solution is known, and it becomes a substantially challenging problem.

The strategy adopted here is to explore less complex models that require a use of less advanced mathematical tools by setting some of the jockeying and/or abandonment parameters equal to zero. The table below (Table 1) lists the models studied in this work and provide some insight of the challenge they imply:

Table 1: Models studied in this thesis, with the active mechanism, the resulting functional-equation type, and the solution status. The head-of-line variants  $B_2^H$  and  $C_2^H$  restrict the mechanism to the customer at the head of Queue 1, which collapses the derivative term to an algebraic one and renders the model solvable in closed form.

| Model   | Active mechanism                        | Equation type                       | Status |
|---------|-----------------------------------------|-------------------------------------|--------|
| A       | none                                    | algebraic PGF equation              | solved |
| B       | bidirectional jockeying                 | PDE, Volterra family                | open   |
| X       | jockeying and abandonment               | PDE, transcendental characteristics | open   |
| $B_2$   | one-way jockeying ( $1 \rightarrow 2$ ) | ODE in $x$ , Kummer                 | solved |
| $B_2^H$ | one-way jockeying, head-of-line         | algebraic PGF equation              | solved |
| $C_2$   | class-1 abandonment                     | ODE in $x$ , Kummer                 | solved |
| $C_2^H$ | class-1 abandonment, head-of-line       | algebraic PGF equation              | solved |

As one can see, some models —in particular, Model- $X$  and Model- $B$ — have been left incomplete, as they involve derivative terms in  $y$ , and some other models such a potential Model- $B_1$ , Model- $C$  or Model- $C_1$  have not even been considered for the same reason: they all reduce to a Volterra integral equation that falls beyond the scope of the work carried out in this thesis. Here Model- $B$  serves as an example of how this sort of problems reduce to such an integral equation, whereas the other models correspond to problems we can solve. The last two entries, Models  $B_2^H$  and  $C_2^H$  (§§10 and 11), are *head-of-line* simplifications in which only the customer at the head of Queue 1 may jockey or abandon: the mechanism then acts at the constant rate  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  or  $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$  rather than the length-proportional  $\gamma_1 n_1$  or  $\theta_1 n_1$ , so the fundamental equation stays algebraic and is solved by the Kernel Method exactly as in Model- $A$ .

## Methods

Three complementary proof strategies are used throughout.

*Kernel method / ODE with integrating factor.* For Model- $A$ , which does not involve derivative terms whatsoever, the algebraic coefficient of  $P(x, y)$  vanishes on a curve (i.e. the *kernel*). Evaluating the equation there lets us determine the rest of the unknowns. Model- $B_2$  and Model- $C_2$ , which do involve derivative terms in  $x$ , reduce to a first-order linear ODE in  $x$  and admit an explicit integrating factor. The unknowns there are obtained by means of a regularity argument at a singularity point of the integrating factor. Both strategies are here explained as one, as the underlying idea is the same. One could say that the latter is the extension of the former when the unknowns are independent from the derivative terms and can be taken outside of the integral.

*Probabilistic argument via Pollaczek–Khinchine and the effective service time.* For Model-*A* and Model-*C*<sub>2</sub>, we have Poisson arrivals for class-2 customers —and therefore the PASTA property holds—, which allows us to utilise the Pollaczek–Khinchine formula, combined with the *effective service time* seen by class-2 customers: an  $M/G/1$  queueing system where the general service times are equivalent to the busy period of the class-1 subsystem, which is an  $M/M/1$  queueing system for Model-*A* and an  $M/M/1 + M$  system for Model-*B*. In both cases, the probabilistic argument confirms the results obtained analytically, allowing to determine the unknown in an independent way. Since jockeying destroys the Poisson structure, this approach is not valid for the other models we studied.

*Partial PGF (PPGF) in the alternative state space  $\tilde{S}$ .* For the baseline Model-*A* we create an alternative state space that looks at the model by taking the number of class-2 customers —difficult distribution— and the total amount of customers —easy, since the total number of customers is preserved and is basically a regular  $M/M/1$  queueing system. This yields a PPGF with a non-homogeneous term. For Model-*A*, a recursion argument derived by Cohen [Coh56] applied to the regular state space. We explore, without much success, the challenges of such an approach on this alternative space and evaluate whether a rough approximation is feasible using Cohen’s trick.

## Structure of the thesis

The sections are ordered to follow the analytical arc baseline → obstruction → synthesis → tractable specialisations → head-of-line collapse → comparison → validation. Section 2 reviews the relevant literature and Section 3 collects the probabilistic and analytic background used throughout this work. Section 4 defines the full model, Model-*X*, and derives the general balance equations and the fundamental PGF equations, both for the regular and the alternative state space. The analysis then proceeds from baseline to synthesis: Section 5 solves the baseline Model-*A* in closed form; Section 6 shows that bidirectional jockeying (Model-*B*) reduces the problem to a Volterra-type integral equation and is left open; and Section 7 returns to Model-*X* to show that the full model inherits both obstructions at once and is likewise left open. Sections 8 and 9 then recover tractability in the one-way-jockeying (Model-*B*<sub>2</sub>) and class-1-abandonment (Model-*C*<sub>2</sub>) specialisations, while Sections 10 and 11 show how the head-of-line variants, Models  $B_2^H$  and  $C_2^H$ , collapse the functional equation to an algebraic one solved in closed form. Finally, Section 12 compares the solved models, Section 13 presents the numerical validation and performance metrics, and Section 14 concludes by abstracting the obstruction taxonomy and setting out the future work.

## 2 Literature

The study of two-class priority queues with departure mechanisms sits at the intersection of three well-developed streams: the exact generating-function analysis of non-preemptive priority queues, the theory of queues with customer impatience (abandonment), and the theory of jockeying and dynamic priority changes. The applied motivation throughout is the management of access to care — elderly-care waiting lists, emergency departments, and transplant registries — in which the two mechanisms studied here carry direct clinical readings: a waiting patient whose condition deteriorates changes priority class (jockeying), while one who leaves the list by death or transfer reneges (abandonment). This thesis contributes an exact, single-server treatment that admits jockeying and abandonment within a unified bivariate-PGF framework, filling a gap that the multi-server and approximate literature has so far left open.

### Non-preemptive priority queues: exact queue-length analysis

The direct methodological ancestor of this work is Cohen [Coh56], who studied a single server loaded by two independent Poisson streams and introduced the partial-generating-function recursion — the “Cohen trick” — that resolves the lower-priority queue length by transforming only one coordinate of the joint distribution while keeping the other discrete. That device is exactly the one reused, in modern notation, in the partial-PGF analysis of Model-*A* on the total-count state space (§5.2). The stationary queue-length law of the two-class non-preemptive  $M/M/1$  priority queue — the content of Theorem 1 (Model-*A*) — is in this sense classical, recoverable by several routes; the contribution of the present work is not that result in isolation but the *unified framework* built around it, into which jockeying and abandonment are introduced one mechanism at a time and solved by a common functional-equation method.

The most recent exact closed-form treatment in this line is that of Zuk & Kirszenblat [ZK23], who obtain the joint and marginal queue-length distributions of a non-preemptive  $M/M/c$  queue with two priority classes and a common service rate by generating-function methods and a quadratic recurrence; their single-server specialisation ( $c = 1$ ) coincides with Model-A. A complementary strand studies dynamic rather than static priority: Stanford et al. [STZ14] analyse the *accumulating-priority*  $M/G/1$  queue, in which each class earns priority linearly in its waiting time so that older customers eventually overtake newer ones, and derive exact waiting-time distributions through a maximum-priority process and Laplace–Stieltjes transforms. Accumulating priority is a continuous-time analogue of the discrete, rate- $\gamma_i$  jockeying studied here; like the present work it is exact and single-server, but its object is the sojourn time rather than the bivariate queue-length PGF.

## Priority queues with abandonment

Abandonment enters queueing theory through the Erlang-A ( $M/M/c + M$ ) model, in which each waiting customer holds an independent exponential patience clock; the modern treatment and its heavy-traffic scaling are due to Garnett et al. [GMR02], and the single-server case  $M/M/1 + M$  supplies the busy-period structure on which Model- $C_2$  rests (§3.3). Exact stationary results for priority queues *with* reneging are comparatively scarce and almost always multi-server: Jouini & Roubos [JR14] treat an  $M/M/s + M$  queue with two non-preemptive classes and per-class exponential patience, deriving Laplace–Stieltjes transforms for conditional and unconditional waiting times together with the probabilities of service and of abandonment — the multi-server analogue of Model- $C_2$ , but with the waiting time, not the queue-length PGF, as its object. Where exact transforms are unavailable the literature turns to fluid and diffusion approximations or to many-server asymptotics, in which the Erlang-A stationary law and its descendants are expressed through confluent hypergeometric and incomplete-gamma functions [ZM05]; the same special-function structure appears exactly, and at finite scale, in the closed form for Model- $C_2$ . The exact multi-server priority results that do exist carry no class-asymmetric reneging — those of Wang et al. [WBSW15] and Zuk & Kirszenblat [ZK23] are abandonment-free, and Baron et al. [BLW25] replace patience with an admission threshold — so that Model- $C_2$  appears to be the first exact, single-server, queue-length treatment of *class-asymmetric* abandonment, with impatience acting on the priority class alone. The clinical reading that licenses this asymmetry is long-standing: Zenios [Zen99] models the organ-transplant waiting list as a queue whose reneging is patient mortality, precisely the regime in which losing high-priority demand is the outcome of interest.

## Jockeying and priority changes

The closest predecessor on the jockeying side is de Waal [dW92], who studies an  $M/G/1$  priority queue in which impatient customers *change priority class* while waiting. Its entry in Table 2 is marked approximate not by choice of method but by necessity: with general service times the governing equations do not close, and the analysis delivers Laplace–Stieltjes-based *approximations* of the waiting-time distributions rather than an exact stationary law. The healthcare incarnation of the same idea is the model of Hu, Chan & Dong [HCD22], who let customers transition between a moderate and an urgent class as their condition deteriorates or improves and characterise a near-optimal  $c\mu/\theta$ -type scheduling index in heavy traffic; the system is multi-server and its results are asymptotic, so it too is approximate in the sense of Table 2. Control-theoretic treatments of the same family, such as the operator-initiated holding and abandonment policies of Shmelev & Zychlinski [SZ25], likewise live in the multi-server, heavy-traffic regime and target scheduling optimality rather than exact distributions. Against this backdrop Model- $B_2$  is the exact, single-server counterpart of the approximate multi-server “priority-change” literature: it retains the deterioration mechanism — a waiting class-1 customer relocating to the class-2 queue at rate  $\gamma_1$  — but pays for exactness with the single-server, Markovian, one-directional restriction, returning a closed-form boundary function in place of an approximation.

## Multi-server extensions and performance design

The remaining references concern multi-server systems studied for performance and design rather than for exact single-server distributions, and they bound the present work from the side of scale. Wang et al. [WBSW15] give the exact queue-length PGF of an  $M/M/c$  queue with two *preemptive* priority classes and class-dependent service rates, a useful contrast to the non-preemptive, common-rate discipline used throughout this thesis. Baron et al. [BLW25] study capacity rationing in an

$M/M/c$  non-preemptive priority queue, in which low-priority customers are admitted to service only when a threshold number of servers is idle, and show that the optimal threshold scales as  $O(\sqrt{c})$  in the Halfin–Whitt regime; rationing is a third kind of inter-class coupling, neither jockeying nor abandonment, noted here to delimit what this thesis does not attempt. Together these works confirm that exact, mechanism-rich analysis is essentially a single-server phenomenon: once  $c > 1$  is combined with a departure mechanism, the literature moves to approximations and asymptotics.

## Methodological lineage

The functional-equation technique used in Sections 8 and 9 belongs to the kernel-method and boundary-value tradition for two-dimensional Markovian queues. The systematic theory — reducing a bivariate balance equation to the curve on which its kernel vanishes, and from there to a boundary-value problem for the unknown boundary functions — is developed by Fayolle, Iasnogorodski & Malyshev [FIM99] for random walks in the quarter-plane. The pinning arguments of this thesis are the elementary, one-dimensional residue of that programme: in Model- $B_2$  the boundary function is fixed by demanding *analyticity* of the PGF at the interior point where the kernel root meets the diagonal (§8), and in Model- $C_2$  by demanding mere *boundedness* at the unit-disc boundary (§9); both select the analytic solution of the reduced equation, exactly the role the boundary conditions play in the general theory. Two narrower tools recur. The confluent hypergeometric (Kummer) function that carries the Model- $C_2$  boundary function is the same special function through which the Erlang-A and reneging-queue stationary laws are expressed [ZM05, DLM]; and the selection of the decaying solution of the three-term recurrence behind the partial-PGF analysis is governed by Pincherle’s theorem, for which we cite the recurrence-relation treatment of Gautschi [Gau67] alongside the DLMF [DLM].

## Position of this thesis

Table 2: Literature overview: models and mechanisms. ✓ = present, – = absent,  $\approx$  = approximate only.

| Reference                                   | Servers  | Non-preem. | Jockeying | Aband. | Queue-len. | Exact     |
|---------------------------------------------|----------|------------|-----------|--------|------------|-----------|
| Cohen (1956)                                | 1        | ✓          | –         | –      | ✓          | ✓         |
| de Waal (1992)                              | 1        | ✓          | ✓         | –      | –          | $\approx$ |
| Jouini & Roubos (2014)                      | $c$      | ✓          | –         | ✓      | –          | ✓         |
| Wang et al. (2015)                          | $c$      | –          | –         | –      | ✓          | ✓         |
| Stanford et al. (2014)                      | 1        | ✓          | ✓         | –      | –          | ✓         |
| Zuk & Kirszenblat (2023)                    | $c$      | ✓          | –         | –      | ✓          | ✓         |
| Baron et al. (2025)                         | $c$      | ✓          | –         | –      | –          | ✓         |
| Hu, Chan & Dong (2022)                      | $c$      | ✓          | ✓         | ✓      | –          | $\approx$ |
| Shmelev & Zychlinski (2025)                 | $c$      | ✓          | –         | ✓      | –          | $\approx$ |
| <b>This thesis (Model-A)</b>                | <b>1</b> | ✓          | –         | –      | ✓          | ✓         |
| <b>This thesis (Model-<math>B_2</math>)</b> | <b>1</b> | ✓          | ✓         | –      | ✓          | ✓         |
| <b>This thesis (Model-<math>C_2</math>)</b> | <b>1</b> | ✓          | –         | ✓      | ✓          | ✓         |

Table 2 places this work against its closest neighbours, and reading it by row makes the gap precise. The Cohen (1956) row is the single-server, exact, non-preemptive queue-length baseline with neither extra mechanism — the ancestor of Model-A. de Waal (1992) adds jockeying (priority change) at a single server, but only approximately and for waiting times rather than queue lengths. Jouini & Roubos (2014) add abandonment exactly, but at  $c$  servers and again for waiting times. Wang et al. (2015) is exact and queue-length-focused, yet multi-server, preemptive, and mechanism-free. Stanford et al. (2014) is single-server and exact with a dynamic-priority (jockeying-like) mechanism, but its object is the sojourn time. Zuk & Kirszenblat (2023) is the exact multi-server queue-length result whose  $c = 1$  case is Model-A, with neither jockeying nor abandonment. Baron et al. (2025) is exact and multi-server, but couples the classes through admission control rather than jockeying or abandonment, and reports no queue-length law. Hu, Chan & Dong (2022) and Shmelev & Zychlinski (2025) carry, respectively, jockeying-as-deterioration and controlled abandonment, but



multi-server and only approximately. No row above combines, at a single server, the exact bivariate queue-length PGF with either jockeying or abandonment — which is precisely the column pattern of the three “This thesis” rows. The gap can therefore be stated in one sentence: prior to this work there is no exact, single-server, queue-length analysis that places non-preemptive priority together with jockeying or abandonment inside a single PGF framework.

The non-claims must be stated with equal precision. The framework is unified but not complete: the full model (Model-*X*, §7), in which jockeying and abandonment act together, and the bidirectional-jockeying model (Model-*B*, §6) are left open, each reducing to a Volterra-type integral equation that this thesis characterises but does not solve in closed form. The exact contributions are confined to the three “This thesis” rows of Table 2 — the baseline Model-*A*, the one-way jockeying Model-*B*<sub>2</sub>, and the class-1 abandonment Model-*C*<sub>2</sub>, together with their head-of-line simplifications. The claim is exactness and unification at a single server with one mechanism active at a time; it is explicitly not a solution of the combined or bidirectional problems, which remain, as in the multi-server literature, beyond exact reach.

### 3 Preliminaries

This section collects background material referenced throughout the derivations. Introducing these building blocks here allows the subsequent arguments to remain focused on their primary objectives.

The central mathematical object in our derivations will be the (P)PGF of the model in question, as well as the idle probability of the system for some corollaries.

#### 3.1 $M/M/1$ queue

The model described here is the canonical  $M/M/1$  queue, well studied in multiple textbooks and manuals and considered the absolute foundation of queueing theory. Here, we will use [AR02] as our reference, but there is a lot of good quality literature and sources about it. Customers arrive at the system according to a Poisson process —i.e., the inter-arrival times are exponentially distributed with parameter  $\lambda$ —, service times are exponentially distributed with parameter  $\mu$ , and a single server operates in first-come-first-serve (FCFS) —also known as first-in-first-out (FIFO)— order. The system is stable if and only if the system’s load  $\rho = \lambda/\mu < 1$ .

Let  $\pi_n$  be the limiting probability of finding  $n$  customers in the system. Under stability, the detailed balance equations  $\lambda \pi_n = \mu \pi_{n+1}$  ( $n \geq 0$ ) together with the normalisation  $\sum_{n=0}^{\infty} \pi_n = 1$  give us the geometric steady-state distribution of the number of customers in the system:

$$\pi_n = (1 - \rho) \rho^n, \quad n \geq 0. \quad (1)$$

By the PASTA property (§3.2), this also corresponds to the distribution seen by the arriving customers. Let  $L$  be the random variable that tells us how many customers—including the one that might be in service—are in the system. The expected number of customers in the system is the following:

$$\mathbb{E}[L] = \frac{\rho}{1 - \rho}.$$

From this, the expected sojourn time  $W$ —the time that elapses between a customer’s arrival and their departure after service completion—is derived by Little’s Law ( $\mathbb{E}[L] = \lambda \mathbb{E}[W]$ ):

$$\mathbb{E}[W] = \frac{\mathbb{E}[L]}{\lambda} = \frac{1/\mu}{1 - \rho}.$$

Another concept, more rare but highly relevant for the content that will be discussed in this thesis, is the Busy Period (*BP*) of the system, i.e., the time elapsed between the arrival of a customer to an idle system and the departure of a customer who leaves the system empty again after service completion. The Laplace-Stieltjes Transform (LST) and the expectation of this system’s busy period are known:

$$\widetilde{BP}(s) = \frac{(\mu + \lambda + s) - \sqrt{(\mu + \lambda + s)^2 - 4\mu\lambda}}{2\lambda}, \quad (2)$$

$$\mathbb{E}[BP] = \frac{1}{\mu(1 - \rho)}. \quad (3)$$



### 3.2 Non-preemptive priority and the PASTA property

Throughout this thesis the two customer classes are served under a *non-preemptive* (head-of-line) priority discipline: whenever the server becomes free it admits a waiting class-1 customer if any are present, and a class-2 customer only otherwise, but a customer already in service is *never* interrupted—a class-1 arrival that finds a class-2 customer in service waits until that service completes. Priority therefore governs only the order in which the queues are drained, not the service in progress. Because the service rate  $\mu$  is common to both classes, the *memorylessness* of the exponential service time makes the residual service of the customer in progress  $\text{Exp}(\mu)$  regardless of its class; hence the class of the in-service customer does not influence the future evolution of the queue lengths, which is why a single in-service customer of either class suffices to mark the server busy.

We also invoke repeatedly the *PASTA* property (*Poisson Arrivals See Time Averages*) [AR02]: if a system is fed by a Poisson arrival process that satisfies the *lack-of-anticipation assumption* (at each instant the future increments of the arrival process are independent of the current system state), then the long-run fraction of arrivals that find the system in a given state equals the stationary probability of that state, so the distribution seen by an arriving customer coincides with the time-stationary one. Both hypotheses hold here: the exogenous class- $i$  arrivals are independent Poisson processes at rates  $\lambda_i$ , and they are independent of the system's internal exponential service, jockeying and abandonment clocks, so they cannot anticipate the future. By Poisson *superposition* the merged arrival stream is itself Poisson at rate  $\lambda = \lambda_1 + \lambda_2$ , so PASTA applies to the aggregate state; and since each class stream is an independent Poisson process—equivalently, an i.i.d.  $\lambda_i/\lambda$  *thinning* of the merged stream—PASTA applies to each class separately.

### 3.3 $M/M/1+M$ queue (Erlang-A)

Another relevant queueing model for this thesis is the so-called  $M/M/1+M$  model, also known as the *Erlang A* model. It is based on the  $M/M/1$  queue—Poisson arrivals at rate  $\lambda$  and exponentially distributed service times with rate  $\mu$ —but adds customer impatience: each *waiting* customer (the one in service is not impatient) independently abandons after an exponentially distributed patience time with rate  $\theta$ . The key difference of this model with respect to the  $M/M/1$  is that the  $M/M/1+M$  does not require  $\rho = \lambda/\mu < 1$  to be stable: it is positive recurrent for all  $\lambda, \mu, \theta > 0$ .

The steady state distribution follows, naturally, from its balance equations. At state  $n \geq 0$ —for  $n$  being the number of customers in the *system*—we have

$$\lambda \pi_n = (\mu + n\theta) \pi_{n+1}, \quad n \geq 0. \quad (4)$$

Iterating yields

$$\pi_n = \pi_0 \frac{(\lambda/\theta)^n}{(\mu/\theta)^{(n)}} \quad n \geq 0; \quad \text{where} \quad a^{(n)} = a(a+1) \cdots (a+n-1) \quad (5)$$

Normalisation gives the non-trivial idle probability  $\pi_0 = [{}_1F_1(1; \mu/\theta; \lambda/\theta)]^{-1}$  (see §3.5 for the Kummer function  ${}_1F_1$ ).

The *Busy Period* of the  $M/M/1+M$  defined here is far more complex than the one expressed in the  $M/M/1$  queueing system. We record it for the class-1 subsystem used in Model- $C_2$ , taking arrival rate  $\lambda_1$ , service rate  $\mu$  and abandonment rate  $\theta_1$ , so that the relevant idle probability is  $\pi_0 = [{}_1F_1(1; \mu/\theta_1; \lambda_1/\theta_1)]^{-1}$ . Writing  $B_C$  for this class-1 busy period, its LST and expectation are

$$\tilde{B}_C(s) = \frac{\mu}{\lambda_1 + \mu + s - \frac{\lambda_1(\mu + \theta_1)}{\lambda_1 + \mu + \theta_1 + s - \frac{\lambda_1(\mu + 2\theta_1)}{\lambda_1 + \mu + 2\theta_1 + s - \cdots}}}, \quad (6)$$

$$\mathbb{E}[B_C] = \frac{1}{\lambda_1} \left( {}_1F_1(1; \mu/\theta_1; \lambda_1/\theta_1) - 1 \right) \quad (7)$$

The continued fraction (6) is generated by a first-passage recursion. Let  $\phi_n(s)$  denote the LST of the first-passage (sub-busy-period) time from  $n$  customers in the system down to  $n-1$ . Conditioning on the first transition out of state  $n$ —an arrival (rate  $\lambda_1$ ) to state  $n+1$ , or a departure by service completion or abandonment (combined rate  $\mu + (n-1)\theta_1$ : one customer in service plus  $n-1$  waiting, each impatient at rate  $\theta_1$ ) to state  $n-1$ —the strong Markov property gives

$$\phi_n(s) = \frac{\mu + (n-1)\theta_1}{\lambda_1 + \mu + (n-1)\theta_1 + s - \lambda_1 \phi_{n+1}(s)}, \quad n \geq 1, \quad \tilde{B}_C(s) = \phi_1(s). \quad (8)$$

Unrolling (8) from  $n = 1$  reproduces the continued fraction (6) term by term. The continued fraction also sums in closed form, and the argument is short enough to give in full. Fix  $s > 0$  and scale the rates by  $\theta_1$ , setting

$$a = \frac{\mu}{\theta_1}, \quad b = \frac{s}{\theta_1}, \quad c = \frac{\lambda_1}{\theta_1},$$

so that (8) reads  $\phi_n(s) = (a + n - 1) / ((a + n - 1) + b + c - c \phi_{n+1}(s))$ . Consider the integrals

$$I(p) = \int_0^1 t^p (1-t)^{b-1} e^{-ct} dt, \quad p > -1, \quad (9)$$

which by the Euler representation (15) are Beta-function multiples of Kummer values,  $I(p) = B(p+1, b) {}_1F_1(p+1; p+1+b; -c)$ . Integrating the exact derivative  $\frac{d}{dt} [t^p (1-t)^b e^{-ct}]$  over  $[0, 1]$  — the boundary terms vanish for  $p, b > 0$  — and splitting  $(1-t)^b = (1-t)^{b-1} - t(1-t)^{b-1}$  in the two terms where it appears yields the three-term recurrence

$$(p+b+c)I(p) = pI(p-1) + cI(p+1), \quad (10)$$

a relation of contiguous type for  ${}_1F_1$  (obtainable equivalently by combining the contiguous relations of [DLM, §13.3]). Dividing (10) at  $p = a + n - 1$  by  $I(a + n - 2)$  shows that the ratios  $R_n := I(a + n - 1) / I(a + n - 2)$  satisfy exactly the recursion of  $\phi_n$ . Two further facts pin the identification. First, by Laplace's method at the endpoint  $t = 1$ ,  $I(p) \sim e^{-c} \Gamma(b) p^{-b}$  as  $p \rightarrow \infty$ : the solution  $\{I(p)\}$  decays only polynomially, whereas the dominant solution of (10) grows superexponentially (its consecutive ratio grows like  $p/c$ ), so  $\{I(p)\}$  is the *minimal (recessive)* solution. Second, by *Pincherle's theorem* the convergent continued fraction equals precisely the ratio built from the minimal solution. Hence

$$\tilde{B}_C(s) = \phi_1(s) = \frac{I(a)}{I(a-1)} = \frac{B(a+1, b) {}_1F_1(a+1; a+1+b; -c)}{B(a, b) {}_1F_1(a; a+b; -c)}, \quad (11)$$

a contiguous ratio of the form (16). This identity is used, with  $s = \lambda_2(1-y)$ , in the boundedness route for Model- $C_2$  (§9, Equation (266)).

### 3.4 Pollaczek–Khinchine formula

The  $M/G/1$  queue [AR02] is a single-server queue with Poisson arrivals at rate  $\lambda$  and generally distributed i.i.d. service times  $B$  with LST  $\tilde{B}(s) = \mathbb{E}[e^{-sB}]$ . For stability purposes, the system load  $\rho_B = \lambda \mathbb{E}[B] < 1$  is required (written  $\rho_B$  to keep the symbol  $\rho$  for the two-class load of this thesis). Under these two conditions — Poisson arrivals and stability — the Pollaczek–Khinchine (PK) formula gives the PGF of the stationary number of customers in the system:

$$L(z) = \frac{(1-\rho_B)(1-z)\tilde{B}(\lambda(1-z))}{\tilde{B}(\lambda(1-z)) - z}. \quad (12)$$

### 3.5 Confluent hypergeometric function

The *Kummer confluent hypergeometric function of the first kind* will come up in some of the derivations of this thesis, and it reads as follows:

$${}_1F_1(a; b; z) = \sum_{n=0}^{\infty} \frac{a^{(n)}}{b^{(n)} n!} z^n, \quad a^{(n)} = \prod_{k=0}^{n-1} (a+k), \quad a^{(0)} = 1, \quad (13)$$

where  $b$  is not 0 or a negative integer —i.e.  $b \notin \{0, -1, -2, \dots\}$ — and the series converges for all  $z \in \mathbb{C}$ . The function  ${}_1F_1(a; b; z)$  is the unique solution of *Kummer's equation* [DLM]:

$$z u'' + (b-z) u' - a u = 0, \quad (14)$$

normalised by  ${}_1F_1(a; b; 0) = 1$ . The *Kummer series* (Equation (13)) converges for all  $z \in \mathbb{C}$  and  $b \notin \{0, -1, -2, \dots\}$ .

For this thesis, two identities related to it are used. First, the *Euler integral representation*, for  $a, b > 0$  we have:

$${}_1F_1(a; a+b; z) = \frac{1}{B(a, b)} \int_0^1 t^{a-1} (1-t)^{b-1} e^{zt} dt, \quad (15)$$

where  $B$  is the *Beta function*  $B(a, b) = \Gamma(a)\Gamma(b)/\Gamma(a+b)$ , and  $\Gamma$  is the *Gamma function*. Secondly, *contiguous ratios*: ratios of two  ${}_1F_1$  values whose parameters differ by integer shifts. Two patterns arise in this thesis. The *simultaneous shift*

$$\frac{{}_1F_1(a+1; a+1+b; z)}{{}_1F_1(a; a+b; z)}, \quad (16)$$

in which both parameters move by one so that their difference  $b$  is preserved, is by the Euler representation (15) a ratio of two weighted Beta integrals differing only in the exponent of  $t$ ; it appears as the closed-form value of the Erlang-A busy-period continued fraction (§3.3, Equation (11)) and hence in the boundary function of Model- $C_2$ . The *first-parameter shift*  ${}_1F_1(a+1; b; z)/{}_1F_1(a; b; z)$ , with the second parameter fixed, appears instead in the boundary function of Model- $B_2$  (§8).

### 3.6 First-order linear ODEs and PDEs

Since jockeying and abandonment add state-dependent rates to the balance equations, derivative terms end showing up in the derivations and they require solving first-order linear ODEs and PDEs for the PGF. The three methods below are invoked at specific steps of the analysis.

#### 3.6.1 Integrating factor

This method is standard to solve —for each fixed  $y$ — a *first-order linear ODE in  $x$* . Such an equation has standard form

$$\frac{dP}{dx} + p(x; y)P = q(x; y). \quad (17)$$

The *integrating factor*  $\mu_I(x; y) = \exp(\int_0^x p(t; y) dt)$  converts (17) to  $\frac{d}{dx}[P \cdot \mu_I] = q \cdot \mu_I$ . Integrating from 0 to  $x$  gives the general solution

$$P(x, y) = \frac{1}{\mu_I(x; y)} \left[ C_0(y) + \int_0^x q(t; y) \mu_I(t; y) dt \right], \quad (18)$$

where  $C_0(y)$  is an arbitrary function of  $y$ , determined by a boundary condition.

#### 3.6.2 Variation of parameters

*Variation of parameters* provides an alternative derivation of (18) that makes the role of the free function explicit. The homogeneous equation  $\frac{dP}{dx} + p(x; y)P = 0$  has general solution  $P_h(x; y) = C/\mu_I(x; y)$  for an arbitrary constant  $C$ . To solve the full equation, one *varies* that constant, setting  $P(x, y) = C(x; y)/\mu_I(x; y)$  and substituting into (17). Since the homogeneous part cancels, one obtains the following.

$$\frac{dC}{dx}(x; y) = q(x; y) \mu_I(x; y), \quad (19)$$

and integrating recovers (18) with  $C_0(y)$ .

#### 3.6.3 Method of Characteristics

This method is standard for solving a *first-order linear PDE in both  $x$  and  $y$*  simultaneously. It reduces the PDE to a family of ODEs along the so-called *characteristic curves* in the  $(x, y)$ -plane; along each curve, the PDE collapses to a single-variable ODE.

A PDE of the form

$$a \frac{\partial P}{\partial x} + b \frac{\partial P}{\partial y} = cP + d \quad (20)$$

is solved by tracing the characteristic curve  $(x(t), y(t))$  satisfying

$$\frac{dx}{dt} = a, \quad \frac{dy}{dt} = b. \quad (21)$$

By the chain rule,  $\frac{d}{dt}P(x(t), y(t)) = aP_x + bP_y$ , so (20) reduces to

$$\frac{dP}{dt} = cP + d, \quad (22)$$

a first-order linear ODE in  $t$ , solved by the integrating factor (§3.6.1). When  $a$  and  $b$  are constants, the characteristics are straight lines and the quantity

$$\xi := bx - ay \quad (23)$$

is preserved along each curve—it is the *first integral* that labels each characteristic. The integration constant of (22) therefore varies between characteristics: it is an arbitrary function  $F(\xi)$ . Re-expressing the solution in the original variables  $(x, y)$  via (21) gives the general solution of (20). In the PGF setting,  $F$  is identified with a boundary generating function, and analyticity of  $P(x, y)$  for  $|x|, |y| \leq 1$  uniquely determines it.

## 4 Model description: Model- $X$

This section defines the general two-class system with both jockeying ( $\gamma_i$ ) and abandonments ( $\theta_i$ ). We refer to this full model as *Model- $X$* . The simpler variants studied are recovered as special cases by zeroing the appropriate parameters, as summarised in the table below (Table 3):

| Model   | Jockeying                        | Abandonment           | Parameter restriction                                                          |
|---------|----------------------------------|-----------------------|--------------------------------------------------------------------------------|
| $X$     | both ( $1 \leftrightarrow 2$ )   | both (1, 2)           | $\gamma_i \geq 0, \theta_i \geq 0$ (general)                                   |
| $A$     | none                             | none                  | $\gamma_1 = \gamma_2 = 0, \theta_1 = \theta_2 = 0$                             |
| $B$     | both ( $1 \leftrightarrow 2$ )   | none                  | $\gamma_i \geq 0, \theta_1 = \theta_2 = 0$                                     |
| $B_2$   | $1 \rightarrow 2$ only           | none                  | $\gamma_1 \geq 0, \gamma_2 = 0, \theta_1 = \theta_2 = 0$                       |
| $C_2$   | none                             | class 1 only          | $\theta_1 \geq 0, \theta_2 = 0, \gamma_1 = \gamma_2 = 0$                       |
| $B_2^H$ | $1 \rightarrow 2$ , head-of-line | none                  | $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}} > 0, \gamma_2 = \theta_1 = \theta_2 = 0$ |
| $C_2^H$ | none                             | class 1, head-of-line | $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}} > 0, \gamma_1 = \gamma_2 = \theta_2 = 0$ |

Table 3: Nested family of models. Model- $X$  is the parent; Models  $A$ ,  $B$ ,  $B_2$  and  $C_2$  are obtained by restricting the jockeying directions and the abandoning classes. The two head-of-line variants  $B_2^H$  and  $C_2^H$  are listed for completeness.<sup>†</sup>

<sup>†</sup>Unlike the others, the head-of-line variants are *not* obtained by zeroing parameters of Model- $X$ : their mechanism fires at the constant head-of-line rate  $\mathbf{1}_{\{n_1 \geq 1\}}$  rather than the length-proportional rate  $n_1$  of the parent (jockeying  $\gamma_1 n_1$ , abandonment  $\theta_1 n_1$ ), so they are standalone simplifications, as explained in §10.

Figure 1 provides a visual overview of Model- $X$  and each of the variants that we will consider:




Figure 1: Physical queue layout for Model- $X$  (left) and its four specialisations (right,  $2 \times 2$  grid). Queue 1 (top buffer) has non-preemptive priority over Queue 2. Solid arrows: Poisson arrivals ( $\lambda_i$ ) and exponential service ( $\mu$ ). Dashed arrows: jockeying ( $\gamma_i$ ), which transfers a waiting customer between queues and conserves the total  $n$ . Dotted arrows: abandonment ( $\theta_i$ ), a true departure that reduces  $n$ .

The model discussed in this section is an  $M/M/1$  queueing system with 2 priority classes, where customers are served on a First-Come, First-Served (FCFS) basis within their own class and those

coming from the priority class (class-1) have non-preemptive priority over those of the non-priority class (class-2); that is, when a class-2 customer is in service, that service will not be interrupted by a potential arrival of a class-1 customer. Beyond basic dynamics, Model- $X$  incorporates two distinct queue-leaving mechanisms. First, *jockeying*: a waiting customer can abandon its own queue and *join the other queue*, at a per-customer rate  $\gamma_1$  for direction  $1 \rightarrow 2$  and  $\gamma_2$  for direction  $2 \rightarrow 1$ , so the total number of customers is preserved. Second, *abandonment* (reneging): a waiting customer may completely leave the *system* at a rate per-customer  $\theta_i$ , so the total number of customers decreases by one. Concretely, each Class- $i$  customer that is waiting in line (i.e. not the one in service) holds an independent  $\text{Exp}(\theta_i)$  patience clock and abandons the system when it expires; consequently, the aggregate Class- $i$  abandonment rate in a state with  $n_i$  waiting customers is the linear term  $\theta_i n_i$  that appears in the balance equations. This is precisely the linear-reneging mechanism of the  $M/M/1 + M$  queue introduced in §3.3.

This model, being an  $M/M/1$  queueing model according to Kendall's notation, features arrivals driven by a Poisson process with arrival rates  $\lambda_i$  for Class- $i$ , exponentially distributed service times with rate  $\mu$ , and a service capacity of 1 customer at a time. In addition, a waiting customer of class-1 jockeys in the class-2 queue at the per-customer rate  $\gamma_1$  and a waiting customer of class-2 jockeys in the class-1 queue at the per-customer rate  $\gamma_2$  (transfers that conserve the total), while a waiting customer of Class- $i$  abandons the system at the per-customer rate  $\theta_i$  (a true departure that reduces the total).

In order to define the stochastic processes that drive this queueing system, it is necessary to introduce the following random variables:

- $\mathcal{B}$ : binary indicator for the server being idle or busy,  $\mathcal{B} \in \{0, 1\}$ ;
- $N_1, N_2$ : number of customers of class 1 and 2, respectively, in their respective queues;
- $N$ : total number of customers in the queues.

Note that the variables  $N_1$ ,  $N_2$  and  $N$  count the number of customers in the queues, as opposed to the customers in the system. This definition is convenient given that we have a common service rate for both priority classes, and hence the states of the system are equivalent regardless of the priority class the customer in service comes from. Since having customers in line directly implies that someone is in service, we will denote the idle state as  $(0)$  and the rest of the states by the number of people in each queue as the tuple  $(n_1, n_2)$ , where  $n_i$  denotes the number of customers in the queue for class- $i$ . This is the most conventional way to define the state space, namely  $S$ , in such a queueing system. However, we can also look at the total number of customers in the queues. We can define an alternative state space  $\tilde{S}$  using the idle state  $(0)$  and the pair  $(n_2, n)$  —we could also choose  $(n_1, n)$ —, which gives that we have  $(n - n_2)$  class-1 customers in the priority queue. Note that, since  $n$  is the total number of customers, we need the constraint  $n \geq n_2$ . Therefore, we define our state spaces as follows:

$$S := \{(0)\} \cup \{(n_1, n_2) \mid n_1, n_2 \geq 0\}, \quad (24)$$

$$\tilde{S} := \{(0)\} \cup \{(n_2, n) \mid n \geq n_2 \geq 0\}. \quad (25)$$

For these state spaces, we define the limiting —as opposed to time-dependent— probabilities of finding the system in that state in the long run as follows:

- $\pi_0$  for the idle state  $(0)$ , which belongs to both spaces  $S$  and  $\tilde{S}$ ;
- $\pi(n_1, n_2)$  for states  $(n_1, n_2) \in S$ , where  $n_1, n_2 \geq 0$ ;
- $\tilde{\pi}(n_2, n)$  for states  $(n_2, n) \in \tilde{S}$ , where  $n \geq n_2 \geq 0$ .

Regarding the parameters of the system, the arrival process for each class follows a Poisson process (i.e., the interarrival times are exponentially distributed). By the properties of Poisson processes, the total number of arrivals will also form a Poisson process. Furthermore, we have a common service rate for exponentially distributed service times. We can organise our notation into class-specific parameters (alongside our common service rate) and class-blind totals. For the class-specific metrics and service rate, we define:

- $\lambda_1, \lambda_2$ : arrival rates of classes 1 and 2, respectively;
- $\mu$ : service rate of a customer, regardless of class;
- $\rho_i = \frac{\lambda_i}{\mu}$ : system load of class  $i \in \{1, 2\}$ ;

- $\gamma_1, \gamma_2$ : per-customer jockeying rates from Class  $1 \rightarrow 2$  and  $2 \rightarrow 1$ , respectively; jockeying transfers a customer between queues, preserving the total number of customers;
- $\theta_1, \theta_2$ : per-customer abandonment (reneging) rates from classes 1 and 2, respectively; abandonment removes a customer from the system, reducing the total  $n$  by one.

For the aggregate system parameters, we define:

- $\lambda = \lambda_1 + \lambda_2$ : total arrival rate;
- $\rho = \frac{\lambda}{\mu}$ : total system load.

**Stability.** The stability conditions depend on the abandonments; jockeying does not affect this condition, since customers are merely relocated. Model- $X$ , having abandonments from both classes, is always stable, as the total number of customers behaves like some sort of  $M/M/1 + M$ —which is always stable, as stated in §3.3. The rate out is  $\mu + \theta_1 n_1 + \theta_2 n_2$ , which grows to infinity in every direction of the  $(n_1, n_2)$ -plane, so the system is always positive-recurrent. In contrast, Model- $A$ , having no abandonments whatsoever, requires  $\rho < 1$ , since the total number of customers behaves exactly like an  $M/M/1$  queueing system (§3.1). It is more complex for models where only one type of abandonment is possible, this will be discussed in the dedicated section for Model- $C_2$  (§9).

#### Computation of $\pi_0$ and $\pi(0, 0)$ .

The following relation between  $\pi_0$  and  $\pi(0, 0)$  holds not only for Model- $X$  but also for every other model we will see in this work. The only transition into state  $(0)$  is a service completion from  $(0, 0)$ , and the only transition out from  $(0, 0)$  is an arrival of a customer from any class to an idle system, leaving us the following balance equation:  $\lambda \pi_0 = \mu \pi(0, 0)$ . Therefore

$$\pi(0, 0) = \frac{\lambda}{\mu} \pi_0 = \rho \pi_0. \quad (26)$$

Note that this holds regardless jockeying and abandonments, as in neither of these states we have waiting customers. However, what does depend on the abandonment parameters—not on the jockeying ones—is the value of  $\pi_0$  itself. For models *without* abandonment, the total number of customers in the system behaves like that of an ordinary  $M/M/1$  single-class system, giving

$$\pi_0 = 1 - \rho = 1 - \frac{\lambda_1 + \lambda_2}{\mu}, \quad \pi(0, 0) = \rho \pi_0 = (1 - \rho)\rho. \quad (27)$$

However, for models *with* abandonments, the total number of customers is no longer conserved. For a busy state, the departure rate is  $\mu + \theta_1 n_1 + \theta_2 n_2$ , so the rate of state  $(n_1, n_2)$  depends on the split between  $n_1$  and  $n_2$ . Indeed, the states  $(n, 0)$  and  $(0, n)$  carry departure rates  $\mu + \theta_1 n$  and  $\mu + \theta_2 n$ , which differ whenever for asymmetric abandonment rates  $\theta_1 \neq \theta_2$ . To determine  $\pi_0$ , we will need our bivariate PGF, which will be defined in Section 4.2.

## 4.1 Balance equations

This subsection presents the state-transition diagrams of Model- $X$  on both state spaces  $S$  and  $\tilde{S}$ , and derives the corresponding sets of balance equations.

### 4.1.1 Balance equations for state space $S$

The state-transition diagram of Model- $X$  on  $S$  is given in Figure 2.




Figure 2: State transition diagram on state space  $S$  for the model with jockeying ( $\gamma_i n_i$ , dashed) and abandonments ( $\theta_i n_i$ , dotted).

$$\begin{aligned}
& [\lambda_1 + \lambda_2 + \mu + (\gamma_1 + \theta_1)n_1 + (\gamma_2 + \theta_2)n_2] \pi(n_1, n_2) \\
&= \mu \pi(n_1 + 1, n_2) \\
&+ \lambda_1 \pi(n_1 - 1, n_2) \\
&+ \lambda_2 \pi(n_1, n_2 - 1) \\
&+ \gamma_1(n_1 + 1) \pi(n_1 + 1, n_2 - 1) \\
&+ \gamma_2(n_2 + 1) \pi(n_1 - 1, n_2 + 1) \\
&+ \theta_1(n_1 + 1) \pi(n_1 + 1, n_2) \\
&+ \theta_2(n_2 + 1) \pi(n_1, n_2 + 1)
\end{aligned}
\quad \text{for } n_1 \geq 1, n_2 \geq 1 \quad (28)$$

$$\begin{aligned}
& [\lambda_1 + \lambda_2 + \mu + (\gamma_2 + \theta_2)n_2] \pi(0, n_2) \\
&= \lambda_2 \pi(0, n_2 - 1) \\
&+ \mu \pi(1, n_2) \\
&+ \mu \pi(0, n_2 + 1) \\
&+ \gamma_1 \pi(1, n_2 - 1) \\
&+ \theta_1 \pi(1, n_2) \\
&+ \theta_2(n_2 + 1) \pi(0, n_2 + 1)
\end{aligned}
\quad \text{for } n_1 = 0, n_2 \geq 1 \quad (29)$$

$$\begin{aligned}
& [\lambda_1 + \lambda_2 + \mu + (\gamma_1 + \theta_1)n_1] \pi(n_1, 0) \\
&= \lambda_1 \pi(n_1 - 1, 0) \\
&+ \mu \pi(n_1 + 1, 0) \\
&+ \gamma_2 \pi(n_1 - 1, 1) \\
&+ \theta_1(n_1 + 1) \pi(n_1 + 1, 0) \\
&+ \theta_2 \pi(n_1, 1)
\end{aligned}
\quad \text{for } n_1 \geq 1, n_2 = 0 \quad (30)$$

$$[\lambda_1 + \lambda_2] \pi(0, 0) = (\mu + \theta_1) \pi(1, 0) + (\mu + \theta_2) \pi(0, 1) \quad (31)$$

$$(\lambda_1 + \lambda_2) \pi_0 = \mu \pi(0, 0) \quad (32)$$



### 4.1.2 Balance equations for state space $\tilde{S}$

Recall the state space  $\tilde{S}$  defined in (25).

For the sake of readability, we will denote the limiting probabilities associated to this state space with  $\tilde{\pi}(n_2, n)$ , for state  $(n_2, n) \in \tilde{S}$ . Figure 3 shows the corresponding state transition diagram.




Figure 3: State transition diagram on state space  $\tilde{S}$ .

The balance equations, after combining the equations where  $\tilde{\pi}_0$  shows up, are:

$$\begin{aligned}
 & [\lambda_1 + \lambda_2 + \mu + (\gamma_1 + \theta_1)(n - n_2) + (\gamma_2 + \theta_2)n_2] \tilde{\pi}(n_2, n) \\
 &= \lambda_1 \tilde{\pi}(n_2, n - 1) \\
 &+ \lambda_2 \tilde{\pi}(n_2 - 1, n - 1) \\
 &+ \mu \tilde{\pi}(n_2, n + 1) \\
 &+ \gamma_1(n - n_2 + 1) \tilde{\pi}(n_2 - 1, n) \\
 &+ \gamma_2(n_2 + 1) \tilde{\pi}(n_2 + 1, n) \\
 &+ \theta_1(n + 1 - n_2) \tilde{\pi}(n_2, n + 1) \\
 &+ \theta_2(n_2 + 1) \tilde{\pi}(n_2 + 1, n + 1)
 \end{aligned}
 \quad \text{for } 1 \leq n_2 \leq n - 1, n \geq 2 \quad (33)$$

$$\begin{aligned}
 & [\lambda_1 + \lambda_2 + \mu + (\gamma_2 + \theta_2)n] \tilde{\pi}(n, n) \\
 &= \lambda_2 \tilde{\pi}(n - 1, n - 1) \\
 &+ \mu \tilde{\pi}(n + 1, n + 1) \\
 &+ \mu \tilde{\pi}(n, n + 1) \\
 &+ \gamma_1 \tilde{\pi}(n - 1, n) \\
 &+ \theta_1 \tilde{\pi}(n, n + 1) \\
 &+ \theta_2(n + 1) \tilde{\pi}(n + 1, n + 1)
 \end{aligned}
 \quad \text{for } n_2 = n, n \geq 1 \quad (34)$$

$$\begin{aligned}
 & [\lambda_1 + \lambda_2 + \mu + (\gamma_1 + \theta_1)n] \tilde{\pi}(0, n) \\
 &= \lambda_1 \tilde{\pi}(0, n - 1) \\
 &+ \mu \tilde{\pi}(0, n + 1) \\
 &+ \gamma_2 \tilde{\pi}(1, n) \\
 &+ \theta_1(n + 1) \tilde{\pi}(0, n + 1) \\
 &+ \theta_2 \tilde{\pi}(1, n + 1)
 \end{aligned}
 \quad \text{for } n_2 = 0, n \geq 1 \quad (35)$$

$$\begin{aligned}
& [\lambda_1 + \lambda_2] \tilde{\pi}(0, 0) \\
& = (\mu + \theta_1) \tilde{\pi}(0, 1) + (\mu + \theta_2) \tilde{\pi}(1, 1)
\end{aligned} \tag{36}$$

$$(\lambda_1 + \lambda_2) \tilde{\pi}_0 = \mu \tilde{\pi}(0, 0) \tag{37}$$

## 4.2 Derivation of equations for (Partial) Probability Generating functions

In order to study the limiting behaviour of this two-class system, we define the bivariate Probability Generating Function (PGF) of the joint number of customers in the queue for each class,  $N_1$  and  $N_2$ :

$$P(x, y) = \mathbb{E}[x^{N_1} y^{N_2}] = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2}. \tag{38}$$

The derivation that will be carried out here includes the boundary terms, where at least one of the two queues is empty:

$$P_x(x) = P(x, 0) = \sum_{n_1=0}^{\infty} \pi(n_1, 0) x^{n_1} \tag{39}$$

$$P_y(y) = P(0, y) = \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2}. \tag{40}$$

Note that both of these terms include the state where the server is busy but both queues are empty, which is represented by  $\pi(0, 0)$ , while the state where the entire system is empty (the idle state) is treated separately.

Another term that arises during the application of the balance equations characterises the boundary dynamics when the class-1 queue contains exactly one waiting customer:

$$P_1(y) = \sum_{n_2=1}^{\infty} \pi(1, n_2) y^{n_2}. \tag{41}$$

This term serves as an intermediate step; as the derivation proceeds, it will be possible to express it in terms of the primary boundary functions  $P_x(x)$  and  $P_y(y)$ .

In the same fashion, we introduce the concept of the **Partial Probability Generating Function (PPGF)**. This technique, notably utilised by Cohen [Coh56], consists of transforming only a subset of the random variables while maintaining the discrete indices of the remainder. In the context of a priority queue, this method is particularly advantageous: it allows us to exploit the well-known properties of the priority class (often a standard  $M/M/1$  system) and focus the analytical effort on the non-trivial PGF of the non-priority class.

While Cohen originally developed this framework for multi-server systems, we apply it here to the single-server case. To avoid the potential confusion stemming from mid-20th-century notation, we provide a modern formulation consistent with the variables defined above.

We define the PPGF with respect to the second variable  $N_2$ , conditioned on the first queue being in state  $n_1$ , as:

$$\tilde{P}(n_1, y) = \mathbb{E}[y^{N_2} |_{\{N_1=n_1\}}] = \sum_{n_2=0}^{\infty} \pi(n_1, n_2) y^{n_2}. \tag{42}$$

Similarly, the transform with respect to the first variable  $N_1$ , while maintaining the discrete state  $n_2$  for the second queue, is defined as:

$$\tilde{P}(x, n_2) = \mathbb{E}[x^{N_1} |_{\{N_2=n_2\}}] = \sum_{n_1=0}^{\infty} \pi(n_1, n_2) x^{n_1}. \tag{43}$$

These partial transforms are linked to the joint PGF by the following relations:

$$P(x, y) = \sum_{n_1=0}^{\infty} \tilde{P}(n_1, y) x^{n_1} = \sum_{n_2=0}^{\infty} \tilde{P}(x, n_2) y^{n_2}. \tag{44}$$

#### 4.2.1 Equation for PGF in state space $S$

**Lemma 1** (Fundamental equation for the PGF on  $S$ ). *Consider the two-class non-preemptive priority queue on the state space  $S = \{(n_1, n_2) : n_1, n_2 \geq 0\}$ , with Poisson arrival rates  $\lambda_1, \lambda_2$ , exponential service rate  $\mu$ , jockeying rates  $\gamma_1, \gamma_2$  and abandonment rates  $\theta_1, \theta_2$ , where  $n_1$  and  $n_2$  count the class-1 and class-2 customers waiting in the queues. Assume the associated CTMC is positive recurrent, so that the stationary distribution  $\{\pi(n_1, n_2)\}$  exists and the joint PGF*

$$P(x, y) = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2}$$

*is analytic on the closed unit polydisc  $|x| \leq 1, |y| \leq 1$ . Then  $P(x, y)$ , together with the boundary function  $P_y(y) = P(0, y) = \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2}$ , satisfies the first-order linear partial differential equation*

$$\begin{aligned} & [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) \\ & + xy [\gamma_1(x - y) + \theta_1(x - 1)] \frac{\partial}{\partial x} P(x, y) \\ & + xy [\gamma_2(y - x) + \theta_2(y - 1)] \frac{\partial}{\partial y} P(x, y) \\ & = \mu(x - y)P_y(y) - \mu x(1 - y)\pi(0, 0). \end{aligned} \tag{45}$$

**Reading the fundamental equation.** Equation (45) repays a moment's mechanical reading, since its structure dictates the entire analysis that follows. The bracketed algebraic coefficient multiplying  $P(x, y)$  is exactly the kernel of the baseline Model-A: it is the flux balance between the arrival contributions  $\lambda_1 x^2 y$  and  $\lambda_2 x y^2$ , the service contribution  $\mu y$ , and the total outflow  $(\lambda_1 + \lambda_2 + \mu)xy$ . The two convection coefficients carry the added mechanisms. In the  $x$ -convection coefficient  $xy [\gamma_1(x - y) + \theta_1(x - 1)]$ , the jockeying factor  $(x - y)$  vanishes on the diagonal  $x = y$ , the analytic signature that jockeying merely relocates a customer and so conserves the total count, while the abandonment factor  $(x - 1)$  vanishes at  $x = 1$ , the signature that abandonment respects normalisation ( $P(1, 1) = 1$ ) yet removes a customer and thereby destroys conservation. The  $y$ -convection coefficient  $xy [\gamma_2(y - x) + \theta_2(y - 1)]$  states the symmetric facts for the class-2 mechanisms, with  $(y - x)$  vanishing on the diagonal and  $(y - 1)$  at  $y = 1$ . The right-hand side carries the two boundary unknowns that any solution must pin down: the marginal  $P_y(y) = P(0, y)$ , the generating function of the states with an empty priority queue, and the scalar  $\pi(0, 0)$ , the probability that the server is busy with both queues empty. The position of these vanishing factors fixes, model by model, where the operative singularity sits and hence which technique applies: with both convection terms absent (Model-A, §5) the equation is algebraic and  $P_y(y)$  is recovered from the kernel root  $x^*(y)$  inside the unit disc; with one-way jockeying alone (Model- $B_2$ , §8) the factor  $(x - y)$  places the singularity at the interior point  $x = y$  with a negative exponent, and  $P_y(y)$  is pinned by analyticity there; with class-1 abandonment alone (Model- $C_2$ , §9) the factor  $(x - 1)$  places it on the boundary  $x = 1$  with a positive exponent, and  $P_y(y)$  is pinned by mere boundedness; and restricting a mechanism to the head of the queue removes the derivative outright, collapsing the equation to the algebraic Model-A form (Models  $B_2^H$  and  $C_2^H$ , §10). These four regimes are gathered side by side in Table 5 of §12.

*Proof.* We multiply each balance equation by  $x^{n_1} y^{n_2}$  and sum over its domain.

From the interior equation (28):

$$\begin{aligned}
& (\lambda_1 + \lambda_2 + \mu) \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2} \\
& + \gamma_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_1 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
& + \gamma_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
& + \theta_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_1 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
& + \theta_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
& = \mu \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1 + 1, n_2) x^{n_1} y^{n_2} \\
& + \lambda_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1 - 1, n_2) x^{n_1} y^{n_2} \\
& + \lambda_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1, n_2 - 1) x^{n_1} y^{n_2} \\
& + \gamma_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_1 + 1) \pi(n_1 + 1, n_2 - 1) x^{n_1} y^{n_2} \\
& + \gamma_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_2 + 1) \pi(n_1 - 1, n_2 + 1) x^{n_1} y^{n_2} \\
& + \theta_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_1 + 1) \pi(n_1 + 1, n_2) x^{n_1} y^{n_2} \\
& + \theta_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_2 + 1) \pi(n_1, n_2 + 1) x^{n_1} y^{n_2} \\
& LHS_1 + LHS_2 + LHS_3 + LHS_4 + LHS_5 \\
& = RHS_1 + RHS_2 + RHS_3 + RHS_4 + RHS_5 + RHS_6 + RHS_7
\end{aligned} \tag{46}$$

$$\begin{aligned}
& LHS_1 + LHS_2 + LHS_3 + LHS_4 + LHS_5 \\
& = RHS_1 + RHS_2 + RHS_3 + RHS_4 + RHS_5 + RHS_6 + RHS_7
\end{aligned} \tag{47}$$

Evaluating each summand:

$$\begin{aligned}
& LHS_1 \\
& = (\lambda_1 + \lambda_2 + \mu) \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2} \\
& = (\lambda_1 + \lambda_2 + \mu) \sum_{n_1=1}^{\infty} \left[ \sum_{n_2=0}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2} - \pi(n_1, 0) x^{n_1} y^0 \right] \\
& = (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2} - \sum_{n_1=1}^{\infty} \pi(n_1, 0) x^{n_1} \right] \\
& = (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \pi(n_1, n_2) x^{n_1} y^{n_2} - \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2} \right] \\
& \quad - (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_1=0}^{\infty} \pi(n_1, 0) x^{n_1} - \pi(0, 0) \right] \\
& = (\lambda_1 + \lambda_2 + \mu) [P(x, y) - P_y(y) - P_x(x) + \pi(0, 0)]
\end{aligned} \tag{48}$$

$LHS_2$

$$\begin{aligned}
&= \gamma_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_1 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
&= \gamma_1 x \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_1 \pi(n_1, n_2) x^{n_1-1} y^{n_2} \\
&= \gamma_1 x \left[ \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} n_1 \pi(n_1, n_2) x^{n_1-1} y^{n_2} - \sum_{n_1=1}^{\infty} n_1 \pi(n_1, 0) x^{n_1-1} \right] \\
&= \gamma_1 x \left[ \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} n_1 \pi(n_1, n_2) x^{n_1-1} y^{n_2} - \sum_{n_1=0}^{\infty} n_1 \pi(n_1, 0) x^{n_1-1} \right] \\
&= \gamma_1 x \left[ \frac{\partial}{\partial x} P(x, y) - \frac{d}{dx} P_x(x) \right]
\end{aligned} \tag{49}$$

$LHS_3$

$$\begin{aligned}
&= \gamma_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
&= \gamma_2 \sum_{n_1=1}^{\infty} \left[ \sum_{n_2=0}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2} - 0 \cdot \pi(n_1, 0) x^{n_1} \right] \\
&= \gamma_2 y \left[ \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2-1} - \sum_{n_2=0}^{\infty} n_2 \pi(0, n_2) y^{n_2-1} \right] \\
&= \gamma_2 y \left[ \frac{\partial}{\partial y} P(x, y) - \frac{d}{dy} P_y(y) \right]
\end{aligned} \tag{50}$$

$LHS_4$

$$\begin{aligned}
&= \theta_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_1 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
&= \theta_1 x \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_1 \pi(n_1, n_2) x^{n_1-1} y^{n_2} \\
&= \theta_1 x \left[ \sum_{n_1=1}^{\infty} \sum_{n_2=0}^{\infty} n_1 \pi(n_1, n_2) x^{n_1-1} y^{n_2} - \sum_{n_1=1}^{\infty} n_1 \pi(n_1, 0) x^{n_1-1} \right] \\
&= \theta_1 x \left[ \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} n_1 \pi(n_1, n_2) x^{n_1-1} y^{n_2} - \sum_{n_1=0}^{\infty} n_1 \pi(n_1, 0) x^{n_1-1} \right] \\
&= \theta_1 x \left[ \frac{\partial}{\partial x} P(x, y) - \frac{d}{dx} P_x(x) \right]
\end{aligned} \tag{51}$$

$LHS_5$

$$\begin{aligned}
&= \theta_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2} \\
&= \theta_2 y \left[ \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} n_2 \pi(n_1, n_2) x^{n_1} y^{n_2-1} - \sum_{n_2=0}^{\infty} n_2 \pi(0, n_2) y^{n_2-1} \right] \\
&= \theta_2 y \left[ \frac{\partial}{\partial y} P(x, y) - \frac{d}{dy} P_y(y) \right]
\end{aligned} \tag{52}$$

$RHS_1$

$$\begin{aligned}
&= \mu \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1+1, n_2) x^{n_1} y^{n_2} \\
&= \mu x^{-1} \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1+1, n_2) x^{n_1+1} y^{n_2} \\
&= \mu x^{-1} \sum_{m=2}^{\infty} \sum_{n_2=1}^{\infty} \pi(m, n_2) x^m y^{n_2} \\
&= \mu x^{-1} \sum_{m=2}^{\infty} \left[ \sum_{n_2=0}^{\infty} \pi(m, n_2) x^m y^{n_2} - \pi(m, 0) x^m \right] \\
&= \mu x^{-1} \left[ \sum_{m=2}^{\infty} \sum_{n_2=0}^{\infty} \pi(m, n_2) x^m y^{n_2} - \sum_{m=2}^{\infty} \pi(m, 0) x^m \right] \\
&= \mu x^{-1} \left[ \sum_{m=0}^{\infty} \sum_{n_2=0}^{\infty} \pi(m, n_2) x^m y^{n_2} - x \sum_{n_2=0}^{\infty} \pi(1, n_2) y^{n_2} - \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2} \right] \\
&\quad - \mu x^{-1} \left[ \sum_{m=0}^{\infty} \pi(m, 0) x^m - x \pi(1, 0) - \pi(0, 0) \right] \\
&= \mu x^{-1} [P(x, y) - x P_y(y|n_1=1) - P_y(y) - P_x(x) + x \pi(1, 0) + \pi(0, 0)]
\end{aligned} \tag{53}$$

$RHS_2$

$$\begin{aligned}
&= \lambda_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1-1, n_2) x^{n_1} y^{n_2} \\
&= \lambda_1 x \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1-1, n_2) x^{n_1-1} y^{n_2} \\
&= \lambda_1 x \sum_{m=0}^{\infty} \sum_{n_2=1}^{\infty} \pi(m, n_2) x^m y^{n_2} \\
&= \lambda_1 x \sum_{m=0}^{\infty} \left[ \sum_{n_2=0}^{\infty} \pi(m, n_2) x^m y^{n_2} - \pi(m, 0) x^m \right] \\
&= \lambda_1 x \left[ \sum_{m=0}^{\infty} \sum_{n_2=0}^{\infty} \pi(m, n_2) x^m y^{n_2} - \sum_{m=0}^{\infty} \pi(m, 0) x^m \right] \\
&= \lambda_1 x [P(x, y) - P_x(x)]
\end{aligned} \tag{54}$$

$RHS_3$

$$\begin{aligned}
&= \lambda_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1, n_2-1) x^{n_1} y^{n_2} \\
&= \lambda_2 y \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} \pi(n_1, n_2-1) x^{n_1} y^{n_2-1} \\
&= \lambda_2 y \sum_{n_1=1}^{\infty} \sum_{m=0}^{\infty} \pi(n_1, m) x^{n_1} y^m \\
&= \lambda_2 y \left[ \sum_{n_1=0}^{\infty} \sum_{m=0}^{\infty} \pi(n_1, m) x^{n_1} y^m - \sum_{m=0}^{\infty} \pi(0, m) y^m \right] \\
&= \lambda_2 y [P(x, y) - P_y(y)]
\end{aligned} \tag{55}$$

$RHS_4$

$$\begin{aligned}
&= \gamma_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_1+1)\pi(n_1+1, n_2-1)x^{n_1}y^{n_2} \\
&= \gamma_1 y \sum_{n_1=1}^{\infty} \sum_{m_2=0}^{\infty} (n_1+1)\pi(n_1+1, m_2)x^{n_1}y^{m_2} \\
&= \gamma_1 y \sum_{m_1=2}^{\infty} \sum_{m_2=0}^{\infty} m_1\pi(m_1, m_2)x^{m_1-1}y^{m_2} \\
&= \gamma_1 y \left[ \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} m_1\pi(m_1, m_2)x^{m_1-1}y^{m_2} - \sum_{m_2=0}^{\infty} \pi(1, m_2)y^{m_2} \right] \\
&= \gamma_1 y \left[ \frac{\partial}{\partial x} P(x, y) - P_y(y|n_1=1) \right]
\end{aligned} \tag{56}$$

$RHS_5$

$$\begin{aligned}
&= \gamma_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_2+1)\pi(n_1-1, n_2+1)x^{n_1}y^{n_2} \\
&= \gamma_2 \sum_{n_1=1}^{\infty} \sum_{m_2=2}^{\infty} m_2\pi(n_1-1, m_2)x^{n_1}y^{m_2-1} \\
&= \gamma_2 x \left[ \sum_{n_1=1}^{\infty} \sum_{m_2=0}^{\infty} m_2\pi(n_1-1, m_2)x^{n_1-1}y^{m_2-1} - \sum_{n_1=1}^{\infty} \pi(n_1-1, 1)x^{n_1-1} \right] \\
&= \gamma_2 x \left[ \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} m_2\pi(m_1, m_2)x^{m_1}y^{m_2-1} - \sum_{m_1=0}^{\infty} \pi(m_1, 1)x^{m_1} \right] \\
&= \gamma_2 x \left[ \frac{\partial}{\partial y} P(x, y) - P_x(x|n_2=1) \right]
\end{aligned} \tag{57}$$

$RHS_6$

$$\begin{aligned}
&= \theta_1 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_1+1)\pi(n_1+1, n_2)x^{n_1}y^{n_2} \\
&= \theta_1 \sum_{m_1=2}^{\infty} \sum_{n_2=1}^{\infty} m_1\pi(m_1, n_2)x^{m_1-1}y^{n_2} \\
&= \theta_1 \left[ \sum_{m_1=0}^{\infty} \sum_{n_2=1}^{\infty} m_1\pi(m_1, n_2)x^{m_1-1}y^{n_2} - \sum_{n_2=1}^{\infty} \pi(1, n_2)y^{n_2} \right] \\
&= \theta_1 \left[ \sum_{m_1=0}^{\infty} \sum_{n_2=0}^{\infty} m_1\pi(m_1, n_2)x^{m_1-1}y^{n_2} - \sum_{m_1=0}^{\infty} m_1\pi(m_1, 0)x^{m_1-1} \right] \\
&\quad - \theta_1 \left[ \sum_{n_2=0}^{\infty} \pi(1, n_2)y^{n_2} - \pi(1, 0) \right] \\
&= \theta_1 \left[ \frac{\partial}{\partial x} P(x, y) - \frac{d}{dx} P_x(x) - P_y(y|n_1=1) + \pi(1, 0) \right]
\end{aligned} \tag{58}$$



$RHS_7$

$$\begin{aligned}
&= \theta_2 \sum_{n_1=1}^{\infty} \sum_{n_2=1}^{\infty} (n_2+1) \pi(n_1, n_2+1) x^{n_1} y^{n_2} \\
&= \theta_2 \sum_{n_1=1}^{\infty} \sum_{m_2=2}^{\infty} m_2 \pi(n_1, m_2) x^{n_1} y^{m_2-1} \\
&= \theta_2 \left[ \sum_{n_1=1}^{\infty} \sum_{m_2=0}^{\infty} m_2 \pi(n_1, m_2) x^{n_1} y^{m_2-1} - \sum_{n_1=1}^{\infty} \pi(n_1, 1) x^{n_1} \right] \\
&= \theta_2 \left[ \sum_{n_1=0}^{\infty} \sum_{m_2=0}^{\infty} m_2 \pi(n_1, m_2) x^{n_1} y^{m_2-1} - \sum_{m_2=0}^{\infty} m_2 \pi(0, m_2) y^{m_2-1} \right] \\
&\quad - \theta_2 \left[ \sum_{n_1=0}^{\infty} \pi(n_1, 1) x^{n_1} - \pi(0, 1) \right] \\
&= \theta_2 \left[ \frac{\partial}{\partial y} P(x, y) - \frac{d}{dy} P_y(y) - P_x(x|n_2=1) + \pi(0, 1) \right]
\end{aligned} \tag{59}$$

Collecting and multiplying through by  $xy$  to clear the  $x^{-1}$  factor:

$$\begin{aligned}
&(\lambda_1 + \lambda_2 + \mu) [P(x, y) - P_y(y) - P_x(x) + \pi(0, 0)] \\
&+ \gamma_1 x \left[ \frac{\partial}{\partial x} P(x, y) - \frac{d}{dx} P_x(x) \right] \\
&+ \gamma_2 y \left[ \frac{\partial}{\partial y} P(x, y) - \frac{d}{dy} P_y(y) \right] \\
&+ \theta_1 x \left[ \frac{\partial}{\partial x} P(x, y) - \frac{d}{dx} P_x(x) \right] \\
&+ \theta_2 y \left[ \frac{\partial}{\partial y} P(x, y) - \frac{d}{dy} P_y(y) \right] \\
&= \mu x^{-1} [P(x, y) - x P_y(y|n_1=1) - P_y(y) - P_x(x) + x \pi(1, 0) + \pi(0, 0)] \\
&\quad + \lambda_1 x [P(x, y) - P_x(x)] \\
&\quad + \lambda_2 y [P(x, y) - P_y(y)] \\
&\quad + \gamma_1 y \left[ \frac{\partial}{\partial x} P(x, y) - P_y(y|n_1=1) \right] \\
&\quad + \gamma_2 x \left[ \frac{\partial}{\partial y} P(x, y) - P_x(x|n_2=1) \right] \\
&\quad + \theta_1 \left[ \frac{\partial}{\partial x} P(x, y) - \frac{d}{dx} P_x(x) - P_y(y|n_1=1) + \pi(1, 0) \right] \\
&\quad + \theta_2 \left[ \frac{\partial}{\partial y} P(x, y) - \frac{d}{dy} P_y(y) - P_x(x|n_2=1) + \pi(0, 1) \right]
\end{aligned}$$

$$\begin{aligned}
& [(\lambda_1 + \lambda_2 + \mu) - \mu x^{-1} - \lambda_1 x - \lambda_2 y] P(x, y) \\
& + [\gamma_1(x - y) + \theta_1(x - 1)] \frac{\partial}{\partial x} P(x, y) \\
& + [\gamma_2(y - x) + \theta_2(y - 1)] \frac{\partial}{\partial y} P(x, y) \\
& = [(\lambda_1 + \lambda_2 + \mu) - \mu x^{-1} - \lambda_1 x] P_x(x) \\
& + [(\lambda_1 + \lambda_2 + \mu) - \mu x^{-1} - \lambda_2 y] P_y(y) \\
& + [-(\lambda_1 + \lambda_2 + \mu) + \mu x^{-1}] \pi(0, 0) \\
& + [\gamma_1 x + \theta_1(x - 1)] \frac{d}{dx} P_x(x) \\
& + [\gamma_2 y + \theta_2(y - 1)] \frac{d}{dy} P_y(y) \\
& - [\mu + \gamma_1 y + \theta_1] P_y(y|n_1 = 1) \\
& - [\gamma_2 x + \theta_2] P_x(x|n_2 = 1) \\
& + \mu \pi(1, 0) \\
& + \theta_1 \pi(1, 0) \\
& + \theta_2 \pi(0, 1) \\
& [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 xy^2] P(x, y) \\
& + xy [\gamma_1(x - y) + \theta_1(x - 1)] \frac{\partial}{\partial x} P(x, y) \\
& + xy [\gamma_2(y - x) + \theta_2(y - 1)] \frac{\partial}{\partial y} P(x, y) \\
& = [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y] P_x(x) \\
& + [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_2 xy^2] P_y(y) \\
& - [(\lambda_1 + \lambda_2 + \mu)xy - \mu y] \pi(0, 0) \\
& + xy [\gamma_1 x + \theta_1(x - 1)] \frac{d}{dx} P_x(x) \\
& + xy [\gamma_2 y + \theta_2(y - 1)] \frac{d}{dy} P_y(y) \\
& - xy [\mu + \gamma_1 y + \theta_1] P_y(y|n_1 = 1) \\
& - xy [\gamma_2 x + \theta_2] P_x(x|n_2 = 1) \\
& + (\mu + \theta_1)xy \pi(1, 0) \\
& + \theta_2 xy \pi(0, 1)
\end{aligned} \tag{60}$$

The  $x$ -boundary equation (30), treated analogously, gives:

$$\begin{aligned}
& \sum_{n_1=1}^{\infty} [\lambda_1 + \lambda_2 + \mu + (\gamma_1 + \theta_1)n_1] \pi(n_1, 0)x^{n_1} \\
& = \lambda_1 \sum_{n_1=1}^{\infty} \pi(n_1 - 1, 0)x^{n_1} + \mu \sum_{n_1=1}^{\infty} \pi(n_1 + 1, 0)x^{n_1} \\
& + \gamma_2 \sum_{n_1=1}^{\infty} \pi(n_1 - 1, 1)x^{n_1} + \theta_1 \sum_{n_1=1}^{\infty} (n_1 + 1)\pi(n_1 + 1, 0)x^{n_1} + \theta_2 \sum_{n_1=1}^{\infty} \pi(n_1, 1)x^{n_1} \\
& (\lambda_1 + \lambda_2 + \mu) \sum_{n_1=1}^{\infty} \pi(n_1, 0)x^{n_1} + (\gamma_1 + \theta_1)x \sum_{n_1=1}^{\infty} n_1 \pi(n_1, 0)x^{n_1-1} \\
& = \lambda_1 x \sum_{m=0}^{\infty} \pi(m, 0)x^m + \mu x^{-1} \sum_{m=2}^{\infty} \pi(m, 0)x^m \\
& + \gamma_2 x \sum_{m=0}^{\infty} \pi(m, 1)x^m + \theta_1 \sum_{m=2}^{\infty} m \pi(m, 0)x^{m-1} + \theta_2 \sum_{m=1}^{\infty} \pi(m, 1)x^m
\end{aligned}$$

$$\begin{aligned}
& (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_1=0}^{\infty} \pi(n_1, 0) x^{n_1} - \pi(0, 0) \right] + (\gamma_1 + \theta_1) x \sum_{n_1=0}^{\infty} n_1 \pi(n_1, 0) x^{n_1-1} \\
&= \lambda_1 x \sum_{m=0}^{\infty} \pi(m, 0) x^m \\
&\quad + \mu x^{-1} \left[ \sum_{m=0}^{\infty} \pi(m, 0) x^m - x \pi(1, 0) - \pi(0, 0) \right] \\
&\quad + \gamma_2 x \sum_{m=0}^{\infty} \pi(m, 1) x^m \\
&\quad + \theta_1 \left[ \sum_{m=0}^{\infty} m \pi(m, 0) x^{m-1} - \pi(1, 0) \right] \\
&\quad + \theta_2 \left[ \sum_{m=0}^{\infty} \pi(m, 1) x^m - \pi(0, 1) \right] \\
& (\lambda_1 + \lambda_2 + \mu) [P_x(x) - \pi(0, 0)] + (\gamma_1 + \theta_1) x \frac{d}{dx} P_x(x) \\
&= \lambda_1 x P_x(x) + \mu x^{-1} [P_x(x) - x \pi(1, 0) - \pi(0, 0)] \\
&\quad + \gamma_2 x P_x(x|n_2 = 1) + \theta_1 \left[ \frac{d}{dx} P_x(x) - \pi(1, 0) \right] + \theta_2 [P_x(x|n_2 = 1) - \pi(0, 1)] \\
& [(\lambda_1 + \lambda_2 + \mu) - \mu x^{-1} - \lambda_1 x] P_x(x) + [\gamma_1 x + \theta_1(x - 1)] \frac{d}{dx} P_x(x) \\
&= [(\lambda_1 + \lambda_2 + \mu) - \mu x^{-1}] \pi(0, 0) - (\mu + \theta_1) \pi(1, 0) - \theta_2 \pi(0, 1) \\
&\quad + [\gamma_2 x + \theta_2] P_x(x|n_2 = 1) \\
& [(\lambda_1 + \lambda_2 + \mu) xy - \mu y - \lambda_1 x^2 y] P_x(x) + xy [\gamma_1 x + \theta_1(x - 1)] \frac{d}{dx} P_x(x) \\
&= [(\lambda_1 + \lambda_2 + \mu) xy - \mu y] \pi(0, 0) \\
&\quad - (\mu + \theta_1) xy \pi(1, 0) \\
&\quad - \theta_2 xy \pi(0, 1) \\
&\quad + xy [\gamma_2 x + \theta_2] P_x(x|n_2 = 1)
\end{aligned} \tag{61}$$

The  $y$ -boundary equation (29) gives:

$$\begin{aligned}
& \sum_{n_2=1}^{\infty} [\lambda_1 + \lambda_2 + \mu + (\gamma_2 + \theta_2) n_2] \pi(0, n_2) y^{n_2} \\
&= \lambda_2 \sum_{n_2=1}^{\infty} \pi(0, n_2 - 1) y^{n_2} \\
&\quad + \mu \sum_{n_2=1}^{\infty} \pi(1, n_2) y^{n_2} \\
&\quad + \mu \sum_{n_2=1}^{\infty} \pi(0, n_2 + 1) y^{n_2} \\
&\quad + \gamma_1 \sum_{n_2=1}^{\infty} \pi(1, n_2 - 1) y^{n_2} \\
&\quad + \theta_1 \sum_{n_2=1}^{\infty} \pi(1, n_2) y^{n_2} \\
&\quad + \theta_2 \sum_{n_2=1}^{\infty} (n_2 + 1) \pi(0, n_2 + 1) y^{n_2}
\end{aligned}$$

$$\begin{aligned}
& (\lambda_1 + \lambda_2 + \mu) \sum_{n_2=1}^{\infty} \pi(0, n_2) y^{n_2} \\
& + (\gamma_2 + \theta_2) y \sum_{n_2=1}^{\infty} n_2 \pi(0, n_2) y^{n_2-1} \\
& = \lambda_2 y \sum_{m=0}^{\infty} \pi(0, m) y^m \\
& + \mu \sum_{n_2=1}^{\infty} \pi(1, n_2) y^{n_2} \\
& + \mu y^{-1} \sum_{m=2}^{\infty} \pi(0, m) y^m \\
& + \gamma_1 y \sum_{m=0}^{\infty} \pi(1, m) y^m \\
& + \theta_1 \sum_{n_2=1}^{\infty} \pi(1, n_2) y^{n_2} \\
& + \theta_2 \sum_{m=2}^{\infty} m \pi(0, m) y^{m-1} \\
& (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2} - \pi(0, 0) \right] \\
& + (\gamma_2 + \theta_2) y \sum_{n_2=0}^{\infty} n_2 \pi(0, n_2) y^{n_2-1} \\
& = \lambda_2 y \sum_{m=0}^{\infty} \pi(0, m) y^m \\
& + \mu \left[ \sum_{n_2=0}^{\infty} \pi(1, n_2) y^{n_2} - \pi(1, 0) \right] \\
& + \mu y^{-1} \left[ \sum_{m=0}^{\infty} \pi(0, m) y^m - y \pi(0, 1) - \pi(0, 0) \right] \\
& + \gamma_1 y \sum_{m=0}^{\infty} \pi(1, m) y^m \\
& + \theta_1 \left[ \sum_{n_2=0}^{\infty} \pi(1, n_2) y^{n_2} - \pi(1, 0) \right] \\
& + \theta_2 \left[ \sum_{m=0}^{\infty} m \pi(0, m) y^{m-1} - \pi(0, 1) \right] \\
& (\lambda_1 + \lambda_2 + \mu) [P_y(y) - \pi(0, 0)] + (\gamma_2 + \theta_2) y \frac{d}{dy} P_y(y) \\
& = \lambda_2 y P_y(y) + \mu [P_y(y|n_1 = 1) - \pi(1, 0)] \\
& + \mu y^{-1} [P_y(y) - y \pi(0, 1) - \pi(0, 0)] + \gamma_1 y P_y(y|n_1 = 1) \\
& + \theta_1 [P_y(y|n_1 = 1) - \pi(1, 0)] + \theta_2 \left[ \frac{d}{dy} P_y(y) - \pi(0, 1) \right] \\
& [(\lambda_1 + \lambda_2 + \mu) - \mu y^{-1} - \lambda_2 y] P_y(y) + [\gamma_2 y + \theta_2(y - 1)] \frac{d}{dy} P_y(y) \\
& = [\mu + \gamma_1 y + \theta_1] P_y(y|n_1 = 1) \\
& - (\mu + \theta_1) \pi(1, 0) - (\mu + \theta_2) \pi(0, 1) \\
& + [(\lambda_1 + \lambda_2 + \mu) - \mu y^{-1}] \pi(0, 0)
\end{aligned}$$

$$\begin{aligned}
& [(\lambda_1 + \lambda_2 + \mu) - \mu y^{-1} - \lambda_2 y] P_y(y) + [\gamma_2 y + \theta_2(y-1)] \frac{d}{dy} P_y(y) \\
&= [\mu + \gamma_1 y + \theta_1] P_y(y|n_1=1) \\
&\quad + [(\lambda_1 + \lambda_2 + \mu) - \mu y^{-1} - (\lambda_1 + \lambda_2)] \pi(0,0) \\
& [(\lambda_1 + \lambda_2 + \mu) - \mu y^{-1} - \lambda_2 y] P_y(y) + [\gamma_2 y + \theta_2(y-1)] \frac{d}{dy} P_y(y) \\
&= [\mu + \gamma_1 y + \theta_1] P_y(y|n_1=1) + [\mu - \mu y^{-1}] \pi(0,0) \\
& [(\lambda_1 + \lambda_2 + \mu)xy - \mu x - \lambda_2 xy^2] P_y(y) + xy [\gamma_2 y + \theta_2(y-1)] \frac{d}{dy} P_y(y) \\
&= [\mu + \gamma_1 y + \theta_1] xy P_y(y|n_1=1) - \mu x(1-y) \pi(0,0)
\end{aligned} \tag{62}$$

Substituting (61) into (60) eliminates all  $P_x$  terms:

$$\begin{aligned}
& [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 xy^2] P(x, y) \\
&+ xy [\gamma_1(x-y) + \theta_1(x-1)] \frac{\partial}{\partial x} P(x, y) \\
&+ xy [\gamma_2(y-x) + \theta_2(y-1)] \frac{\partial}{\partial y} P(x, y) \\
&= [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_2 xy^2] P_y(y) \\
&\quad + xy [\gamma_2 y + \theta_2(y-1)] \frac{d}{dy} P_y(y) \\
&\quad - xy [\mu + \gamma_1 y + \theta_1] P_y(y|n_1=1)
\end{aligned} \tag{63}$$

Isolating  $[\mu + \gamma_1 y + \theta_1] xy P_y(y|n_1=1)$  from (62) and substituting into (63):

$$\begin{aligned}
& [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 xy^2] P(x, y) \\
&+ xy [\gamma_1(x-y) + \theta_1(x-1)] \frac{\partial}{\partial x} P(x, y) \\
&+ xy [\gamma_2(y-x) + \theta_2(y-1)] \frac{\partial}{\partial y} P(x, y) \\
&= [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_2 xy^2] P_y(y) \\
&\quad + xy [\gamma_2 y + \theta_2(y-1)] \frac{d}{dy} P_y(y) \\
&\quad - \left\{ [(\lambda_1 + \lambda_2 + \mu)xy - \mu x - \lambda_2 xy^2] P_y(y) \right. \\
&\quad \left. + xy [\gamma_2 y + \theta_2(y-1)] \frac{d}{dy} P_y(y) + \mu x(1-y) \pi(0,0) \right\} \\
& [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 xy^2] P(x, y) \\
&+ xy [\gamma_1(x-y) + \theta_1(x-1)] \frac{\partial}{\partial x} P(x, y) \\
&+ xy [\gamma_2(y-x) + \theta_2(y-1)] \frac{\partial}{\partial y} P(x, y) \\
&= \mu(x-y) P_y(y) - \mu x(1-y) \pi(0,0)
\end{aligned} \tag{64}$$

□

#### 4.2.2 Equation for PPGF in state space $\tilde{S}$

**Lemma 2** (Fundamental equation for the PPGF on  $\tilde{S}$ ). *Consider the same system on the state space  $\tilde{S} = \{(n_2, n) : 0 \leq n_2 \leq n\}$ , where  $n$  is the total number of customers waiting in the queues and  $n_2$  the number of waiting class-2 customers. Assume positive recurrence, so that the stationary distribution  $\{\tilde{\pi}(n_2, n)\}$  exists, and define the partial PGF (transforming only  $n_2$ , keeping  $n$  discrete)*

$$\tilde{P}(y, n) = \sum_{n_2=0}^n \tilde{\pi}(n_2, n) y^{n_2},$$

a polynomial of degree  $n$  in  $y$ . Then, for every  $n \geq 1$ , the family  $\{\tilde{P}(y, n)\}_{n \geq 0}$  satisfies the differential-difference equation

$$\begin{aligned}
& [-(\gamma_2 + \gamma_1 y)(1 - y) + (\theta_2 - \theta_1)y] \frac{d}{dy} \tilde{P}(y, n) + [\lambda_1 + \lambda_2 + \mu + \gamma_1 n(1 - y) + \theta_1 n] \tilde{P}(y, n) \\
& = (\lambda_1 + \lambda_2 y) \tilde{P}(y, n - 1) + [\mu + \theta_1(n + 1)] \tilde{P}(y, n + 1) \\
& + (\theta_2 - \theta_1 y) \frac{d}{dy} \tilde{P}(y, n + 1) + \mu y^n (1 - y) \tilde{\pi}(n + 1, n + 1).
\end{aligned} \tag{65}$$

*Proof.* Multiply the interior equation (33) by  $y^{n_2}$  and sum over  $1 \leq n_2 \leq n - 1$ :

$$\begin{aligned}
& (\lambda_1 + \lambda_2 + \mu) \sum_{n_2=1}^{n-1} \tilde{\pi}(n_2, n) y^{n_2} \\
& + \gamma_1 \sum_{n_2=1}^{n-1} (n - n_2) \tilde{\pi}(n_2, n) y^{n_2} \\
& + \gamma_2 \sum_{n_2=1}^{n-1} n_2 \tilde{\pi}(n_2, n) y^{n_2} \\
& + \theta_1 \sum_{n_2=1}^{n-1} (n - n_2) \tilde{\pi}(n_2, n) y^{n_2} \\
& + \theta_2 \sum_{n_2=1}^{n-1} n_2 \tilde{\pi}(n_2, n) y^{n_2} \\
& = \lambda_1 \sum_{n_2=1}^{n-1} \tilde{\pi}(n_2, n - 1) y^{n_2} \\
& + \lambda_2 \sum_{n_2=1}^{n-1} \tilde{\pi}(n_2 - 1, n - 1) y^{n_2} \\
& + \mu \sum_{n_2=1}^{n-1} \tilde{\pi}(n_2, n + 1) y^{n_2} \\
& + \gamma_1 \sum_{n_2=1}^{n-1} (n - n_2 + 1) \tilde{\pi}(n_2 - 1, n) y^{n_2} \\
& + \gamma_2 \sum_{n_2=1}^{n-1} (n_2 + 1) \tilde{\pi}(n_2 + 1, n) y^{n_2} \\
& + \theta_1 \sum_{n_2=1}^{n-1} (n + 1 - n_2) \tilde{\pi}(n_2, n + 1) y^{n_2} \\
& + \theta_2 \sum_{n_2=1}^{n-1} (n_2 + 1) \tilde{\pi}(n_2 + 1, n + 1) y^{n_2}
\end{aligned}$$

Adding the diagonal equation (34) multiplied by  $y^n$  extends most summation limits to  $n$ ; the  $\lambda_1$ - and  $\gamma_2$ -terms on the RHS remain restricted to  $n_2 \leq n - 1$ , and a spare  $\mu \tilde{\pi}(n + 1, n + 1) y^n$

survives:

$$\begin{aligned}
& (\lambda_1 + \lambda_2 + \mu) \sum_{n_2=1}^n \tilde{\pi}(n_2, n) y^{n_2} \\
& + \gamma_1 \sum_{n_2=1}^n (n - n_2) \tilde{\pi}(n_2, n) y^{n_2} \\
& + \gamma_2 \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2} \\
& + \theta_1 \sum_{n_2=1}^n (n - n_2) \tilde{\pi}(n_2, n) y^{n_2} \\
& + \theta_2 \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2} \\
& = \lambda_1 \sum_{n_2=1}^{n-1} \tilde{\pi}(n_2, n-1) y^{n_2} \\
& + \lambda_2 \sum_{n_2=1}^n \tilde{\pi}(n_2-1, n-1) y^{n_2} \\
& + \mu \sum_{n_2=1}^n \tilde{\pi}(n_2, n+1) y^{n_2} \\
& + \mu \tilde{\pi}(n+1, n+1) y^n \\
& + \gamma_1 \sum_{n_2=1}^n (n - n_2 + 1) \tilde{\pi}(n_2-1, n) y^{n_2} \\
& + \gamma_2 \sum_{n_2=1}^{n-1} (n_2 + 1) \tilde{\pi}(n_2+1, n) y^{n_2} \\
& + \theta_1 \sum_{n_2=1}^n (n+1 - n_2) \tilde{\pi}(n_2, n+1) y^{n_2} \\
& + \theta_2 \sum_{n_2=1}^n (n_2 + 1) \tilde{\pi}(n_2+1, n+1) y^{n_2}.
\end{aligned}$$

Evaluating each summand:

$LHS_1$

$$\begin{aligned}
& = (\lambda_1 + \lambda_2 + \mu) \sum_{n_2=1}^n \tilde{\pi}(n_2, n) y^{n_2} \\
& = (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_2=0}^n \tilde{\pi}(n_2, n) y^{n_2} - \tilde{\pi}(0, n) y^0 \right] \\
& = (\lambda_1 + \lambda_2 + \mu) \left[ \tilde{P}(y, n) - \tilde{\pi}(0, n) \right]
\end{aligned} \tag{66}$$

$LHS_2$

$$\begin{aligned}
& = \gamma_1 \sum_{n_2=1}^n (n - n_2) \tilde{\pi}(n_2, n) y^{n_2} \\
& = \gamma_1 \left[ n \sum_{n_2=1}^n \tilde{\pi}(n_2, n) y^{n_2} - \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2} \right] \\
& = \gamma_1 \left[ n \left( \sum_{n_2=0}^n \tilde{\pi}(n_2, n) y^{n_2} - \tilde{\pi}(0, n) y^0 \right) - y \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2-1} \right] \\
& = \gamma_1 \left[ n(\tilde{P}(y, n) - \tilde{\pi}(0, n)) - y \frac{d}{dy} \tilde{P}(y, n) \right]
\end{aligned} \tag{67}$$

$$\begin{aligned}
& LHS_3 \\
&= \gamma_2 \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2} \\
&= \gamma_2 \left[ y \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2-1} \right] \\
&= \gamma_2 y \frac{d}{dy} \tilde{P}(y, n)
\end{aligned} \tag{68}$$

$$\begin{aligned}
& LHS_4 \\
&= \theta_1 \sum_{n_2=1}^n (n - n_2) \tilde{\pi}(n_2, n) y^{n_2} \\
&= \theta_1 \left[ n \left( \tilde{P}(y, n) - \tilde{\pi}(0, n) \right) - y \frac{d}{dy} \tilde{P}(y, n) \right]
\end{aligned} \tag{69}$$

$$\begin{aligned}
& LHS_5 \\
&= \theta_2 \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n) y^{n_2} \\
&= \theta_2 y \frac{d}{dy} \tilde{P}(y, n)
\end{aligned} \tag{70}$$

$$\begin{aligned}
& RHS_1 \\
&= \lambda_1 \sum_{n_2=1}^{n-1} \tilde{\pi}(n_2, n-1) y^{n_2} \\
&= \lambda_1 \left[ \sum_{n_2=1}^n \tilde{\pi}(n_2, n-1) y^{n_2} - \tilde{\pi}(n, n-1) y^n \right] \\
&= \lambda_1 \left[ \sum_{n_2=0}^n \tilde{\pi}(n_2, n-1) y^{n_2} - \tilde{\pi}(n, n-1) y^n - \tilde{\pi}(0, n-1) y^0 \right] \\
&= \lambda_1 \left[ \tilde{P}(y, n-1) - \tilde{\pi}(n, n-1) y^n - \tilde{\pi}(0, n-1) \right]
\end{aligned} \tag{71}$$

$$\begin{aligned}
& RHS_2 \\
&= \lambda_2 \sum_{n_2=1}^n \tilde{\pi}(n_2-1, n-1) y^{n_2} \\
&= \lambda_2 \left[ y \sum_{n_2=1}^n \tilde{\pi}(n_2-1, n-1) y^{n_2-1} \right] \\
&= \lambda_2 \left[ y \sum_{n_2=0}^{n-1} \tilde{\pi}(n_2, n-1) y^{n_2} \right] \\
&= \lambda_2 \left[ y \left( \sum_{n_2=0}^n \tilde{\pi}(n_2, n-1) y^{n_2} - \tilde{\pi}(n, n-1) y^n \right) \right] \\
&= \lambda_2 y \left[ \tilde{P}(y, n-1) - \tilde{\pi}(n, n-1) y^n \right]
\end{aligned} \tag{72}$$

$$\begin{aligned}
& RHS_3 \\
&= \mu \tilde{\pi}(n+1, n+1) y^n + \mu \sum_{n_2=1}^n \tilde{\pi}(n_2, n+1) y^{n_2} \\
&= \mu \left[ \tilde{\pi}(n+1, n+1) y^n + \sum_{n_2=1}^n \tilde{\pi}(n_2, n+1) y^{n_2} \right] \\
&= \mu \left[ \tilde{\pi}(n+1, n+1) y^n + \sum_{n_2=0}^{n+1} \tilde{\pi}(n_2, n+1) y^{n_2} - \tilde{\pi}(0, n+1) y^0 - \tilde{\pi}(n+1, n+1) y^{n+1} \right] \\
&= \mu \left[ \tilde{P}(y, n+1) - \tilde{\pi}(0, n+1) + \tilde{\pi}(n+1, n+1) y^n (1-y) \right]
\end{aligned} \tag{73}$$



$RHS_4$

$$\begin{aligned}
&= \gamma_1 \sum_{n_2=1}^n (n - n_2 + 1) \tilde{\pi}(n_2 - 1, n) y^{n_2} \\
&= \gamma_1 \left[ y \sum_{n_2=1}^n (n - (n_2 - 1)) \tilde{\pi}(n_2 - 1, n) y^{n_2-1} \right] \\
&= \gamma_1 \left[ y \sum_{k=0}^{n-1} (n - k) \tilde{\pi}(k, n) y^k \right] \\
&= \gamma_1 \left[ y \sum_{k=0}^n (n - k) \tilde{\pi}(k, n) y^k \right] \quad (\text{since the } k = n \text{ term is } 0) \\
&= \gamma_1 y \left[ n \sum_{k=0}^n \tilde{\pi}(k, n) y^k - y \sum_{k=1}^n k \tilde{\pi}(k, n) y^{k-1} \right] \\
&= \gamma_1 y \left[ n \tilde{P}(y, n) - y \frac{d}{dy} \tilde{P}(y, n) \right]
\end{aligned} \tag{74}$$

$RHS_5$

$$\begin{aligned}
&= \gamma_2 \sum_{n_2=1}^{n-1} (n_2 + 1) \tilde{\pi}(n_2 + 1, n) y^{n_2} \\
&= \gamma_2 \sum_{k=2}^n k \tilde{\pi}(k, n) y^{k-1} \quad (\text{letting } k = n_2 + 1) \\
&= \gamma_2 \left[ \sum_{k=1}^n k \tilde{\pi}(k, n) y^{k-1} - 1 \cdot \tilde{\pi}(1, n) y^0 \right] \\
&= \gamma_2 \left[ \frac{d}{dy} \tilde{P}(y, n) - \tilde{\pi}(1, n) \right]
\end{aligned} \tag{75}$$

$RHS_6$

$$\begin{aligned}
&= \theta_1 \sum_{n_2=1}^n (n + 1 - n_2) \tilde{\pi}(n_2, n + 1) y^{n_2} \\
&= \theta_1 \left[ (n + 1) \sum_{n_2=1}^n \tilde{\pi}(n_2, n + 1) y^{n_2} - \sum_{n_2=1}^n n_2 \tilde{\pi}(n_2, n + 1) y^{n_2} \right] \\
&= \theta_1 \left[ (n + 1) \left( \sum_{n_2=0}^{n+1} \tilde{\pi}(n_2, n + 1) y^{n_2} - \tilde{\pi}(0, n + 1) - \tilde{\pi}(n + 1, n + 1) y^{n+1} \right) \right] \\
&\quad - \theta_1 \left[ y \left( \sum_{n_2=0}^{n+1} n_2 \tilde{\pi}(n_2, n + 1) y^{n_2-1} - (n + 1) \tilde{\pi}(n + 1, n + 1) y^n \right) \right] \\
&= \theta_1 \left[ (n + 1) (\tilde{P}(y, n + 1) - \tilde{\pi}(0, n + 1)) - (n + 1) \tilde{\pi}(n + 1, n + 1) y^{n+1} \right] \\
&\quad - \theta_1 \left[ y \frac{d}{dy} \tilde{P}(y, n + 1) - (n + 1) \tilde{\pi}(n + 1, n + 1) y^{n+1} \right] \\
&= \theta_1 \left[ (n + 1) (\tilde{P}(y, n + 1) - \tilde{\pi}(0, n + 1)) - y \frac{d}{dy} \tilde{P}(y, n + 1) \right]
\end{aligned} \tag{76}$$

$RHS_7$

$$\begin{aligned}
&= \theta_2 \sum_{n_2=1}^n (n_2 + 1) \tilde{\pi}(n_2 + 1, n + 1) y^{n_2} \\
&= \theta_2 \sum_{k=2}^{n+1} k \tilde{\pi}(k, n + 1) y^{k-1} \quad (\text{letting } k = n_2 + 1) \\
&= \theta_2 \left[ \sum_{k=1}^{n+1} k \tilde{\pi}(k, n + 1) y^{k-1} - 1 \cdot \tilde{\pi}(1, n + 1) y^0 \right] \\
&= \theta_2 \left[ \frac{d}{dy} \tilde{P}(y, n + 1) - \tilde{\pi}(1, n + 1) \right]
\end{aligned} \tag{77}$$

Collecting:

$$\begin{aligned}
&(\lambda_1 + \lambda_2 + \mu) \left[ \tilde{P}(y, n) - \tilde{\pi}(0, n) \right] \\
&+ \gamma_1 \left[ n(\tilde{P}(y, n) - \tilde{\pi}(0, n)) - y \frac{d}{dy} \tilde{P}(y, n) \right] \\
&+ \gamma_2 y \frac{d}{dy} \tilde{P}(y, n) \\
&+ \theta_1 \left[ n(\tilde{P}(y, n) - \tilde{\pi}(0, n)) - y \frac{d}{dy} \tilde{P}(y, n) \right] \\
&+ \theta_2 y \frac{d}{dy} \tilde{P}(y, n) \\
&= \lambda_1 \left[ \tilde{P}(y, n - 1) - \tilde{\pi}(n, n - 1) y^n - \tilde{\pi}(0, n - 1) \right] \\
&+ \lambda_2 y \left[ \tilde{P}(y, n - 1) - \tilde{\pi}(n, n - 1) y^n \right] \\
&+ \mu \left[ \tilde{P}(y, n + 1) - \tilde{\pi}(0, n + 1) + \tilde{\pi}(n + 1, n + 1) y^n (1 - y) \right] \\
&+ \gamma_1 y \left[ n \tilde{P}(y, n) - y \frac{d}{dy} \tilde{P}(y, n) \right] \\
&+ \gamma_2 \left[ \frac{d}{dy} \tilde{P}(y, n) - \tilde{\pi}(1, n) \right] \\
&+ \theta_1 \left[ (n + 1) (\tilde{P}(y, n + 1) - \tilde{\pi}(0, n + 1)) - y \frac{d}{dy} \tilde{P}(y, n + 1) \right] \\
&+ \theta_2 \left[ \frac{d}{dy} \tilde{P}(y, n + 1) - \tilde{\pi}(1, n + 1) \right]
\end{aligned}$$

Regrouping:

$$\begin{aligned}
&[-(\gamma_2 + \gamma_1 y)(1 - y) + (\theta_2 - \theta_1)y] \frac{d}{dy} \tilde{P}(y, n) + [\lambda_1 + \lambda_2 + \mu + \gamma_1 n(1 - y) + \theta_1 n] \tilde{P}(y, n) \\
&= (\lambda_1 + \lambda_2 y) \tilde{P}(y, n - 1) + [\mu + \theta_1(n + 1)] \tilde{P}(y, n + 1) + (\theta_2 - \theta_1 y) \frac{d}{dy} \tilde{P}(y, n + 1) \\
&+ \{[\lambda_1 + \lambda_2 + \mu + (\gamma_1 + \theta_1)n] \tilde{\pi}(0, n) - \lambda_1 \tilde{\pi}(0, n - 1) - [\mu + \theta_1(n + 1)] \tilde{\pi}(0, n + 1) \\
&\quad - \gamma_2 \tilde{\pi}(1, n) - \theta_2 \tilde{\pi}(1, n + 1)\} \\
&- y^n (\lambda_1 + \lambda_2 y) \tilde{\pi}(n, n - 1) + \mu y^n (1 - y) \tilde{\pi}(n + 1, n + 1)
\end{aligned}$$

The  $y$ -boundary equation (35) makes the bracketed terms vanish; since  $\tilde{\pi}(n, n - 1) = 0$  (outside  $\tilde{S}$ ):

$$\begin{aligned}
&[-(\gamma_2 + \gamma_1 y)(1 - y) + (\theta_2 - \theta_1)y] \frac{d}{dy} \tilde{P}(y, n) + [\lambda_1 + \lambda_2 + \mu + \gamma_1 n(1 - y) + \theta_1 n] \tilde{P}(y, n) \\
&= (\lambda_1 + \lambda_2 y) \tilde{P}(y, n - 1) + [\mu + \theta_1(n + 1)] \tilde{P}(y, n + 1) + (\theta_2 - \theta_1 y) \frac{d}{dy} \tilde{P}(y, n + 1) \\
&+ \mu y^n (1 - y) \tilde{\pi}(n + 1, n + 1)
\end{aligned} \tag{78}$$

□

## 5 Analysis of Model-A ( $\gamma_1 = \gamma_2 = 0$ , $\theta_1 = \theta_2 = 0$ )

Model-A is the baseline of this collection of models: all jockeying and abandonment parameters are set to zero, leaving an ordinary two-class  $M/M/1$  queueing model with non-preemptive priority. The analysis is carried out in both state spaces  $S$  and  $\tilde{S}$ : on the former, two complete proofs are presented; while on the latter, an approximate solution and its verification are provided.

The balance equations are obtained by plugging  $\gamma_1 = \gamma_2 = 0$  and  $\theta_1 = \theta_2 = 0$  into Equations (28), (30), (29), (31), and (32) for state space  $S$ , and Equations (33), (34), (35), (36), and (37) for state space  $\tilde{S}$ . The corresponding starting points are Lemma 1 for  $S$  and Lemma 2 for  $\tilde{S}$ .

**Stability.** Since there is no abandonment, the system is positive recurrent if and only if the total traffic intensity satisfies

$$\rho = \frac{\lambda_1 + \lambda_2}{\mu} < 1. \quad (79)$$

Figure 4 shows the queueing diagram of Model-A.




Figure 4: Queue layout for Model-A. Only Poisson arrivals and exponential service are present; all jockeying and abandonment rates are zero.

### 5.1 Analysis of Model-A: state space $S$

We begin from Equation (45) with  $\gamma_1 = \gamma_2 = \theta_1 = \theta_2 = 0$ , in which the marginal  $P_y(y) = P(0, y)$  is still to be determined:

$$[(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) = \mu(x - y)P_y(y) - \mu x(1 - y)\pi(0, 0). \quad (80)$$

The following theorem summarises the main result of this subsection; two standalone proofs are given, one analytical and one probabilistic.

**Theorem 1.** *Consider a queueing system with two priority classes with Poisson arrival rates  $\lambda_1$  and  $\lambda_2$ , where class-1 customers have non-preemptive priority over those of class 2, and exponentially distributed service times with rate  $\mu$  independently of the class. Under the stability condition (79), the joint PGF  $P(x, y) = \mathbb{E}[x^{N_1} y^{N_2}]$  is*

$$P(x, y) = [(1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2]^{-1} \cdot y(1 - y) \left[ \frac{x - x^*(y)}{x^*(y) - y} \right] \cdot \rho(1 - \rho), \quad (81)$$

where

$$x^*(y) = \frac{[\mu + \lambda_1 + \lambda_2(1 - y)] - \sqrt{[\mu + \lambda_1 + \lambda_2(1 - y)]^2 - 4\lambda_1\mu}}{2\lambda_1}. \quad (82)$$

**Corollary 1.** *Under the stability condition (79), the empty-state probabilities for Model-A are*

$$\pi_0 = 1 - \rho, \quad \pi(0, 0) = \rho(1 - \rho). \quad (83)$$

In particular,  $P(1, 1) = 1 - \pi_0 = \rho$  and the diagonal generating function satisfies  $P(z, z) = \pi(0, 0)/(1 - \rho z)$ .

*Proof of Corollary 1.* Setting  $x = y = z$  in the fundamental equation (80), the boundary term  $\mu(x - y)P_y(y)$  vanishes:

$$[(\lambda_1 + \lambda_2 + \mu)z^2 - \mu z - \lambda_1 z^3 - \lambda_2 z^3] P(z, z) = -\mu z(1 - z)\pi(0, 0).$$

Dividing both sides by  $z$  and factoring the bracket:

$$-(1-z)[\mu - (\lambda_1 + \lambda_2)z]P(z, z) = -\mu(1-z)\pi(0, 0).$$

Cancelling  $(1-z) \neq 0$  for  $z \in [0, 1)$  and rearranging:

$$P(z, z) = \frac{\mu \pi(0, 0)}{\mu - \lambda z} = \frac{\pi(0, 0)}{1 - \rho z}, \quad (84)$$

where  $\lambda = \lambda_1 + \lambda_2$  and the second equality follows by dividing numerator and denominator by  $\mu$ .

Setting  $z = 1$  and using the normalisation  $P(1, 1) = 1 - \pi_0$ :

$$1 - \pi_0 = \frac{\pi(0, 0)}{1 - \rho}. \quad (85)$$

The global balance equation for the idle state gives  $\lambda \pi_0 = \mu \pi(0, 0)$ , i.e.  $\pi(0, 0) = \rho \pi_0$ . Substituting into (85):

$$1 - \pi_0 = \frac{\rho \pi_0}{1 - \rho} \implies (1 - \pi_0)(1 - \rho) = \rho \pi_0 \implies \pi_0 = 1 - \rho,$$

and therefore  $\pi(0, 0) = \rho(1 - \rho)$ .  $\square$

### 5.1.1 Analytical approach for Model-A in state space $S$

*Proof.* We begin from the fundamental equation for Model-A (80):

$$[(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) = \mu(x - y)P_y(y) - \mu x(1 - y)\pi(0, 0).$$

We apply the *Kernel Method*: fix  $y \in (0, 1]$  and find  $x^*(y)$  such that the LHS of the expression above is zero. Since  $P(x, y)$  is a PGF, it is bounded. At any such root the LHS vanishes, so the RHS must also vanish:

$$\mu[(x^*(y) - y)P_y(y) - x^*(y)(1 - y)\pi(0, 0)] = 0.$$

To find the roots of the coefficient of  $P(x, y)$ , divide by  $y \neq 0$  and set

$$\begin{aligned} 0 &= (\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 x y \\ 0 &= \lambda_1 x^2 - [\mu + \lambda_1 + \lambda_2(1 - y)]x + \mu, \end{aligned}$$

whose roots are

$$x^\pm(y) = \frac{[\mu + \lambda_1 + \lambda_2(1 - y)] \pm \sqrt{[\mu + \lambda_1 + \lambda_2(1 - y)]^2 - 4\mu\lambda_1}}{2\lambda_1}.$$

We label the root with the negative sign  $x^*(y)$  and the root with the positive sign  $x^+(y)$ . At  $y = 1$ :

$$x^*(1) = 1 \quad \text{and} \quad x^+(1) = \mu/\lambda_1 > 1.$$

We require our root to lie inside the closed unit disk for  $y \in [0, 1)$ . Computing the derivative of  $x^*(y)$ :

$$\frac{d}{dy}x^*(y) = \frac{\lambda_2}{2\lambda_1} \left[ -1 + \frac{\mu + \lambda_1 + \lambda_2(1 - y)}{\sqrt{(\mu + \lambda_1 + \lambda_2(1 - y))^2 - 4\mu\lambda_1}} \right] > 0, \quad (86)$$

since  $(\mu + \lambda_1 + \lambda_2(1 - y))^2 > (\mu + \lambda_1 + \lambda_2(1 - y))^2 - 4\mu\lambda_1$  whenever  $\lambda_1\mu > 0$ . Therefore  $x^*(y)$  is strictly increasing on  $[0, 1]$ , and since  $x^*(1) = 1$  we have  $x^*(y) < 1$  for  $y \in [0, 1)$ . By Vieta's formulas for the kernel polynomial  $\lambda_1 x^2 - [\mu + \lambda_1 + \lambda_2(1 - y)]x + \mu = 0$ , the product of the two roots satisfies  $x^*(y)x^+(y) = \mu/\lambda_1 > 1$ , so  $x^+(y) = \mu/(\lambda_1 x^*(y)) > 1$  throughout  $[0, 1)$ . Hence the only root strictly inside the unit disk for  $y \in [0, 1)$  is  $x^*(y)$ :

$$x^*(y) = \frac{(\mu + \lambda_1 + \lambda_2(1 - y)) - \sqrt{(\mu + \lambda_1 + \lambda_2(1 - y))^2 - 4\mu\lambda_1}}{2\lambda_1}. \quad (87)$$

Since subsequent steps involve L'Hôpital's rule at  $y \rightarrow 1$ , we evaluate  $\frac{d}{dy}x^*(y)$  from (86) at  $y = 1$ .

Under stability,  $\rho_1 = \lambda_1/\mu < \rho < 1$  implies  $\mu > \lambda_1$ , so  $\sqrt{(\mu - \lambda_1)^2} = \mu - \lambda_1$ :

$$\frac{d}{dy}x^*(1) = \frac{\lambda_2}{2\lambda_1} \left[ -1 + \frac{\mu + \lambda_1}{\sqrt{(\mu + \lambda_1)^2 - 4\lambda_1\mu}} \right] = \frac{\lambda_2}{2\lambda_1} \left[ -1 + \frac{\mu + \lambda_1}{\sqrt{(\mu - \lambda_1)^2}} \right] = \frac{\lambda_2}{2\lambda_1} \left[ \frac{\mu + \lambda_1 - (\mu - \lambda_1)}{\mu - \lambda_1} \right] = \frac{\lambda_2}{\mu - \lambda_1}.$$

For future reference:

$$\frac{d}{dy}x^*(1) = \frac{\lambda_2}{\mu - \lambda_1} = \frac{\rho_2}{1 - \rho_1}. \quad (88)$$

Returning to the fundamental equation (80) at  $x = x^*(y)$ , the RHS also equals zero, giving

$$\mu \cdot [(x^*(y) - y)P_y(y) - x^*(y)(1 - y)\pi(0, 0)] = 0 \implies P_y(y) = \frac{x^*(y)(1 - y)}{x^*(y) - y}\pi(0, 0). \quad (89)$$

By Corollary 1,  $\pi(0, 0) = \rho(1 - \rho)$ . Substituting this and Equation (89) into the fundamental equation (80):

$$\begin{aligned} [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) &= \mu(x - y) \frac{x^*(y)(1 - y)}{x^*(y) - y} \rho(1 - \rho) - \mu x(1 - y)\rho(1 - \rho) \\ [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) &= \mu(1 - y) \left[ \frac{x^*(y)(x - y)}{x^*(y) - y} - x \right] \rho(1 - \rho) \\ [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) &= \mu(1 - y) \left[ \frac{y(x - x^*(y))}{x^*(y) - y} \right] \rho(1 - \rho) \\ [(1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2] P(x, y) &= y(1 - y) \left[ \frac{x - x^*(y)}{x^*(y) - y} \right] \rho(1 - \rho). \end{aligned}$$

Multiplying both sides by  $[(1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2]^{-1}$  isolates  $P(x, y)$  and concludes the proof.  $\square$

### 5.1.2 Probabilistic approach for Model-A in state space $S$

*Proof.* The proof is similar to the analytical approach, but deriving the class-2 marginal PGF  $P_y(y)$  by means of probabilistic arguments. We begin by stating the fundamental global balance equation for Model-A (Equation (80)):

$$[(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) = \mu[(x - y)P_y(y) - x(1 - y)\pi(0, 0)]. \quad (90)$$

**Class-2 conditional distribution via  $M/G/1$ .** Since class 1 has non-preemptive priority, class-2 customers will not be taken into service whenever  $N_1 > 0$ ; their *effective service time*, namely  $B$ , is therefore the duration of a complete Busy Period, of the single-class  $M/M/1$  sub-queue with arrival rate  $\lambda_1$  and service rate  $\mu$ , so from class-2's viewpoint the system is an  $M/G/1$  queue with arrival rate  $\lambda_2$  and generally distributed service times  $B$  with mean  $\mathbb{E}[B]$ . The LST and mean of  $B$  are presented in §3.1 (Equations (2)–(3)), now with  $\lambda_1$  instead of  $\lambda$ :

$$\tilde{B}(s) = \frac{(\mu + \lambda_1 + s) - \sqrt{(\mu + \lambda_1 + s)^2 - 4\mu\lambda_1}}{2\lambda_1}, \quad (91)$$

$$\mathbb{E}[B] = \frac{1}{\mu(1 - \rho_1)}. \quad (92)$$

Applying the Pollaczek–Khinchine (PK) formula (§3.4, Equation (12)) to this  $M/G/1$  system, with  $\tilde{B}(\lambda_2(1 - y)) = x^*(y)$  (cf. (87)) and  $\lambda_2 \mathbb{E}[B] = \rho_2/(1 - \rho_1)$ :

$$L_2(y) = \frac{(1 - \lambda_2 \mathbb{E}[B])(1 - y)\tilde{B}(\lambda_2(1 - y))}{\tilde{B}(\lambda_2(1 - y)) - y} = \frac{1 - \rho}{1 - \rho_1} \cdot \frac{x^*(y)(1 - y)}{x^*(y) - y}. \quad (93)$$

By the PASTA property, the time-average distribution of  $N_2$  conditional on (busy,  $N_1 = 0$ ) coincides with the  $M/G/1$  steady-state distribution, so  $\mathbb{E}[y^{N_2} \mid \text{busy}, N_1 = 0] = L_2(y)$ .

By the law of total expectation,  $P_y(y) = \mathbb{P}(\text{busy}, N_1 = 0) \cdot \mathbb{E}[y^{N_2} \mid \text{busy}, N_1 = 0]$ . So

$$P_y(y) = \mathbb{P}(\text{busy}, N_1 = 0) \cdot \mathbb{E}[y^{N_2} \mid \text{busy}, N_1 = 0] = \mathbb{P}(\text{busy}, N_1 = 0) \cdot L_2(y).$$

We now can obtain  $\mathbb{P}(\text{busy}, N_1 = 0)$ , from  $\mathbb{P}(\text{busy}) = \rho$  and  $\mathbb{P}(N_1 = 0 \mid \text{busy}) = 1 - \rho_1$ . The latter holds because, under non-preemptive priority with a common service rate  $\mu$ , the class-1 queue-plus-service process is exactly the queue-length process of an autonomous  $M/M/1(\lambda_1, \mu)$ : a class-1 arrival and a class-1 departure occur at rates  $\lambda_1$  and  $\mu$  regardless of the class currently in service, since the residual service time is  $\text{Exp}(\mu)$  by memorylessness whether the server holds

a class-1 or class-2 customer (cf. §3.2); the class-2 occupancy is thus exchangeable with class-1 occupancy for the purpose of counting class-1 customers, so the count obeys the geometric law and  $\mathbb{P}(N_1 = 0 \mid \text{busy}) = 1 - \rho_1$ . So

$$\mathbb{P}(\text{busy}, N_1 = 0) = \rho(1 - \rho_1)$$

and therefore

$$P_y(y) = \rho(1 - \rho_1) \cdot L_2(y) = \rho(1 - \rho_1) \cdot \frac{1 - \rho}{1 - \rho_1} \cdot \frac{x^*(y)(1 - y)}{x^*(y) - y} = \frac{x^*(y)(1 - y)}{x^*(y) - y} \cdot (1 - \rho)\rho \quad (94)$$

Substituting (94) and  $\pi(0, 0) = \rho(1 - \rho)$  (Corollary 1) into (90):

$$\begin{aligned} [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) &= \mu(x - y) \frac{x^*(y)(1 - y)}{x^*(y) - y} \rho(1 - \rho) - \mu x(1 - y) \rho(1 - \rho) \\ [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) &= \mu(1 - y) \left[ \frac{x^*(y)(x - y)}{x^*(y) - y} - x \right] \rho(1 - \rho) \\ [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) &= \mu(1 - y) \left[ \frac{y(x - x^*(y))}{x^*(y) - y} \right] \rho(1 - \rho) \\ [(1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2] P(x, y) &= y(1 - y) \left[ \frac{x - x^*(y)}{x^*(y) - y} \right] \rho(1 - \rho). \end{aligned}$$

Multiplying both sides by  $[(1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2]^{-1}$  isolates  $P(x, y)$  and concludes the proof.  $\square$

### 5.1.3 Partial-PGF recurrence approach for Model-A in state space $S$

This proof follows Cohen [Coh56]. It consists in keeping the class-1 index  $n_1$  as a discrete parameter and forming a generating function only with respect to  $N_2$ , namely the PPGF  $\tilde{P}(n_1, y) = \sum_{n_2 \geq 0} \pi(n_1, n_2) y^{n_2}$  introduced in (42). Multiplying the balance equations by  $y^{n_2}$  and summing over  $n_2$  turns the two-dimensional balance equations into a second-order linear recurrence in  $n_1$ , whose characteristic roots fix the geometric structure of the solution in the priority class index.

*Proof.* We start by stating the balance equations for Model-A, with all jockeying and abandonment parameters set to zero (i.e.  $\gamma_1 = \gamma_2 = \theta_1 = \theta_2 = 0$ ) in the interior balance equation (Equation (28)) and our  $x$ -boundary equation (Equation (30)). We have

$$(\lambda_1 + \lambda_2 + \mu) \pi(n_1, n_2) = \mu \pi(n_1 + 1, n_2) + \lambda_1 \pi(n_1 - 1, n_2) + \lambda_2 \pi(n_1, n_2 - 1), \quad n_1 \geq 1, n_2 \geq 1, \quad (95)$$

$$(\lambda_1 + \lambda_2 + \mu) \pi(n_1, 0) = \mu \pi(n_1 + 1, 0) + \lambda_1 \pi(n_1 - 1, 0), \quad n_1 \geq 1, n_2 = 0. \quad (96)$$

$$\begin{aligned} & (\lambda_1 + \lambda_2 + \mu) \sum_{n_2=1}^{\infty} \pi(n_1, n_2) y^{n_2} \\ &= \mu \sum_{n_2=1}^{\infty} \pi(n_1 + 1, n_2) y^{n_2} + \lambda_1 \sum_{n_2=1}^{\infty} \pi(n_1 - 1, n_2) y^{n_2} + \lambda_2 \sum_{n_2=1}^{\infty} \pi(n_1, n_2 - 1) y^{n_2} \\ LHS_1 &= RHS_1 + RHS_2 + RHS_3 \\ LHS_1 &= (\lambda_1 + \lambda_2 + \mu) \sum_{n_2=1}^{\infty} \pi(n_1, n_2) y^{n_2} \\ &= (\lambda_1 + \lambda_2 + \mu) \left[ \sum_{n_2=0}^{\infty} \pi(n_1, n_2) y^{n_2} - \pi(n_1, 0) y^0 \right] \\ &= (\lambda_1 + \lambda_2 + \mu) \left[ \tilde{P}(n_1, y) - \pi(n_1, 0) \right] \end{aligned}$$

$$\begin{aligned}
& RHS_1 \\
&= \mu \sum_{n_2=1}^{\infty} \pi(n_1+1, n_2) y^{n_2} \\
&= \mu \left[ \sum_{n_2=0}^{\infty} \pi(n_1+1, n_2) y^{n_2} - \pi(n_1+1, 0) y^0 \right] \\
&= \mu \left[ \tilde{P}(n_1+1, y) - \pi(n_1+1, 0) \right] \\
& RHS_2 \\
&= \lambda_1 \sum_{n_2=1}^{\infty} \pi(n_1-1, n_2) y^{n_2} \\
&= \lambda_1 \left[ \sum_{n_2=0}^{\infty} \pi(n_1-1, n_2) y^{n_2} - \pi(n_1-1, 0) y^0 \right] \\
&= \lambda_1 \left[ \tilde{P}(n_1-1, y) - \pi(n_1-1, 0) \right] \\
& RHS_3 \\
&= \lambda_2 \sum_{n_2=1}^{\infty} \pi(n_1, n_2-1) y^{n_2} \\
&= \lambda_2 y \sum_{n_2=1}^{\infty} \pi(n_1, n_2-1) y^{n_2-1} \\
&= \lambda_2 y \sum_{m=0}^{\infty} \pi(n_1, m) y^m \\
&= \lambda_2 y \tilde{P}(n_1, y) \\
& (\lambda_1 + \lambda_2 + \mu) [\tilde{P}(n_1, y) - \pi(n_1, 0)] \\
&= \mu [\tilde{P}(n_1+1, y) - \pi(n_1+1, 0)] \\
&\quad + \lambda_1 [\tilde{P}(n_1-1, y) - \pi(n_1-1, 0)] \\
&\quad + \lambda_2 y \tilde{P}(n_1, y) \\
& [\mu + \lambda_1 + \lambda_2(1-y)] \tilde{P}(n_1, y) - \mu \tilde{P}(n_1+1, y) - \lambda_1 \tilde{P}(n_1-1, y) \\
&= (\lambda_1 + \lambda_2 + \mu) \pi(n_1, 0) - \mu \pi(n_1+1, 0) - \lambda_1 \pi(n_1-1, 0)
\end{aligned} \tag{97}$$

Note that, using our  $x$ -boundary balance equation (Equation (96)), the right-hand side of the expression above (Equation (97)) vanishes completely. We obtain the homogeneous recurrence

$$[\mu + \lambda_1 + \lambda_2(1-y)] \tilde{P}(n_1, y) = \mu \tilde{P}(n_1+1, y) + \lambda_1 \tilde{P}(n_1-1, y), \quad n_1 \geq 1. \tag{98}$$

**Characteristic roots and mode selection.** Equation (98) is a *second-order linear recurrence* in  $n_1$  with constant coefficients in  $n_1$ . So its general solution is a linear combination of two geometric sequences  $[\tilde{x}^{\pm}(y)]^{n_1}$ , where  $\tilde{x}^{\pm}(y)$  solve the characteristic equation

$$\mu \tilde{x}^2 - [\mu + \lambda_1 + \lambda_2(1-y)] \tilde{x} + \lambda_1 = 0, \tag{99}$$

that is,

$$\tilde{x}^{\pm}(y) = \frac{(\mu + \lambda_1 + \lambda_2(1-y)) \pm \sqrt{(\mu + \lambda_1 + \lambda_2(1-y))^2 - 4\lambda_1\mu}}{2\mu}. \tag{100}$$

The *characteristic polynomial* (Equation (99)) is the reciprocal to that of the bivariate PGF approaches (i.e. the reciprocal of  $\lambda_1 x^2 - [\mu + \lambda_1 + \lambda_2(1-y)]x + \mu$ ), thus its roots are the reciprocals of that kernel's roots: the positive root is the reciprocal of the negative one ( $\tilde{x}^+(y) = 1/x^*(y)$ ), and conversely  $\tilde{x}^-(y) = 1/\tilde{x}^+(y)$ . By Vieta's formulas,  $\tilde{x}^-(y) \tilde{x}^+(y) = \lambda_1/\mu = \rho_1 < 1$ . Recall from §5.1.1 that  $x^*(y) < 1 < x^+(y)$  for  $y \in [0, 1]$ . The selection of the negative-sign root is motivated by the fact that we need the sequence  $\{\tilde{P}(n_1, y)\}_{n_1 \geq 0}$  to be summable. In fact, we need

$$\sum_{n_1=0}^{\infty} \tilde{P}(n_1, y) = P(1, y) \leq P(1, 1) = 1 - \pi_0 = \rho < \infty.$$

At  $y = 1$ , we have

$$\tilde{x}^-(1) = \rho_1 \quad \text{and} \quad \tilde{x}^+(1) = 1,$$

the latter making the geometric sequence  $[\tilde{x}^+(y)]^{n_1}$  non-summable. We therefore define  $\tilde{x}^*(y) := \tilde{x}^-(y)$ ,

$$\tilde{x}^*(y) := \tilde{x}^-(y) = \frac{(\mu + \lambda_1 + \lambda_2(1-y)) - \sqrt{(\mu + \lambda_1 + \lambda_2(1-y))^2 - 4\lambda_1\mu}}{2\mu} = \rho_1 x^*(y). \quad (101)$$

The last identity holds because  $\tilde{x}^*$  and  $x^*$  share the same numerator and differ only in the denominator ( $2\mu$  versus  $2\lambda_1$ ). Next,

$$\tilde{P}(n_1, y) = \tilde{P}(0, y) [\tilde{x}^*(y)]^{n_1}, \quad n_1 \geq 0, \quad (102)$$

where the relation also holds at  $n_1 = 0$  trivially. Recall that at  $y = 1$ , we have  $\tilde{x}^*(1) = \lambda_1/\mu = \rho_1$ . Thus,  $\tilde{P}(n_1, 1) = \tilde{P}(0, 1) \rho_1^{n_1}$  and  $\tilde{P}(0, 1) = P_y(1) = \rho(1 - \rho_1)$ . The constant  $\tilde{P}(0, y)$  is resolved by its definition

$$\tilde{P}(0, y) = \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2} = P(0, y) = P_y(y),$$

as determined in the bivariate PGF approaches. Summing the geometric law (Equation (102)) over  $n_1 \geq 0$  for  $x \in (0, 1]$  recovers

$$P(x, y) = \sum_{n_1=0}^{\infty} \tilde{P}(n_1, y) x^{n_1} = P_y(y) \sum_{n_1=0}^{\infty} [\tilde{x}^*(y) x]^{n_1} = \frac{P_y(y)}{1 - \tilde{x}^*(y) x}. \quad (103)$$

Now, in order to bring this to the form stated in Theorem 1, we define the kernel of the fundamental equation divided by  $\mu$  as

$$K(x, y) := (1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2 = -y\rho_1(x - x^*(y))(x - x^+(y)), \quad (104)$$

where  $x^*(y)$  and  $x^+(y)$  are the roots found in §5.1.1. By Vieta's product formula,  $x^*(y)x^+(y) = \mu/\lambda_1 = 1/\rho_1$ . This, together with  $\tilde{x}^*(y)$  (Equation (101)), gives  $\tilde{x}^*(y) = \rho_1 x^*(y) = 1/x^+(y)$ . Now, looking at the denominator of our reconstructed PGF (Equation (103)), and using the factorisation of  $K(x, y)$  (Equation (104)), we see that

$$1 - \tilde{x}^*(y) x = 1 - \frac{x}{x^+(y)} = \frac{x^+(y) - x}{x^+(y)} = \frac{K(x, y)}{y\rho_1 x^+(y)(x - x^*(y))}.$$

Substituting this into our reconstructed  $P(x, y)$  (Equation (103)), using  $P_y(y) = \frac{x^*(y)(1-y)}{x^*(y)-y} \rho(1-\rho)$  and also noting that  $\rho_1 x^*(y) x^+(y) = 1$ , we obtain

$$\begin{aligned} P(x, y) &= \frac{x^*(y)(1-y)}{x^*(y)-y} \rho(1-\rho) \cdot \frac{y\rho_1 x^+(y)(x - x^*(y))}{K(x, y)} \\ &= \underbrace{\rho_1 x^*(y) x^+(y)}_{=1} \frac{1}{K(x, y)} y(1-y) \frac{x - x^*(y)}{x^*(y)-y} \rho(1-\rho) \\ &= [(1 + \rho_1 + \rho_2)xy - y - \rho_1 x^2 y - \rho_2 x y^2]^{-1} \cdot y(1-y) \frac{x - x^*(y)}{x^*(y)-y} \rho(1-\rho), \end{aligned}$$

which recovers Equation (81) from Theorem 1 and concludes the proof.  $\square$

## 5.2 Analysis of Model-A on $\tilde{S}$ : scope and limitations of the partial-PGF recursion

Setting  $\gamma_1 = \gamma_2 = \theta_1 = \theta_2 = 0$  in the  $\tilde{S}$  balance equations (33)–(37) and forming the PPGF  $\tilde{P}(y, n) = \sum_{n_2=0}^n \tilde{\pi}(n_2, n) y^{n_2}$  yields the inhomogeneous recurrence:

$$\begin{aligned} &[\lambda_1 + \lambda_2 + \mu] \tilde{P}(y, n) \\ &= (\lambda_1 + \lambda_2 y) \tilde{P}(y, n-1) + \mu \tilde{P}(y, n+1) + \mu y^n (1-y) \tilde{\pi}(n+1, n+1) \end{aligned} \quad (105)$$

The forcing term in (105) is controlled by the diagonal probabilities  $\tilde{\pi}(n+1, n+1) = \pi(0, n+1)$ , whose decay we first establish from the exact  $S$ -solution.



**Lemma 3** (Geometric decay of the boundary probabilities). *For Model-A the boundary probabilities  $\pi(0, n)$  satisfy  $\pi(0, n) = O(r^{-n})$  for every  $r < 1/\rho$ ; equivalently they decay geometrically with rate  $\rho$ .*

*Proof.* By definition  $P_y(y) = P(0, y) = \sum_{n \geq 0} \pi(0, n) y^n$  is the generating function of  $\{\pi(0, n)\}$ , and from (89) with  $\pi(0, 0) = \rho(1 - \rho)$ ,

$$P_y(y) = \rho(1 - \rho) \frac{x^*(y)(1 - y)}{x^*(y) - y}.$$

Evaluating the kernel quadratic (82) at  $x = y$  gives  $\lambda_1 y^2 - [\mu + \lambda_1 + \lambda_2(1 - y)]y + \mu = (\lambda y - \mu)(y - 1)$  with  $\lambda = \lambda_1 + \lambda_2$ , so  $x^*(y) = y$  holds only at  $y = 1$  and  $y = \mu/\lambda = 1/\rho$ . At  $y = 1$  the simple zeros of  $(1 - y)$  and  $(x^*(y) - y)$  cancel and  $P_y$  is analytic with  $P_y(1) = \rho$ ; the branch point of  $x^*$  lies beyond  $1/\rho$ . Hence the singularity of  $P_y$  nearest the origin is the simple pole at  $y = 1/\rho > 1$ , so  $P_y$  is analytic on the disc  $|y| < 1/\rho$ . By the Cauchy coefficient estimates, for every  $r < 1/\rho$  there is  $C_r < \infty$  with  $\pi(0, n) \leq C_r r^{-n}$ , i.e. geometric decay at rate  $\rho$ .  $\square$

**Approximation 1** (Heuristic PPGF on  $\tilde{S}$ ). *Neglecting the boundary-diagonal forcing term  $\mu y^n(1 - y)\tilde{\pi}(n + 1, n + 1)$  in (105) — which is exponentially small by Lemma 3 — the homogeneous recurrence admits the geometric solution*

$$\tilde{P}(y, n) \approx \rho(1 - \rho) [\tilde{y}^*(y)]^n, \quad (106)$$

where

$$\tilde{y}^*(y) = \frac{(\lambda + \mu) - \sqrt{(\lambda + \mu)^2 - 4\mu(\lambda_1 + \lambda_2 y)}}{2\mu}, \quad \lambda = \lambda_1 + \lambda_2. \quad (107)$$

At  $y = 1$  this recovers  $\tilde{P}(1, n) = (1 - \rho)\rho^{n+1}$ , consistent with the  $M/M/1(\lambda, \mu)$  steady-state marginal for  $n$  customers waiting.

*Justification of Approximation 1.* The inhomogeneous term  $\mu y^n(1 - y)\tilde{\pi}(n + 1, n + 1)$  prevents the direct application of a geometric-sequence ansatz. The diagonal probabilities  $\tilde{\pi}(n, n) = \pi(0, n)$  correspond to states where all  $n$  waiting customers are class 2, and by Lemma 3 they decay geometrically in  $n$  at rate  $\rho$ ; the forcing term is therefore exponentially small, and neglecting it yields an approximation whose accuracy is quantified below. The resulting homogeneous recurrence is

$$(\lambda_1 + \lambda_2 + \mu)\tilde{P}(y, n) = (\lambda_1 + \lambda_2 y)\tilde{P}(y, n - 1) + \mu\tilde{P}(y, n + 1). \quad (108)$$

The characteristic polynomial of this recurrence is

$$\mu[\tilde{y}(y)]^2 - (\lambda_1 + \lambda_2 + \mu)\tilde{y}(y) + (\lambda_1 + \lambda_2 y) = 0, \quad (109)$$

with roots

$$\tilde{y}^\pm(y) = \frac{(\lambda_1 + \lambda_2 + \mu) \pm \sqrt{(\lambda_1 + \lambda_2 + \mu)^2 - 4\mu(\lambda_1 + \lambda_2 y)}}{2\mu}. \quad (110)$$

At  $y = 1$  the roots are  $\tilde{y}_+(1) = 1$  and  $\tilde{y}_-(1) = \rho$ . The root  $\tilde{y}_+ = 1$  would produce a non-summable geometric series as  $n \rightarrow \infty$ , so only the negative-sign root is admissible:

$$\tilde{y}^*(y) = \frac{(\lambda_1 + \lambda_2 + \mu) - \sqrt{(\lambda_1 + \lambda_2 + \mu)^2 - 4\mu(\lambda_1 + \lambda_2 y)}}{2\mu}. \quad (111)$$

The geometric ansatz gives  $\tilde{P}(y, n) = \tilde{P}(y, 0) \cdot [\tilde{y}^*(y)]^n$ . Using  $\tilde{P}(y, 0) = \tilde{\pi}(0, 0) = \pi(0, 0) = \rho(1 - \rho)$  (Corollary 1):

$$\tilde{P}(y, n) = \rho(1 - \rho) \cdot [\tilde{y}^*(y)]^n. \quad (112)$$

Since  $n$  counts customers *waiting* (the customer in service is not counted),  $n + 1$  is the total number in the system, so evaluating at  $y = 1$ :

$$\tilde{P}(1, n) = \rho(1 - \rho) \cdot \rho^n = (1 - \rho)\rho^{n+1}, \quad (113)$$

$\square$

which matches the  $M/M/1(\lambda, \mu)$  stationary distribution for  $n + 1$  customers in the system [AR02].

**Failure mechanism.** The neglected forcing  $\mu y^n(1-y)\tilde{\pi}(n+1, n+1)$  feeds in the diagonal states, in which all waiting customers are class 2; using  $\tilde{\pi}(n, n) = \pi(0, n)$  and the rate just proved in Lemma 3, this term decays in  $n$  like  $\rho^n$  (modulated by  $y^n$ ), whereas the homogeneous solution decays like  $[\tilde{y}^*(y)]^n$ . The approximation is therefore accurate exactly when the forcing rate is the faster of the two,  $\rho < \tilde{y}^*(y)$  on the relevant range of  $y$  — which holds when the priority class carries most of the load, i.e. when  $\rho_1/\rho$  is large and  $\rho$  moderate, and this is precisely the empirical validity region mapped in Figure 22. The obstruction is structural and recurs throughout this thesis: an unknown boundary/diagonal trace — here the sequence  $\pi(0, n)$  — couples into an otherwise solvable recursion, exactly as the unknown diagonal  $P(z, z)$  enters the Volterra forcing of Models  $B$  and  $X$  (§6, §7). A rigorous closure would retain the forcing and solve (105) by variation of constants in  $n$ , which reintroduces the very diagonal sequence  $\{\pi(0, n)\}$  as an unknown to be determined self-consistently; see §14.5.

### 5.3 Mean queue lengths and mean waiting times

#### 5.3.1 Class-1 marginal PGF and mean queue length

Setting  $y = 1$  in the fundamental equation (80) gives

$$[-\lambda_1(x-1)(x-1/\rho_1)]P(x, 1) = \mu(x-1)P_y(1). \quad (114)$$

Cancelling the common factor  $(x-1)$  for  $x \neq 1$  and rearranging:

$$P(x, 1) = \frac{\mu P_y(1)}{\mu - \lambda_1 x}. \quad (115)$$

The constant  $P_y(1)$  is pinned by the normalisation  $P(1, 1) = 1 - \pi_0 = \rho$ :

$$\rho = \frac{\mu P_y(1)}{\mu - \lambda_1} \implies P_y(1) = \rho(1 - \rho_1), \quad (116)$$

so that

$$P(x, 1) = \frac{\rho(1 - \rho_1)}{1 - \rho_1 x}. \quad (117)$$

Differentiating by the quotient rule and evaluating at  $x = 1$ :

$$\mathbb{E}[N_1] = \left. \frac{d}{dx} P(x, 1) \right|_{x=1} = \left. \frac{\rho(1 - \rho_1) \cdot \rho_1}{(1 - \rho_1 x)^2} \right|_{x=1} = \frac{\rho(1 - \rho_1) \rho_1}{(1 - \rho_1)^2} = \frac{\rho \rho_1}{1 - \rho_1}. \quad (118)$$

#### 5.3.2 Total queue length from the diagonal

Setting  $x = y = z$  in the kernel polynomial  $\lambda_1 x^2 - (\lambda_1 + \lambda_2 + \mu)x + \mu$  shows that  $x^*(z) = z$  iff  $z \in \{1, 1/\rho\}$ ; hence  $x^*(z) \neq z$  for all  $z \in (0, 1)$  with  $\rho < 1$ . Therefore the ratio  $(z - x^*(z))/(x^*(z) - z) = -1$  throughout  $(0, 1)$ , and the formula of Theorem 1 reduces to:

$$P(z, z) = \frac{z(1-z)\rho(1-\rho)}{D(z, z)} \cdot (-1). \quad (119)$$

The denominator at  $x = y = z$  is  $D(z, z) = (1 + \rho)z^2 - z - \rho z^3 = -\rho z(z-1)(z-1/\rho)$ , so:

$$\begin{aligned} P(z, z) &= \frac{-z(1-z)\rho(1-\rho)}{-\rho z(z-1)(z-1/\rho)} \\ &= \frac{(1-z)(1-\rho)}{(z-1)(z-1/\rho)} = \frac{-(1-\rho)}{z-1/\rho} = \frac{\rho(1-\rho)}{1-\rho z}, \end{aligned} \quad (120)$$

confirming the diagonal identity of Corollary 1. Since  $\left. \frac{d}{dz} P(z, z) \right|_{z=1} = \frac{\partial P}{\partial x}(1, 1) + \frac{\partial P}{\partial y}(1, 1) = \mathbb{E}[N_1] + \mathbb{E}[N_2]$ , differentiating by the quotient rule and evaluating at  $z = 1$  gives:

$$\mathbb{E}[N_1] + \mathbb{E}[N_2] = \left. \frac{d}{dz} \frac{\rho(1-\rho)}{1-\rho z} \right|_{z=1} = \left. \frac{\rho(1-\rho) \cdot \rho}{(1-\rho z)^2} \right|_{z=1} = \frac{\rho^2(1-\rho)}{(1-\rho)^2} = \frac{\rho^2}{1-\rho}. \quad (121)$$

This is the  $M/M/1(\lambda, \mu)$  queue-length formula for the combined system [AR02], as expected: the two classes together behave as a single  $M/M/1$  queue.

### 5.3.3 Class-2 mean queue length by subtraction

$$\begin{aligned}
\mathbb{E}[N_2] &= (\mathbb{E}[N_1] + \mathbb{E}[N_2]) - \mathbb{E}[N_1] = \frac{\rho^2}{1-\rho} - \frac{\rho\rho_1}{1-\rho_1} \\
&= \frac{\rho^2(1-\rho_1) - \rho\rho_1(1-\rho)}{(1-\rho)(1-\rho_1)} = \frac{\rho[\rho(1-\rho_1) - \rho_1(1-\rho)]}{(1-\rho)(1-\rho_1)} \\
&= \frac{\rho(\rho - \rho_1)}{(1-\rho)(1-\rho_1)} = \frac{\rho\rho_2}{(1-\rho_1)(1-\rho)}.
\end{aligned} \tag{122}$$

### 5.3.4 Mean waiting times via Little's Law

**Little's Law** applied to the *waiting sub-system* of each class: since  $N_i$  counts only customers *waiting* in queue (excluding the customer in service), Little's Law gives  $\mathbb{E}[N_i] = \lambda_i \mathbb{E}[W_i]$ , where  $\mathbb{E}[W_i]$  is the mean waiting time in queue (time from arrival until service start, not including the service duration):

$$\mathbb{E}[W_1] = \frac{\mathbb{E}[N_1]}{\lambda_1} = \frac{\rho\rho_1}{\lambda_1(1-\rho_1)} = \frac{\rho \cdot (\lambda_1/\mu)}{\lambda_1(1-\rho_1)} = \frac{\rho}{\mu(1-\rho_1)}, \tag{123}$$

$$\mathbb{E}[W_2] = \frac{\mathbb{E}[N_2]}{\lambda_2} = \frac{\rho\rho_2}{\lambda_2(1-\rho_1)(1-\rho)} = \frac{\rho \cdot (\lambda_2/\mu)}{\lambda_2(1-\rho_1)(1-\rho)} = \frac{\rho}{\mu(1-\rho_1)(1-\rho)}. \tag{124}$$

Class-1 benefits from priority:  $\mathbb{E}[W_2]$  carries the additional penalty factor  $1/(1-\rho)$  relative to  $\mathbb{E}[W_1]$ , reflecting the delay that class-2 customers incur while class-1 customers in the queue are served first. Note that both  $\mathbb{E}[W_1]$  and  $\mathbb{E}[W_2]$  depend on  $\rho_2$  through the total load  $\rho = \rho_1 + \rho_2$ ; class-2 affects class-1 waiting times only via server utilisation, not through a direct queueing penalty.

## 6 Analysis of Model-B ( $\gamma_1, \gamma_2 > 0, \theta_1 = \theta_2 = 0$ )

Model-B introduces bidirectional jockeying ( $\gamma_1, \gamma_2 > 0$ ) while retaining no abandonment ( $\theta_1 = \theta_2 = 0$ ). Jockeying conserves the total number of customers in the system, so the stationary distribution still exists under the same condition as Model-A. However, jockeying breaks the per-class Poisson/renewal structure: a class-2 customer can no longer identify its effective service time with a class-1 busy period, so the Pollaczek-Khinchine probabilistic route is unavailable. Applying the method of characteristics to the fundamental equation yields a general solution, but closing it — i.e., determining the unknown boundary function  $P_y(y)$  — requires solving a Volterra-type integral equation that does not admit a closed form. Model-B is therefore intentionally left open, in structural parallel with Model-X; its role is to motivate the tractable specialisation Model-B<sub>2</sub>.

**Stability.**  $\rho = (\lambda_1 + \lambda_2)/\mu < 1$ . Jockeying conserves the total customer count and does not affect this condition. Figure 5 shows the queue layout.




Figure 5: Queue layout for Model-B. Dashed arrows indicate bidirectional jockeying:  $\gamma_1$  moves a class-1 customer down to Queue 2 and  $\gamma_2$  moves a class-2 customer up to Queue 1. Both transfers conserve the total  $n$ .

**Corollary 2.** *The empty-system probability  $\pi_0$  (server idle, no one waiting) and the both-queues-empty probability  $\pi(0,0)$  (server busy,  $N_1 = N_2 = 0$ ) of Model-B coincide with the Model-A baseline,*

$$\pi_0 = 1 - \rho, \quad \pi(0,0) = \rho(1 - \rho), \tag{125}$$

and the diagonal of the PGF is  $P(z, z) = \pi(0, 0)/(1 - \rho z)$ .

*Proof.* Setting  $x = y$  in the fundamental equation (45) with  $\theta_1 = \theta_2 = 0$ , both jockeying terms carry the factor  $(x - y)$  and vanish, as does the boundary term  $\mu(x - y)P_y(y)$ ; dividing the surviving terms by the common factor  $x$  leaves

$$[(\lambda_1 + \lambda_2 + \mu)x - \mu - (\lambda_1 + \lambda_2)x^2]P(x, x) = -\mu(1 - x)\pi(0, 0). \quad (126)$$

Writing  $\lambda = \lambda_1 + \lambda_2$ , the bracket factors,

$$(\lambda + \mu)x - \mu - \lambda x^2 = -(\lambda x^2 - (\lambda + \mu)x + \mu) = -(\lambda x - \mu)(x - 1) = (\mu - \lambda x)(x - 1);$$

cancelling  $(x - 1)$  against  $-(1 - x) = (x - 1)$  on the right gives

$$P(x, x) = \frac{\mu \pi(0, 0)}{\mu - \lambda x} = \frac{\pi(0, 0)}{1 - \rho x}. \quad (127)$$

The idle state (0) exchanges probability flux only with the both-queues-empty busy state, so the global balance across the idle-busy cut reads  $\lambda\pi_0 = \mu\pi(0, 0)$  (an arrival at total rate  $\lambda$  leaves (0); a service completion at rate  $\mu$  returns to it), whence  $\pi(0, 0) = \rho\pi_0$ . Evaluating (127) at  $x = 1$  with  $P(1, 1) = 1 - \pi_0$  gives  $1 - \pi_0 = \rho\pi_0/(1 - \rho)$ , so  $(1 - \pi_0)(1 - \rho) = \rho\pi_0$ , i.e.  $\pi_0 = 1 - \rho$  and  $\pi(0, 0) = \rho(1 - \rho)$ . This matches Model-A (83): jockeying conserves the total-customer process and cannot change the marginal idle probability. The same identities hold in Models  $B_2$  and  $C_2$  for the same conservation reason (27).  $\square$

**Claim 1.** *Under stability,  $\rho < 1$ , the bivariate PGF of Model-B satisfies, on each characteristic curve  $\gamma_2 x + \gamma_1 y = C_1$  of the fundamental equation, the integral representation*

$$P(x, y(x)) = P_h(x, y(x)) \int_0^x \frac{h(t; C_1)}{P_h(t, y(t))} dt, \quad (128)$$

where  $P_h$  is the homogeneous solution (140),  $y(t) = (C_1 - \gamma_2 t)/\gamma_1$  is the characteristic parametrisation, and  $h(t; C_1)$  is an explicit forcing function depending on the unknown boundary trace  $P_y(y) = P(0, y)$ . Evaluating (128) along the diagonal  $x = y = z$  and using the conservation identity  $P(z, z) = \pi(0, 0)/(1 - \rho z)$  reduces the problem to a Volterra integral equation of the first kind for  $P_y$ :

$$\mathcal{F}(z) = \int_0^z \mathcal{K}(z, t) P_y(y(t; z)) dt, \quad (129)$$

where  $\mathcal{K}$  and  $\mathcal{F}$  are fully explicit (defined in (160)–(162) below). The kernel  $\mathcal{K}$  does not factorise for general  $(\gamma_1, \gamma_2)$ , so (129) does not reduce to a closed-form solution for  $P_y$ . Model-B is therefore left open.

*Proof of Claim 1.* The proof proceeds in five steps: (i) homogeneous solution via characteristics, (ii) rigorous sign analysis of the exponent  $E$ , (iii) particular solution via variation of parameters, (iv) pinning the free function from the boundary condition, and (v) diagonal evaluation yielding the Volterra equation.

#### Step 1: Homogeneous solution via the method of characteristics.

The fundamental equation (45) with  $\theta_1 = \theta_2 = 0$  is a first-order linear PDE in  $P(x, y)$ . Setting the right-hand side to zero and dividing through by  $y$  yields the homogeneous PDE

$$\gamma_1 x(x - y) \frac{\partial P}{\partial x} - \gamma_2 x(x - y) \frac{\partial P}{\partial y} = -[(\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy]P. \quad (130)$$

Following the method of characteristics (§3.6.3), we associate to this PDE the characteristic system

$$\frac{dx}{\gamma_1 x(x - y)} = \frac{dy}{-\gamma_2 x(x - y)} = \frac{dP}{-[(\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy]P}. \quad (131)$$

Taking the ratio of the first two differentials, the  $x(x - y)$  factors cancel, giving  $dx/\gamma_1 = dy/(-\gamma_2)$ . Integrating yields the first integral

$$C_1 = \gamma_2 x + \gamma_1 y \quad (\text{constant along each characteristic}). \quad (132)$$

The characteristics are therefore the straight lines of slope  $-\gamma_2/\gamma_1$  in the  $(x, y)$ -plane. Fix one such line and parametrise  $y$  by  $y(x) = (C_1 - \gamma_2 x)/\gamma_1$ . To extract the equation for  $P$  along this curve we need the coefficient  $A(x, y)$  and the denominator  $x(x - y)$  after substitution. Setting

$$A(x, y) := (\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy, \quad (133)$$

and substituting  $y = (C_1 - \gamma_2 x)/\gamma_1$  so that  $\lambda_2 xy = (\lambda_2 C_1/\gamma_1)x - (\lambda_2 \gamma_2/\gamma_1)x^2$ , we obtain, collecting by powers of  $x$ ,

$$\begin{aligned} A(x, y(x)) &= (\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \frac{\lambda_2 C_1}{\gamma_1}x + \frac{\lambda_2 \gamma_2}{\gamma_1}x^2 \\ &= -\frac{\gamma_1 \lambda_1 - \lambda_2 \gamma_2}{\gamma_1}x^2 + \frac{\gamma_1(\lambda_1 + \lambda_2 + \mu) - \lambda_2 C_1}{\gamma_1}x - \mu. \end{aligned} \quad (134)$$

Similarly,  $x - y(x) = [(\gamma_1 + \gamma_2)x - C_1]/\gamma_1$ , so  $\gamma_1 x(x - y(x)) = x[(\gamma_1 + \gamma_2)x - C_1]$ . The  $dP/P$  equation from the characteristic system therefore becomes

$$\frac{dP}{P} = \frac{-A(x, y(x))}{\gamma_1 x(x - y(x))} dx = \frac{(\gamma_1 \lambda_1 - \lambda_2 \gamma_2)x^2 - [\gamma_1(\lambda_1 + \lambda_2 + \mu) - \lambda_2 C_1]x + \gamma_1 \mu}{\gamma_1 x[(\gamma_1 + \gamma_2)x - C_1]} dx. \quad (135)$$

To integrate this, we decompose the integrand via partial fractions. Writing the integrand as  $(1/\gamma_1)I(x)dx$  with  $I(x) = K + M/x + N(C_1)/[(\gamma_1 + \gamma_2)x - C_1]$ , the partial fraction identity requires

$$K(\gamma_1 + \gamma_2)x^2 + [M(\gamma_1 + \gamma_2) + N - KC_1]x - MC_1 = (\gamma_1 \lambda_1 - \lambda_2 \gamma_2)x^2 - [\gamma_1(\lambda_1 + \lambda_2 + \mu) - \lambda_2 C_1]x + \gamma_1 \mu.$$

Matching the  $x^2$ ,  $x^0$ , and  $x^1$  coefficients in turn:

$$K = \frac{\gamma_1 \lambda_1 - \lambda_2 \gamma_2}{\gamma_1 + \gamma_2}, \quad M = -\frac{\gamma_1 \mu}{C_1}, \quad N(C_1) = \gamma_1 \left[ \frac{\lambda_1 + \lambda_2}{\gamma_1 + \gamma_2} C_1 - (\lambda_1 + \lambda_2 + \mu) + \frac{\mu(\gamma_1 + \gamma_2)}{C_1} \right]. \quad (136)$$

For  $N$ , the  $x^1$  match gives  $N = -\gamma_1(\lambda_1 + \lambda_2 + \mu) + \lambda_2 C_1 + KC_1 - M(\gamma_1 + \gamma_2)$ ; substituting  $K$  and  $M$ , writing  $\lambda = \lambda_1 + \lambda_2$ ,  $\Gamma = \gamma_1 + \gamma_2$ , and collecting the two  $C_1$ -proportional terms over the common denominator  $\Gamma$ ,

$$\begin{aligned} N &= -\gamma_1(\lambda + \mu) + C_1 \frac{\lambda_2 \Gamma + \gamma_1 \lambda_1 - \lambda_2 \gamma_2}{\Gamma} + \frac{\gamma_1 \mu \Gamma}{C_1} \\ &= -\gamma_1(\lambda + \mu) + \frac{\gamma_1 \lambda}{\Gamma} C_1 + \frac{\gamma_1 \mu \Gamma}{C_1}, \end{aligned}$$

using  $\lambda_2 \Gamma + \gamma_1 \lambda_1 - \lambda_2 \gamma_2 = \lambda_2 \gamma_1 + \gamma_1 \lambda_1 = \gamma_1 \lambda$  in the middle term; factoring out  $\gamma_1$  yields (136). Integrating  $dP/P = (1/\gamma_1)I(x)dx$  term by term gives

$$\ln P = \frac{K}{\gamma_1}x - \frac{\mu}{C_1} \ln x + \frac{N(C_1)}{\gamma_1(\gamma_1 + \gamma_2)} \ln[(\gamma_1 + \gamma_2)x - C_1] + \ln f(C_1), \quad (137)$$

where  $f(C_1)$  is an arbitrary function of the characteristic constant. Now observe that

$$(\gamma_1 + \gamma_2)x - C_1 = (\gamma_1 + \gamma_2)x - (\gamma_2 x + \gamma_1 y) = \gamma_1(x - y), \quad (138)$$

so  $\ln[(\gamma_1 + \gamma_2)x - C_1] = \ln \gamma_1 + \ln(x - y)$ . Exponentiating (137) term by term,

$$P = f(C_1) \exp\left(\frac{K}{\gamma_1}x\right) \cdot x^{-\mu/C_1} \cdot \gamma_1^{N(C_1)/(\gamma_1 \Gamma)} (x - y)^{N(C_1)/(\gamma_1 \Gamma)};$$

the factor  $\gamma_1^{N(C_1)/(\gamma_1 \Gamma)}$  depends on  $(x, y)$  only through  $C_1$  and is absorbed, together with  $f$ , into a redefined arbitrary function  $F(C_1)$ . Reading off the exponent of  $(x - y)$ :

$$E(x, y) := \frac{N(C_1)}{\gamma_1(\gamma_1 + \gamma_2)} \Big|_{C_1 = \gamma_2 x + \gamma_1 y} = \frac{\lambda_1 + \lambda_2}{(\gamma_1 + \gamma_2)^2} (\gamma_2 x + \gamma_1 y) - \frac{\lambda_1 + \lambda_2 + \mu}{\gamma_1 + \gamma_2} + \frac{\mu}{\gamma_2 x + \gamma_1 y}. \quad (139)$$

Exponentiating (137) yields the general homogeneous solution

$$P_h(x, y) = F(\gamma_2 x + \gamma_1 y) \cdot \exp\left(\frac{\gamma_1 \lambda_1 - \lambda_2 \gamma_2}{\gamma_1(\gamma_1 + \gamma_2)}x\right) \cdot x^{-\mu/(\gamma_2 x + \gamma_1 y)} \cdot (x - y)^{E(x, y)}, \quad (140)$$

where  $F$  is an arbitrary differentiable function to be determined.

**Step 2: The exponent  $E(x, y)$  is strictly positive on  $(0, 1)^2$ .**

Recall  $\lambda = \lambda_1 + \lambda_2$  and  $\Gamma = \gamma_1 + \gamma_2$ . Along any fixed characteristic,  $C_1 = \gamma_2 x + \gamma_1 y$  is constant, so the exponent (139) depends only on  $u := C_1 \in (0, \Gamma)$ :

$$E(u) = \frac{\lambda}{\Gamma^2} u - \frac{\lambda + \mu}{\Gamma} + \frac{\mu}{u}. \quad (141)$$

(For  $(x, y) \in (0, 1)^2$  we have  $u = \gamma_2 x + \gamma_1 y \in (0, \Gamma)$ ;  $E(u)$  is continuous there, with  $E(u) \rightarrow +\infty$  as  $u \rightarrow 0^+$  and  $E(u) \rightarrow 0^+$  as  $u \rightarrow \Gamma^-$ , so it is finite at every interior point although not uniformly bounded.) Multiplying through by  $u\Gamma^2 > 0$  gives the equivalent quadratic

$$q(u) := \lambda u^2 - (\lambda + \mu)\Gamma u + \mu\Gamma^2. \quad (142)$$

The sign of  $E$  on  $(0, \Gamma)$  equals the sign of  $q(u)$  there. We factor  $q$  by finding its roots. The discriminant is  $\Delta = (\lambda + \mu)^2\Gamma^2 - 4\lambda\mu\Gamma^2 = \Gamma^2(\mu - \lambda)^2$ , so (using  $\rho < 1 \Rightarrow \mu > \lambda \Rightarrow \mu - \lambda > 0$ )

$$u_{1,2} = \frac{(\lambda + \mu)\Gamma \pm \Gamma(\mu - \lambda)}{2\lambda} \implies u_1 = \frac{2\lambda\Gamma}{2\lambda} = \Gamma, \quad u_2 = \frac{2\mu\Gamma}{2\lambda} = \frac{\Gamma}{\rho} > \Gamma. \quad (143)$$

Hence

$$q(u) = \lambda(u - \Gamma)\left(u - \frac{\Gamma}{\rho}\right). \quad (144)$$

For every  $u \in (0, \Gamma)$  both factors are strictly negative ( $(u - \Gamma) < 0$  and  $(u - \Gamma/\rho) < 0$  since  $\Gamma/\rho > \Gamma$ ), so  $q(u) = \lambda \cdot (\text{neg}) \cdot (\text{neg}) > 0$ , and therefore

$$E(x, y) > 0 \quad \text{for all } (x, y) \in (0, 1)^2. \quad (145)$$

Note that  $E \rightarrow 0$  as  $u \rightarrow \Gamma^-$  (i.e. as  $(x, y) \rightarrow (1, 1)$ ), while  $E$  is bounded and bounded away from zero in any compact subset of  $(0, 1)^2$ .

*Sign convention and diagonal limit.* Because  $E > 0$ , the factor  $(x - y)^E$  in  $P_h$  carries the complex phase  $e^{i\pi E}$  whenever  $x < y$ . This phase is absorbed into the arbitrary function  $F(C_1)$ , and the ratio  $P_h(x, y(x))/P_h(t, y(t))$  that appears in the integral representation is always real and positive: both numerator and denominator lie on the same characteristic (same  $C_1$ , same constant value of  $E$ ), so the phases cancel exactly. Crucially, since  $E > 0$  the factor  $(y(t; z) - t)^E \rightarrow 0$  as  $t \rightarrow z^-$ , meaning  $P_h$  formally vanishes on the diagonal  $x = y$ . The diagonal equation that appears in Step 5 below is therefore understood as the  $\lim_{x \rightarrow z^-}$  of (151); the limit-by-limit analysis carried out there, using the exact linear relation  $y(t; z) - t = (\Gamma/\gamma_1)(z - t)$ , confirms that the  $0 \cdot \infty$  limit is finite and equals  $\pi(0, 0)/(1 - \rho z)$ .

**Step 3: Particular solution via variation of parameters.**

We now work with the full (inhomogeneous) fundamental equation restricted to a fixed characteristic. Along the curve  $y(x) = (C_1 - \gamma_2 x)/\gamma_1$ , the total derivative satisfies  $\frac{d}{dx}P(x, y(x)) = \partial_x P + \frac{dy}{dx}\partial_y P = \partial_x P - \frac{\gamma_2}{\gamma_1}\partial_y P$ , so the jockeying terms in (45) become

$$\gamma_1 xy(x - y)\partial_x P + \gamma_2 xy(y - x)\partial_y P = \gamma_1 xy(x - y)\left(\partial_x P - \frac{\gamma_2}{\gamma_1}\partial_y P\right) = \gamma_1 xy(x - y)\frac{d}{dx}P(x, y(x)).$$

The fundamental equation along the characteristic therefore reads, with its algebraic coefficient factorised as  $yA(x, y)$  (cf. (133)),

$$yA(x, y)P + \gamma_1 xy(x - y)\frac{dP}{dx} = G(x, y),$$

and dividing through by  $\gamma_1 xy(x - y(x))$  reduces the PDE to the first-order linear ODE (for fixed  $C_1$ )

$$\frac{dP}{dx} = f(x; C_1)P + h(x; C_1), \quad (146)$$

where the coefficient  $f$  and the forcing term  $h$  are

$$f(x; C_1) = -\frac{A(x, y(x))}{\gamma_1 x(x - y(x))}, \quad (147)$$

$$h(x; C_1) = \frac{G(x, y(x))}{\gamma_1 xy(x)(x - y(x))}, \quad (148)$$

with  $A$  as in (133) and

$$G(x, y) := \mu(x - y) P_y(y) - \mu x(1 - y) \pi(0, 0). \quad (149)$$

Since  $P_h(x, y(x))$  satisfies the homogeneous equation  $\frac{d}{dx} P_h = f(x; C_1) P_h$ , it is the fundamental solution of ODE (146). Applying variation of parameters (§3.6.2), the general solution of the inhomogeneous ODE is

$$P(x, y(x)) = P_h(x, y(x)) \left[ F(C_1) + \int_{x_0}^x \frac{h(t; C_1)}{P_h(t, y(t))} dt \right]. \quad (150)$$

**Step 4: Pinning the free function from the boundary condition at  $x = 0$ .**

Along any characteristic with constant  $C_1 > 0$ , the endpoint at  $x = 0$  is  $y(0) = C_1/\gamma_1$ , giving

$$P\left(0, \frac{C_1}{\gamma_1}\right) = P_y\left(\frac{C_1}{\gamma_1}\right),$$

which is finite. However,  $P_h$  contains the factor  $x^{-\mu/C_1} \rightarrow +\infty$  as  $x \rightarrow 0^+$  (since  $\mu/C_1 > 0$  for  $C_1 > 0$ ), while the integral in (150) vanishes as  $x_0 \rightarrow 0$  (the integration interval collapses). Therefore

$$\lim_{x \rightarrow 0^+} P(x, y(x)) = \lim_{x \rightarrow 0^+} P_h(x, y(x)) \cdot F(C_1) = +\infty \cdot F(C_1),$$

which can only be finite if  $F(C_1) = 0$ . Setting  $x_0 = 0$  and  $F(C_1) = 0$  in (150) gives the integral representation of Claim 1:

$$P(x, y(x)) = P_h(x, y(x)) \int_0^x \frac{h(t; C_1)}{P_h(t, y(t))} dt. \quad (151)$$

**Step 5: Reduction to the Volterra equation.**

By Corollary 2 the empty-state probabilities are  $\pi_0 = 1 - \rho$  and  $\pi(0, 0) = \rho(1 - \rho)$ , with diagonal value  $P(z, z) = \pi(0, 0)/(1 - \rho z)$  by (127); these are the inputs to the diagonal evaluation below.

We simplify the forcing function  $h$  on the characteristic. Substituting  $G(t, y(t))$  from (149) into (148):

$$\begin{aligned} h(t; C_1) &= \frac{\mu(t - y(t)) P_y(y(t)) - \mu t(1 - y(t)) \pi(0, 0)}{\gamma_1 t y(t) (t - y(t))} \\ &= \frac{\mu P_y(y(t))}{\gamma_1 t y(t)} - \frac{\mu(1 - y(t)) \pi(0, 0)}{\gamma_1 y(t) (t - y(t))}. \end{aligned} \quad (152)$$

We will evaluate (151) along the diagonal characteristic, which passes through  $(z, z)$  and has  $C_1 = (\gamma_1 + \gamma_2)z$ . The parametrisation along that curve is  $y(t; z) = ((\gamma_1 + \gamma_2)z - \gamma_2 t)/\gamma_1$ , so  $y(0; z) = (\gamma_1 + \gamma_2)z/\gamma_1 > z$  and  $y(z; z) = z$ . Therefore  $y(t; z) > t$  for all  $t \in [0, z]$ , which means  $t - y(t; z) < 0$  throughout the integration interval. Writing  $t - y(t; z) = -(y(t; z) - t)$  in the denominator of (152):

$$h(t; (\gamma_1 + \gamma_2)z) = \frac{\mu}{\gamma_1 t y(t; z)} P_y(y(t; z)) + \frac{\mu(1 - y(t; z))}{\gamma_1 y(t; z) (y(t; z) - t)} \pi(0, 0). \quad (153)$$

As established in Step 2,  $E > 0$ , so the homogeneous factor  $(x - y)^E$  vanishes on the diagonal and  $P_h(x, y(x; z)) \rightarrow 0$  as  $x \rightarrow z^-$ ; simultaneously the integrand in (151) develops a non-integrable singularity  $\sim (y(t; z) - t)^{-1-E}$  as  $t \rightarrow z^-$ , so the integral diverges. Their product is the finite  $0 \cdot \infty$  limit

$$\frac{\pi(0, 0)}{1 - \rho z} = \lim_{x \rightarrow z^-} P_h(x, y(x; z)) \int_0^x \frac{h(t; (\gamma_1 + \gamma_2)z)}{P_h(t, y(t; z))} dt, \quad (154)$$

whose value is  $P(z, z) = \pi(0, 0)/(1 - \rho z)$  by Corollary 2. Substituting (153) and separating the term containing the unknown  $P_y$  from the known remainder, this limit splits as

$$\begin{aligned} \frac{\pi(0, 0)}{1 - \rho z} &= \lim_{x \rightarrow z^-} \frac{\mu P_h(x, y(x; z))}{\gamma_1} \int_0^x \frac{P_y(y(t; z))}{t \cdot y(t; z) \cdot P_h(t, y(t; z))} dt \\ &\quad + \lim_{x \rightarrow z^-} \frac{\mu P_h(x, y(x; z)) \pi(0, 0)}{\gamma_1} \int_0^x \frac{1 - y(t; z)}{y(t; z) \cdot (y(t; z) - t) \cdot P_h(t, y(t; z))} dt. \end{aligned} \quad (155)$$



We now evaluate the two limits in (155) one by one; this both clarifies in what sense the kernel pairing below is to be read and provides an independent consistency check of the representation. The geometry on the diagonal characteristic is exactly linear,

$$y(t; z) - t = \frac{(\gamma_1 + \gamma_2)z - \gamma_2 t}{\gamma_1} - t = \frac{\Gamma}{\gamma_1} (z - t), \quad (156)$$

and along this characteristic ( $C_1 = \Gamma z$ ) the exponent of Step 2 takes the closed form

$$E_z := E(\Gamma z) = \frac{\lambda z^2 - (\lambda + \mu)z + \mu}{\Gamma z} = \frac{(\mu - \lambda z)(1 - z)}{\Gamma z} > 0, \quad z \in (0, 1). \quad (157)$$

Every factor of  $P_h$  in (140) other than the exponential, the power of  $x$ , and  $(x - y)^E$  depends on  $(x, y)$  only through  $C_1$ , so along the characteristic the homogeneous ratio is *exact*:

$$\frac{P_h(x, y(x; z))}{P_h(t, y(t; z))} = R(x, t) \left( \frac{z - x}{z - t} \right)^{E_z}, \quad R(x, t) := e^{\frac{\kappa}{\gamma_1}(x-t)} \left( \frac{t}{x} \right)^{\mu/(\Gamma z)}, \quad (158)$$

with  $R$  smooth and  $R(z, z) = 1$ . Substituting (158) into the *second* limit of (155) and using (156), its integrand behaves as  $w(z)(z - t)^{-1-E_z}$  near  $t = z$ , with  $w(z) = \mu \pi(0, 0)(1 - z)/(\Gamma z)$ ; the truncated integral diverges like  $(z - x)^{-E_z}/E_z$ , the prefactor vanishes like  $(z - x)^{E_z}$ , and the product converges:

$$\begin{aligned} \lim_{x \rightarrow z^-} (z - x)^{E_z} \int^x w(z) (z - t)^{-1-E_z} dt &= \frac{w(z)}{E_z} = \frac{\mu \pi(0, 0)(1 - z)}{\Gamma z} \cdot \frac{\Gamma z}{(\mu - \lambda z)(1 - z)} \\ &= \frac{\mu \pi(0, 0)}{\mu - \lambda z} = \frac{\pi(0, 0)}{1 - \rho z}. \end{aligned} \quad (159)$$

The second limit alone thus reproduces the full diagonal value — an independent confirmation that the representation (151) is consistent with Corollary 2. The *first* limit, by contrast, vanishes: its integrand behaves as  $(z - t)^{-E_z}$  near  $t = z$ , so for  $E_z < 1$  the integral converges and is annihilated by the vanishing prefactor  $(z - x)^{E_z}$ , while for  $E_z \geq 1$  the truncated integral grows only like  $(z - x)^{1-E_z}$  (logarithmically at  $E_z = 1$ ) and the product is still  $O((z - x)^{\min(E_z, 1)} \ln(z - x)) \rightarrow 0$ . Hence *at leading order the limit (155) reduces to the identity*  $\pi(0, 0)/(1 - \rho z) = \pi(0, 0)/(1 - \rho z)$  *and carries no information about*  $P_y$ : the dependence on the boundary function enters only at the next order in  $(z - x)$ . Each limit is accordingly a *Hadamard finite part*: writing  $P_h^{\text{diag}}(z) := \lim_{x \rightarrow z^-} P_h(x, y(x; z))$  (which vanishes pointwise), the kernel pairing below is the extraction of the truncation-independent, subleading part of the divergent integrals, and neither factor is meaningful in isolation. Writing the kernel and known term *formally* as

$$\mathcal{K}(z, t) := \frac{\mu \cdot P_h^{\text{diag}}(z)}{\gamma_1 \cdot t \cdot y(t; z) \cdot P_h(t, y(t; z))}, \quad (160)$$

$$\mathcal{B}(z) := \frac{\mu \cdot P_h^{\text{diag}}(z) \cdot \pi(0, 0)}{\gamma_1} \int_0^z \frac{1 - y(t; z)}{y(t; z) \cdot (y(t; z) - t) \cdot P_h(t, y(t; z))} dt, \quad (161)$$

*to be read through the regularised limit (155) rather than pointwise*, and setting  $\mathcal{F}(z) = \pi(0, 0)/(1 - \rho z) - \mathcal{B}(z)$ , equation (155) becomes the Volterra integral equation of the first kind

$$\mathcal{F}(z) = \int_0^z \mathcal{K}(z, t) P_y(y(t; z)) dt, \quad (162)$$

confirming Claim 1. The finite-part reading is not a decoration. As the limit-by-limit analysis shows, the displayed  $\mathcal{K}$  and  $\mathcal{B}$  are formal — the prefactor  $P_h^{\text{diag}}(z)$  vanishes pointwise — and the equation for  $P_y$  lives entirely at subleading order in  $(z - x)$ . Giving that extraction a rigorous meaning, whether by matched asymptotics near  $t = z$  or by a change of variable removing the  $(z - t)^{-1-E_z}$  singularity before passing to the limit, is itself the analytical obstruction: it is this step, together with the fact that the kernel  $\mathcal{K}$  depends on  $P_h$  through both arguments and does not simplify for general  $(\gamma_1, \gamma_2)$ , that places Model-B beyond the scope of this thesis. Equation (162) is recorded as the precise form of the open problem, not solved here; resolving it would require Volterra inversion methods together with the regularisation just described.  $\square$



Model- $B$  is therefore left open at this point. Its role in the hierarchy is structural: it identifies the precise obstruction — the Volterra equation for  $P_y$  — that the tractable special cases  $B_2$  and  $C_2$  circumvent, each by a different mechanism. Setting  $\gamma_2 = 0$  collapses the characteristic family from a two-parameter set of oblique lines to a one-parameter set of vertical lines, degenerating the PDE to an ODE in  $x$  alone; the boundary function  $P_y(y)$  is then pinned by an analyticity argument at the interior singular point  $x = y$ . Model- $C_2$  removes jockeying entirely and pins  $P_y$  by a boundedness argument at the boundary of the domain. Both specialisations are analysed in the following sections.

## 7 Analysis of Model- $X$ ( $\gamma_1, \gamma_2 > 0$ , $\theta_1, \theta_2 > 0$ )

This section analyses the full model, Model- $X$ , which retains *both* bidirectional jockeying ( $\gamma_1, \gamma_2 > 0$ ) *and* two-class abandonment ( $\theta_1, \theta_2 > 0$ ), and from which every other model in this work is recovered by zeroing parameters. No probabilistic route is available: jockeying breaks the per-class Poisson/renewal structure—a class-2 customer cannot identify its effective service time with a class-1 busy period—so the Pollaczek–Khinchine relation does not apply, while abandonment, though it leaves each arrival stream Poisson, couples the two queues through the state-dependent rates  $\theta_i n_i$ . The analysis therefore proceeds analytically. The method of characteristics yields the homogeneous solution of the fundamental equation; variation of parameters produces the general solution; the boundary behaviour at  $x = 0$  pins the free function; and evaluating the result along the characteristic through a diagonal point reduces the determination of the unknown boundary function  $P_y(y)$  to a Volterra integral equation of the first kind. There the analysis stops, for two reasons established below: the kernel of that equation is built from a non-elementary quadrature along transcendental characteristics, and its right-hand side is itself coupled to an unknown diagonal gradient. Model- $X$  is therefore left open; its role in the hierarchy is to expose the precise obstructions that its tractable specialisations circumvent, and its queue layout is the left panel of Figure 1.

**Stability.** Model- $X$  has abandonment from both classes, so the total departure rate in state  $(n_1, n_2)$  is  $\mu + \theta_1 n_1 + \theta_2 n_2$ , which grows without bound in every direction of the  $(n_1, n_2)$ -plane while the total arrival rate stays fixed at  $\lambda = \lambda_1 + \lambda_2$ : far from the origin the chain drifts strictly towards smaller queues, so for any  $\theta_1, \theta_2 > 0$  no traffic-intensity restriction of the form  $\rho < 1$  is to be expected (cf. the Erlang-A queue, §3.3, which is stable at all loads). Throughout this section we therefore work under the hypothesis that the chain is positive recurrent, so that the stationary distribution exists. Setting  $\theta_1 = \theta_2 = 0$  removes this unbounded downward drift, and the condition reverts to  $\rho = (\lambda_1 + \lambda_2)/\mu < 1$  as in the non-abandoning models (§§5–6).

**Corollary 3.** *The empty-system probability  $\pi_0$  (server idle, no one waiting) and the both-queues-empty busy probability  $\pi(0, 0)$  (server busy,  $N_1 = N_2 = 0$ ) of Model- $X$  satisfy the universal idle balance*

$$\pi(0, 0) = \rho \pi_0, \quad \rho = \frac{\lambda_1 + \lambda_2}{\mu}, \quad (163)$$

*as in every model of this work (26). With abandonment active, however, the total-customer process is a state-dependent birth–death chain rather than an  $M/M/1$ , so  $\pi_0 \neq 1 - \rho$ : the idle probability depends on  $\theta_1, \theta_2$  and is fixed only by the global normalisation  $P(1, 1) = 1 - \pi_0$  once  $P$  is known. Correspondingly, the diagonal of the PGF no longer collapses to  $\pi(0, 0)/(1 - \rho z)$ ; setting  $x = y = z$  in the fundamental equation leaves the abandonment-corrected relation*

$$(\mu - \lambda z) P(z, z) + z[\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)} = \mu \pi(0, 0), \quad \lambda = \lambda_1 + \lambda_2. \quad (164)$$

*As  $\theta_1, \theta_2 \rightarrow 0^+$  the gradient terms drop out and (164) returns the identities  $P(z, z) = \pi(0, 0)/(1 - \rho z)$ ,  $\pi_0 = 1 - \rho$  and  $\pi(0, 0) = \rho(1 - \rho)$  of the non-abandoning models (125), (83).*

*Proof.* Set  $x = y = z$  in the fundamental equation (45) and track its four groups of terms separately. The two jockeying contributions carry the factor  $(x - y)$  and vanish, as does the boundary term  $\mu(x - y)P_y(y)$ ; the abandonment contributions, carrying instead the factors  $(x - 1)$  and  $(y - 1)$ , survive:

$$\begin{aligned} xy[\gamma_1(x - y) + \theta_1(x - 1)] \partial_x P \Big|_{x=y=z} &= z^2 \theta_1(z - 1) \partial_x P(z, z), \\ xy[\gamma_2(y - x) + \theta_2(y - 1)] \partial_y P \Big|_{x=y=z} &= z^2 \theta_2(z - 1) \partial_y P(z, z). \end{aligned}$$

The algebraic coefficient of  $P$  factorises, with  $\lambda = \lambda_1 + \lambda_2$ ,

$$\begin{aligned} (\lambda_1 + \lambda_2 + \mu)z^2 - \mu z - \lambda_1 z^3 - \lambda_2 z^3 &= z[-\lambda z^2 + (\lambda + \mu)z - \mu] \\ &= -z(\lambda z - \mu)(z - 1) = z(z - 1)(\mu - \lambda z), \end{aligned}$$

and the surviving right-hand side is  $-\mu z(1 - z)\pi(0, 0) = \mu z(z - 1)\pi(0, 0)$ . Collecting,

$$z(z - 1)(\mu - \lambda z)P(z, z) + z^2(z - 1)[\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)} = \mu z(z - 1)\pi(0, 0), \quad (165)$$

and dividing by the common factor  $z(z - 1) \neq 0$  (for  $z \in (0, 1)$ ) gives (164). The idle state (0) exchanges probability flux only with the both-queues-empty busy state, so global balance across the idle-busy cut reads  $\lambda\pi_0 = \mu\pi(0, 0)$  (32) (an arrival at total rate  $\lambda$  leaves (0); a service completion at rate  $\mu$  returns to it), whence  $\pi(0, 0) = \rho\pi_0$ , which is (163). When  $\theta_1 = \theta_2 = 0$  the gradient terms in (164) are absent, so  $P(z, z) = \pi(0, 0)/(1 - \rho z)$ ; evaluating at  $z = 1$  with  $P(1, 1) = 1 - \pi_0$  then forces  $(1 - \pi_0)(1 - \rho) = \rho\pi_0$ , i.e.  $\pi_0 = 1 - \rho$ . With abandonment those gradient terms are present and *not* known, so the diagonal no longer determines  $P(z, z)$ , and  $\pi_0$  cannot be read off in closed form: it follows only from the full solution via normalisation. The same gradient coupling resurfaces in Step 5 of the analysis below, and again, in scalar form, for Model- $C_2$  (§9).  $\square$

**Claim 2.** *Under positive recurrence, the bivariate PGF of Model- $X$  satisfies, on each characteristic curve  $\Phi(x, y) = C_1$  of the fundamental equation, the integral representation*

$$P(x, y(x)) = P_h(x, y(x)) \int_0^x \frac{h(t; C_1)}{P_h(t, y(t))} dt, \quad (166)$$

where  $P_h$  is the homogeneous solution (179),  $y(t) = y(t; C_1)$  is the characteristic parametrisation,  $\Phi$  is the first integral (177) (a transcendental function of  $(x, y)$ ), and  $h(t; C_1)$  is a forcing function depending on the unknown boundary trace  $P_y(y) = P(0, y)$ . Evaluating (166) along the characteristic through a diagonal point  $(z, z)$  reduces the problem to a Volterra integral equation of the first kind for  $P_y$ :

$$\mathcal{F}(z) = \int_0^z \mathcal{K}(z, t) P_y(y(t; z)) dt, \quad (167)$$

whose kernel  $\mathcal{K}$  and right-hand side  $\mathcal{F}$  are defined in (190)–(193) below. The kernel  $\mathcal{K}$  is built from the non-elementary homogeneous solution  $P_h$  and does not simplify for general  $(\gamma_1, \gamma_2, \theta_1, \theta_2)$ ; moreover, the right-hand side  $\mathcal{F}$  is not fully known — it is coupled through (164) to the diagonal gradient  $[\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)}$ . Equation (167) therefore does not reduce to a closed-form solution for  $P_y$ , and Model- $X$  is left open.

*Proof of Claim 2.* The proof proceeds in five steps: (i) homogeneous solution via the method of characteristics, (ii) behaviour of  $P_h$  at the boundaries, (iii) particular solution via variation of parameters, (iv) pinning the free function from the boundary condition, and (v) diagonal evaluation yielding the Volterra equation.

### Step 1: Homogeneous solution via the method of characteristics.

The fundamental equation (45) is a first-order linear PDE in  $P(x, y)$ . Every one of its coefficients contains the overall factor  $y$ : the algebraic coefficient factorises as

$$(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 xy^2 = y A(x, y), \quad A(x, y) := (\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy, \quad (168)$$

and both convection coefficients carry the prefactor  $xy$ . Setting the right-hand side to zero and dividing through by  $y$  therefore yields the homogeneous PDE

$$x[\gamma_1(x - y) + \theta_1(x - 1)] \frac{\partial P}{\partial x} + x[\gamma_2(y - x) + \theta_2(y - 1)] \frac{\partial P}{\partial y} = -A(x, y) P. \quad (169)$$

Following the method of characteristics (§3.6.3), we associate to (169) the characteristic system

$$\frac{dx}{x[\gamma_1(x - y) + \theta_1(x - 1)]} = \frac{dy}{x[\gamma_2(y - x) + \theta_2(y - 1)]} = \frac{dP}{-A(x, y) P}. \quad (170)$$

Taking the ratio of the first two differentials, the common factor  $x$  cancels:

$$\frac{dy}{dx} = \frac{\gamma_2(y - x) + \theta_2(y - 1)}{\gamma_1(x - y) + \theta_1(x - 1)} = \frac{-\gamma_2 x + (\gamma_2 + \theta_2)y - \theta_2}{(\gamma_1 + \theta_1)x - \gamma_1 y - \theta_1}. \quad (171)$$

When  $\theta_1 = \theta_2 = 0$  the two affine forms in (171) share the factor  $(x - y)$ , the slope collapses to the constant  $-\gamma_2/\gamma_1$ , and the characteristics are the straight lines  $\gamma_2 x + \gamma_1 y = \text{const}$  (the jockeying-only case of §6). With abandonment active no common factor survives: (171) is a genuine first-order ODE whose right-hand side is a ratio of two affine forms, and its integral curves are not straight lines.

Both affine forms in (171) vanish simultaneously only at the single point  $(x, y) = (1, 1)$  (the unique intersection of the lines  $(\gamma_1 + \theta_1)x - \gamma_1 y - \theta_1 = 0$  and  $-\gamma_2 x + (\gamma_2 + \theta_2)y - \theta_2 = 0$ ), which is therefore the lone singular point of the characteristic field. Shifting to it via  $X = x - 1$ ,  $Y = y - 1$  removes the constant terms and renders (171) homogeneous:

$$\frac{dY}{dX} = \frac{-\gamma_2 X + (\gamma_2 + \theta_2)Y}{(\gamma_1 + \theta_1)X - \gamma_1 Y}. \quad (172)$$

The right-hand side of (172) depends on  $(X, Y)$  only through the ratio  $v := Y/X$ , so we substitute  $Y = vX$ ,  $\frac{dY}{dX} = v + X \frac{dv}{dX}$ , and isolate  $X dv/dX$  over the common denominator:

$$\begin{aligned} v + X \frac{dv}{dX} &= \frac{-\gamma_2 + (\gamma_2 + \theta_2)v}{(\gamma_1 + \theta_1) - \gamma_1 v} \\ \implies X \frac{dv}{dX} &= \frac{[-\gamma_2 + (\gamma_2 + \theta_2)v] - v[(\gamma_1 + \theta_1) - \gamma_1 v]}{(\gamma_1 + \theta_1) - \gamma_1 v} = \frac{\gamma_1 v^2 + (\gamma_2 + \theta_2 - \gamma_1 - \theta_1)v - \gamma_2}{(\gamma_1 + \theta_1) - \gamma_1 v}. \end{aligned}$$

Naming the quadratic in the numerator  $Q(v)$ , the variables separate:

$$\frac{(\gamma_1 + \theta_1) - \gamma_1 v}{Q(v)} dv = \frac{dX}{X}, \quad Q(v) := \gamma_1 v^2 + (\gamma_2 + \theta_2 - \gamma_1 - \theta_1)v - \gamma_2. \quad (173)$$

The quadratic  $Q$  has discriminant  $D = (\gamma_2 + \theta_2 - \gamma_1 - \theta_1)^2 + 4\gamma_1\gamma_2 > 0$  and real roots of opposite sign (their product is  $-\gamma_2/\gamma_1 < 0$ ),

$$v_{\pm} = \frac{-(\gamma_2 + \theta_2 - \gamma_1 - \theta_1) \pm \sqrt{D}}{2\gamma_1}, \quad v_- < 0 < v_+. \quad (174)$$

Decompose the left-hand side of (173) by partial fractions over  $Q(v) = \gamma_1(v - v_+)(v - v_-)$ : writing

$$\frac{(\gamma_1 + \theta_1) - \gamma_1 v}{\gamma_1(v - v_+)(v - v_-)} = \frac{a_+}{v - v_+} + \frac{a_-}{v - v_-} \quad (175)$$

and clearing denominators,

$$(\gamma_1 + \theta_1) - \gamma_1 v = \gamma_1 [a_+(v - v_-) + a_-(v - v_+)].$$

Evaluating at  $v = v_+$  and  $v = v_-$  in turn isolates each coefficient, while comparing the coefficients of  $v$  on the two sides gives  $-\gamma_1 = \gamma_1(a_+ + a_-)$ :

$$a_{\pm} = \frac{(\gamma_1 + \theta_1) - \gamma_1 v_{\pm}}{\gamma_1(v_{\pm} - v_{\mp})}, \quad a_+ + a_- = -1 \quad (176)$$

(equivalently,  $a_+ + a_-$  is read off from the  $1/v$  decay of the integrand at  $v = \infty$ ). Integrating (173) term by term and exponentiating,

$$\begin{aligned} a_+ \ln(v - v_+) + a_- \ln(v - v_-) &= \ln X + \text{const} \\ \implies (v - v_+)^{a_+} (v - v_-)^{a_-} &= C X. \end{aligned}$$

Returning to  $(X, Y)$  via  $v - v_{\pm} = (Y - v_{\pm}X)/X$ , the powers of  $X$  collect into a single factor that cancels against the right-hand side:

$$\begin{aligned} (v - v_+)^{a_+} (v - v_-)^{a_-} &= (Y - v_+X)^{a_+} (Y - v_-X)^{a_-} X^{-(a_+ + a_-)} \\ &= (Y - v_+X)^{a_+} (Y - v_-X)^{a_-} X \quad (\text{by } a_+ + a_- = -1), \end{aligned}$$

so  $(Y - v_+X)^{a_+} (Y - v_-X)^{a_-} = C$ . With  $X = x - 1$ ,  $Y = y - 1$  we obtain the first integral

$$\Phi(x, y) := [(y - 1) - v_+(x - 1)]^{a_+} [(y - 1) - v_-(x - 1)]^{a_-} = C_1, \quad (177)$$

constant along each characteristic. The first integral (177) is a product of non-integer powers of two linear forms — a transcendental level curve, not a linear one. As a consistency check, let  $\theta_1, \theta_2 \rightarrow 0^+$ : the quadratic factorises,

$$Q(v) \rightarrow \gamma_1 v^2 + (\gamma_2 - \gamma_1)v - \gamma_2 = (\gamma_1 v + \gamma_2)(v - 1),$$

so  $v_+ \rightarrow 1$  and  $v_- \rightarrow -\gamma_2/\gamma_1$ , and the closed forms (176) give

$$a_+ \rightarrow \frac{\gamma_1 - \gamma_1 \cdot 1}{\gamma_1(v_+ - v_-)} = 0, \quad a_- \rightarrow \frac{\gamma_1 - \gamma_1 v_-}{\gamma_1(v_- - v_+)} = \frac{1 - v_-}{v_- - 1} = -1.$$

The  $v_+$  factor of  $\Phi$  then carries exponent 0 and drops out, leaving  $\Phi(x, y) \rightarrow [(y - 1) + (\gamma_2/\gamma_1)(x - 1)]^{-1}$ , whose level curves are exactly the straight lines  $\gamma_2 x + \gamma_1 y = \text{const}$  of the jockeying-only case.

To find  $P$  along a characteristic we take the ratio of the first and third differentials in (170),

$$\frac{dP}{P} = \frac{-A(x, y(x))}{x[\gamma_1(x - y(x)) + \theta_1(x - 1)]} dx, \quad (178)$$

where  $y(x) = y(x; C_1)$  is the characteristic (177). Integrating yields the general homogeneous solution

$$P_h(x, y) = F(C_1) \exp\left(-\int_{x_*}^x \frac{A(s, y(s))}{s[\gamma_1(s - y(s)) + \theta_1(s - 1)]} ds\right), \quad C_1 = \Phi(x, y), \quad (179)$$

where  $F$  is an arbitrary differentiable function of the characteristic constant and  $x_*$  is an arbitrary base point (its choice is absorbed into  $F$ ). Here the closed-form algebra ends: because  $y(s)$  is defined only implicitly by the transcendental level set (177), the integrand in (179) is not a rational function of  $s$ , and the quadrature is not elementary. No explicit closed form for  $P_h$  exists.

## Step 2: Boundary behaviour of $P_h$ .

Although the quadrature (179) is not elementary, the two qualitative features needed downstream can be established directly from the local behaviour of its integrand.

(a) *Blow-up at  $x = 0$ .* A characteristic meets the axis  $x = 0$  at a finite height  $y_0 := y(0; C_1) \in (0, 1]$ . Evaluating the two ingredients of the exponent integrand there: from (168),  $A(0, y_0) = -\mu$ , and the convection bracket is  $\gamma_1(0 - y_0) + \theta_1(0 - 1) = -(\gamma_1 y_0 + \theta_1) < 0$ . Hence, as  $s \rightarrow 0^+$ ,

$$-\frac{A(s, y(s))}{s[\gamma_1(s - y(s)) + \theta_1(s - 1)]} \sim -\frac{-\mu}{s \cdot [-(\gamma_1 y_0 + \theta_1)]} = -\frac{\mu}{(\gamma_1 y_0 + \theta_1)s},$$

a simple pole whose integral contributes  $-\mu/(\gamma_1 y_0 + \theta_1) \cdot \ln x$  to the exponent in (179), so

$$P_h(x, y(x)) \sim x^{-\mu/(\gamma_1 y_0 + \theta_1)} \rightarrow +\infty \quad (x \rightarrow 0^+), \quad (180)$$

with a strictly negative exponent because  $\gamma_1 y_0 + \theta_1 > 0$ . This blow-up drives the pinning of  $F$  in Step 4.

(b) *Regularity on the diagonal.* On the diagonal  $x = y = z$  the  $x$ -convection coefficient reduces to

$$x[\gamma_1(x - y) + \theta_1(x - 1)] \Big|_{x=y=z} = z\theta_1(z - 1) \neq 0 \quad \text{for } z \in (0, 1):$$

the abandonment term  $\theta_1(x - 1)$  keeps the characteristic field away from zero there, even though the jockeying part  $\gamma_1(x - y)$  vanishes. Consequently the integrand in (179) is regular as  $x \rightarrow z$  along the diagonal-crossing characteristic, and

$$P_h^{\text{diag}}(z) := \lim_{x \rightarrow z} P_h(x, y(x; z)) \quad \text{is finite and non-zero.} \quad (181)$$

Were  $\theta_1$  zero, the coefficient would vanish on the diagonal and equip  $P_h$  with an interior singularity there — the singularity that dominates the jockeying-only analyses (§§6–8). With abandonment present no such singularity arises; as Step 5 shows, the obstruction migrates instead to the right-hand side of the Volterra equation.

*Branch convention.* The non-integer powers in (177) and the quadrature in (179) may carry a complex phase off the real region; this phase is absorbed into the arbitrary function  $F(C_1)$

and cancels exactly in the ratio  $P_h(x, y(x))/P_h(t, y(t))$  appearing in (166), since numerator and denominator lie on the same characteristic (same  $C_1$ ).

**Step 3: Particular solution via variation of parameters.**

We now restrict the full (inhomogeneous) fundamental equation (45) to a fixed characteristic  $y = y(x; C_1)$ . Along it the total derivative of  $P$  is

$$\frac{d}{dx} P(x, y(x)) = \partial_x P + y'(x) \partial_y P,$$

and the characteristic ODE (171), cleared of its denominator, states precisely that

$$\gamma_2(y - x) + \theta_2(y - 1) = [\gamma_1(x - y) + \theta_1(x - 1)] y'(x).$$

Substituting this for the second convection bracket, the two convection terms combine into a single total derivative:

$$\begin{aligned} & xy[\gamma_1(x - y) + \theta_1(x - 1)] \partial_x P + xy[\gamma_2(y - x) + \theta_2(y - 1)] \partial_y P \\ &= xy[\gamma_1(x - y) + \theta_1(x - 1)] (\partial_x P + y'(x) \partial_y P) = xy[\gamma_1(x - y) + \theta_1(x - 1)] \frac{d}{dx} P(x, y(x)). \end{aligned}$$

The fundamental equation along the characteristic therefore reads, using the factorisation (168) of its algebraic coefficient,

$$y A(x, y) P + xy[\gamma_1(x - y) + \theta_1(x - 1)] \frac{dP}{dx} = G(x, y),$$

and dividing through by  $xy[\gamma_1(x - y) + \theta_1(x - 1)]$  reduces the PDE to the first-order linear ODE (for fixed  $C_1$ )

$$\frac{dP}{dx} = f(x; C_1) P + h(x; C_1), \quad (182)$$

with coefficient and forcing

$$f(x; C_1) = -\frac{A(x, y(x))}{x[\gamma_1(x - y(x)) + \theta_1(x - 1)]}, \quad (183)$$

$$h(x; C_1) = \frac{G(x, y(x))}{x y(x) [\gamma_1(x - y(x)) + \theta_1(x - 1)]}, \quad (184)$$

where  $A$  is as in (168) and  $G$  denotes the right-hand side of the fundamental equation,

$$G(x, y) := \mu(x - y) P_y(y) - \mu x(1 - y) \pi(0, 0). \quad (185)$$

Note that the denominator of both (183) and (184) is the  $x$ -convection coefficient itself: every analytic feature of the problem established in Step 2 — the blow-up at  $x = 0$ , the regularity on the diagonal — traces back to its zeros. Since  $P_h$  from (179) satisfies the homogeneous equation  $\frac{d}{dx} P_h = f(x; C_1) P_h$ , it is the fundamental solution of (182); applying variation of parameters (§3.6.2), the general solution of the inhomogeneous ODE is

$$P(x, y(x)) = P_h(x, y(x)) \left[ F(C_1) + \int_{x_0}^x \frac{h(t; C_1)}{P_h(t, y(t))} dt \right]. \quad (186)$$

**Step 4: Pinning the free function from the boundary condition at  $x = 0$ .**

Along any characteristic the endpoint at  $x = 0$  is  $y(0) = y_0$ , giving  $P(0, y_0) = P_y(y_0)$ , which is finite. However, by (180) the homogeneous factor blows up,  $P_h(x, y(x)) \rightarrow +\infty$  as  $x \rightarrow 0^+$ , while the integral in (186) vanishes as  $x_0 \rightarrow 0$  (the integration interval collapses). Therefore

$$\lim_{x \rightarrow 0^+} P(x, y(x)) = \lim_{x \rightarrow 0^+} P_h(x, y(x)) \cdot F(C_1) = +\infty \cdot F(C_1),$$

which can be finite only if  $F(C_1) = 0$ . Setting  $x_0 = 0$  and  $F(C_1) = 0$  in (186) gives the integral representation of Claim 2,

$$P(x, y(x)) = P_h(x, y(x)) \int_0^x \frac{h(t; C_1)}{P_h(t, y(t))} dt. \quad (187)$$

The blow-up exponent  $\mu/(\gamma_1 y_0 + \theta_1)$  of (180) is strictly positive on every characteristic, so the conclusion  $F \equiv 0$  holds for the entire family.

**Step 5: Reduction to the Volterra equation.**

We simplify the forcing  $h$  on the characteristic. Substituting  $G$  from (185) into (184), then splitting the fraction and cancelling one factor of  $t$  in the second term,

$$\begin{aligned} h(t; C_1) &= \frac{\mu(t - y(t)) P_y(y(t)) - \mu t(1 - y(t)) \pi(0, 0)}{t y(t) [\gamma_1(t - y(t)) + \theta_1(t - 1)]} \\ &= \frac{\mu(t - y(t))}{t y(t) [\gamma_1(t - y(t)) + \theta_1(t - 1)]} P_y(y(t)) - \frac{\mu(1 - y(t))}{y(t) [\gamma_1(t - y(t)) + \theta_1(t - 1)]} \pi(0, 0). \end{aligned} \quad (188)$$

We evaluate (187) along the characteristic through a diagonal point  $(z, z)$ , that is, the curve with  $C_1 = \Phi(z, z)$ ; its parametrisation  $y(t; z)$  satisfies  $y(z; z) = z$ . Setting  $x = z$ ,

$$P(z, z) = P_h(z, z) \int_0^z \frac{h(t; \Phi(z, z))}{P_h(t, y(t; z))} dt. \quad (189)$$

By the regularity (181),  $P_h(z, z) = P_h^{\text{diag}}(z)$  is finite and non-zero and the integrand is regular as  $t \rightarrow z$  (its numerator  $t - y(t; z) \rightarrow 0$ ), so (189) is a *proper* integral and no finite-part regularisation is needed. Substituting the two pieces of (188) and separating the term containing the unknown  $P_y$  from the known remainder, define the kernel and remainder

$$\mathcal{K}(z, t) := \frac{\mu(t - y(t; z)) P_h^{\text{diag}}(z)}{t y(t; z) [\gamma_1(t - y(t; z)) + \theta_1(t - 1)] P_h(t, y(t; z))}, \quad (190)$$

$$\mathcal{R}(z) := -\mu \pi(0, 0) P_h^{\text{diag}}(z) \int_0^z \frac{1 - y(t; z)}{y(t; z) [\gamma_1(t - y(t; z)) + \theta_1(t - 1)] P_h(t, y(t; z))} dt, \quad (191)$$

so that (189) reads  $P(z, z) = \int_0^z \mathcal{K}(z, t) P_y(y(t; z)) dt + \mathcal{R}(z)$ . It remains to express the left-hand side  $P(z, z)$ . Solving the diagonal relation (164) for  $P(z, z)$  and using  $\mu/(\mu - \lambda z) = 1/(1 - \rho z)$ ,

$$\begin{aligned} (\mu - \lambda z) P(z, z) &= \mu \pi(0, 0) - z [\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)} \\ \Rightarrow P(z, z) &= \frac{\pi(0, 0)}{1 - \rho z} - \frac{z}{\mu - \lambda z} [\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)}. \end{aligned} \quad (192)$$

Setting

$$\mathcal{F}(z) := P(z, z) - \mathcal{R}(z) = \underbrace{\frac{\pi(0, 0)}{1 - \rho z}}_{\theta_1 = \theta_2 = 0 \text{ value}} - \underbrace{\frac{z}{\mu - \lambda z} [\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)}}_{\text{abandonment coupling}} - \mathcal{R}(z), \quad (193)$$

equation (189) becomes the Volterra integral equation of the first kind (167), confirming Claim 2. Two features distinguish (167) from its Model- $B$  analogue, and both make Model- $X$  strictly less tractable. First, the kernel (190) is built from the non-elementary homogeneous solution  $P_h$  along transcendental characteristics, so it has no closed form for general parameters (in Model- $B$ ,  $P_h$  and the straight-line characteristics were explicit). Second, and decisively, the right-hand side (193) is *not* known data: through (164) it is coupled to the diagonal gradient  $[\theta_1 \partial_x P + \theta_2 \partial_y P]_{(z, z)}$ , which is precisely the quantity that abandonment prevents the diagonal from determining (278). Where Model- $B$  left a clean first-kind Volterra equation with explicit kernel and known forcing (itself already beyond closed-form inversion), Model- $X$  leaves one whose kernel is implicit and whose forcing is entangled with the unknown solution. Resolving it lies well beyond the scope of this thesis.  $\square$

Model- $X$  is therefore left open at this point. Its role in the hierarchy is to make the two obstructions explicit and to show how each tractable specialisation removes them. The limit  $\theta_1 = \theta_2 \rightarrow 0$  (Model- $B$ ) restores both at once: the characteristics (177) straighten to the lines  $\gamma_2 x + \gamma_1 y = \text{const}$ , so the kernel (190) becomes explicit, and the gradient coupling in (193) vanishes, so  $\mathcal{F}(z) = \pi(0, 0)/(1 - \rho z) - \mathcal{R}(z)$  becomes known data — leaving the clean (still unsolved) Volterra equation of §6. Removing one mechanism at a time then closes the problem entirely: setting in addition  $\gamma_2 = 0$  (Model- $B_2$ ) collapses the characteristic family to vertical lines and degenerates the PDE to an ODE in  $x$  alone, whose boundary function  $P_y(y)$  is pinned by an analyticity argument

at the interior singular point  $x = y$  (§8); removing jockeying entirely while keeping only class-1 abandonment (Model- $C_2$ ) likewise degenerates the PDE to an ODE in  $x$ , whose  $P_y(y)$  is pinned by a boundedness argument at  $x = 1$  (§9); and zeroing all four parameters returns the fully solvable baseline Model- $A$  (§5). The full model  $X$ , retaining all mechanisms, retains both obstructions simultaneously and is the one case that does not close.

## 8 Analysis of Model- $B_2$ ( $\gamma_1 > 0$ , $\gamma_2 = 0$ , $\theta_1 = \theta_2 = 0$ )

Model- $B_2$  restricts jockeying to the direction  $1 \rightarrow 2$  only ( $\gamma_1 > 0$ ,  $\gamma_2 = 0$ ) and retains no abandonment ( $\theta_1 = \theta_2 = 0$ ). Setting  $\gamma_2 = 0$  eliminates the  $\partial P / \partial y$  term from the fundamental equation, reducing it to a first-order linear ODE in  $x$  for each fixed  $y \in [0, 1]$ . An integrating factor then yields a formal integral expression for  $P(x, y)$ ; the boundary function  $P_y(y)$  is pinned by requiring  $P(\cdot, y)$  to be holomorphic at the interior singular point  $x = y$ , where the integrating factor carries a branch singularity with negative exponent. This analyticity condition replaces the boundedness argument used for Model- $C_2$ .

**Stability.** As in Models  $A$  and  $B$ ,  $\rho = (\lambda_1 + \lambda_2) / \mu < 1$ . Jockeying conserves the total customer count and does not affect the stability condition. Figure 6 shows the queue layout.




Figure 6: Queue layout for Model- $B_2$ . Only the  $1 \rightarrow 2$  jockeying direction is active (dashed, downward): class-1 customers may defect to Queue 2, but no customer can move in the reverse direction ( $\gamma_2 = 0$ ).

### 8.1 Reduced fundamental equation

Setting  $\gamma_2 = 0$  and  $\theta_1 = \theta_2 = 0$  in the general fundamental equation (45) (Lemma 1) annihilates the  $\partial P / \partial y$  term, whose coefficient  $xy[\gamma_2(y - x) + \theta_2(y - 1)]$  vanishes identically, and leaves only the  $\gamma_1$  contribution to the  $\partial P / \partial x$  term:

$$\begin{aligned} & [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P + \gamma_1 x y (x - y) \frac{\partial P}{\partial x} \\ & = \mu(x - y) P_y(y) - \mu x(1 - y) \pi(0, 0). \end{aligned} \quad (194)$$

For each fixed  $y \in (0, 1]$  this is a first-order linear ODE in  $x$ ; the entire analysis of this section is built on it.

**Theorem 2.** *Consider the two-class non-preemptive priority queueing system of Model- $B_2$  with per-customer jockeying rate  $\gamma_1 > 0$  from Queue 1 to Queue 2 ( $\gamma_2 = 0$ ,  $\theta_1 = \theta_2 = 0$ ). Under the stability condition  $\rho = \lambda / \mu < 1$ ,  $\lambda = \lambda_1 + \lambda_2$ , the boundary generating function is*

$$P_y(y) = \frac{\mu \pi(0, 0)}{\mu - \lambda y} \cdot \frac{{}_1F_1(\alpha(y) + 1; b^*(y); -\lambda_1 y / \gamma_1)}{{}_1F_1(\alpha(y); b^*(y); -\lambda_1 y / \gamma_1)}, \quad (195)$$

and the bivariate PGF, for  $0 \leq x \leq y$ , is

$$P(x, y) = \frac{\mu e^{\lambda_1 x / \gamma_1} x^{-\alpha} (y - x)^{-\beta}}{\gamma_1 y} \left[ P_y(y) \mathcal{H}_1(x, y) + (1 - y) \pi(0, 0) \mathcal{H}_2(x, y) \right], \quad (196)$$

where  ${}_1F_1$  is Kummer's confluent hypergeometric function, the exponents are

$$\alpha(y) = \frac{\mu}{\gamma_1 y}, \quad \beta(y) = \frac{(1 - y)(\lambda y - \mu)}{\gamma_1 y}, \quad b^*(y) = \alpha + \beta + 1 = \frac{\mu + \lambda(1 - y) + \gamma_1}{\gamma_1}, \quad (197)$$



the auxiliary integrals are

$$\mathcal{H}_1(x, y) = \int_0^x e^{-\lambda_1 t / \gamma_1} t^{\alpha-1} (y-t)^\beta dt, \quad (198)$$

$$\mathcal{H}_2(x, y) = \int_0^x e^{-\lambda_1 t / \gamma_1} t^\alpha (y-t)^{\beta-1} dt, \quad (199)$$

and the empty-state probability  $\pi(0, 0) = \rho(1 - \rho)$  is given by Corollary 4 below. On  $y < x \leq 1$  the same expression (196) holds with  $\mathcal{H}_1, \mathcal{H}_2$  replaced by their finite-part continuations.

**Corollary 4.** *The empty-system probability  $\pi_0$  (server idle, no one waiting) and the both-queues-empty probability  $\pi(0, 0)$  (server busy,  $N_1 = N_2 = 0$ ) of Model-B<sub>2</sub> are*

$$\pi_0 = 1 - \rho, \quad \pi(0, 0) = \rho(1 - \rho), \quad (200)$$

identical to the Model-A baseline.

*Proof.* Because one-way jockeying conserves the total customer count and there is no abandonment, the total-count process  $L$  (equal to the number in queue  $N$  plus the single customer in service whenever the server is busy) is itself a birth-death chain: it increases at rate  $\lambda = \lambda_1 + \lambda_2$  on any arrival and decreases at rate  $\mu$  on any service completion, irrespective of class. Thus  $L$  is an  $M/M/1$  queue, and by the memorylessness of the exponential clocks its stationary law is geometric,  $\mathbb{P}(L = k) = (1 - \rho)\rho^k$ ; in particular  $\pi_0 = \mathbb{P}(L = 0) = 1 - \rho$ .

For  $\pi(0, 0)$ , set  $x = y$  in the reduced fundamental equation (194). The convection term  $\gamma_1 xy(x - y)\partial_x P$  and the boundary term  $\mu(x - y)P_y(y)$  both carry the factor  $(x - y)$  and vanish on the diagonal; dividing the surviving terms by the common factor  $x$  leaves

$$[(\lambda_1 + \lambda_2 + \mu)x - \mu - (\lambda_1 + \lambda_2)x^2]P(x, x) = -\mu(1 - x)\pi(0, 0). \quad (201)$$

Factoring the bracket and cancelling  $(x - 1)$  against  $-(1 - x) = (x - 1)$  on the right:

$$\begin{aligned} [-\lambda x^2 + (\lambda + \mu)x - \mu]P(x, x) &= (\mu - \lambda x)(x - 1)P(x, x) \\ &= \mu(x - 1)\pi(0, 0), \end{aligned} \quad (202)$$

so that

$$P(x, x) = \frac{\mu\pi(0, 0)}{\mu - \lambda x} = \frac{\pi(0, 0)}{1 - \rho x}. \quad (203)$$

Evaluating at  $x = 1$  with  $P(1, 1) = 1 - \pi_0 = \rho$  (using  $\pi_0 = 1 - \rho$  just established) yields  $\pi(0, 0) = \rho(1 - \rho)$ .  $\square$

*Proof of Theorem 2. Stage 1: Standard form and the integrating factor.*

For any fixed  $y \neq 0$ , divide the reduced fundamental equation (194) through by the coefficient  $\gamma_1 xy(x - y)$  of  $\partial P / \partial x$ . The factor  $y$  cancels against the overall factor  $y$  in the bracket multiplying  $P$ , putting the equation in standard first-order form  $\frac{dP}{dx} + p(x; y)P = q(x; y)$  with

$$p(x; y) = \frac{(\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy}{\gamma_1 x(x - y)}, \quad (204)$$

$$q(x; y) = \frac{\mu P_y(y)}{\gamma_1 xy} - \frac{\mu(1 - y)\pi(0, 0)}{\gamma_1 y(x - y)}. \quad (205)$$

Write the numerator of  $p$  as a polynomial in  $x$ ,

$$N(x) = -\lambda_1 x^2 + [(\lambda_1 + \lambda_2 + \mu) - \lambda_2 y]x - \mu = -\lambda_1(x - x^*(y))(x - x^+(y)),$$

which factors through the roots  $x^*(y), x^+(y)$  of the Model-A kernel (Equation (87)), with  $\lambda_1 x^* x^+ = \mu$ . The partial-fraction coefficients of  $p = N(x) / [\gamma_1 x(x - y)]$  follow by residues:

$$\begin{aligned} A &= -\lambda_1 && \text{(ratio of leading } x^2 \text{ terms),} \\ B &= \frac{N(0)}{0 - y} = \frac{-\mu}{-y} = \frac{\mu}{y}, \\ C &= \frac{N(y)}{y} = \frac{-(\lambda y - \mu)(y - 1)}{y} = \frac{(1 - y)(\lambda y - \mu)}{y}, \end{aligned} \quad (206)$$



where  $N(y) = -\lambda y^2 + (\lambda + \mu)y - \mu = -(\lambda y - \mu)(y - 1)$ . Hence

$$p(x; y) = \frac{1}{\gamma_1} \left[ -\lambda_1 + \frac{\mu/y}{x} + \frac{(1-y)(\lambda y - \mu)/y}{x-y} \right]. \quad (207)$$

We henceforth work on the principal triangle  $0 \leq x \leq y$ , on which  $x - y \leq 0$ ; the complementary region  $y < x \leq 1$  is recovered at the end by analytic continuation. There the real antiderivative of the simple-pole term is  $\beta \ln|x - y| = \beta \ln(y - x)$ , so integrating the partial-fraction form (207) term by term gives, up to an additive constant,

$$\int^x p(t; y) dt = \frac{1}{\gamma_1} \left[ -\lambda_1 x + \frac{\mu}{y} \ln x + \frac{(1-y)(\lambda y - \mu)}{y} \ln(y - x) \right].$$

Exponentiating (an integrating factor is determined only up to a multiplicative constant, which we drop) yields the *real* integrating factor

$$\mu_I(x; y) = e^{-\lambda_1 x / \gamma_1} x^{\alpha(y)} (y - x)^{\beta(y)}, \quad (208)$$

with exponents  $\alpha(y) = \mu/(\gamma_1 y) > 0$  and  $\beta(y) = (1-y)(\lambda y - \mu)/(\gamma_1 y)$  as in (197); one verifies directly that  $\mu'_I/\mu_I = -\lambda_1/\gamma_1 + \alpha/x + \beta/(x-y) = p(x; y)$ , using  $\frac{d}{dx} \ln(y-x)^\beta = \beta/(x-y)$ . Working with  $(y-x) > 0$  in place of the complex base  $(x-y)$  keeps every quantity manifestly real throughout the principal triangle. Since  $\alpha > 0$  we have  $\mu_I \rightarrow 0$  as  $x \rightarrow 0^+$ , so the boundary term  $P(0, y) \mu_I(0; y)$  vanishes when we integrate  $\frac{d}{dx}[P \cdot \mu_I] = q \cdot \mu_I$  over  $(0, x]$ :

$$P(x, y) \mu_I(x; y) = \int_0^x q(t; y) \mu_I(t; y) dt. \quad (209)$$

It remains to evaluate the right-hand side. Multiplying  $q$  from (205) by  $\mu_I$  and rewriting the second term with the elementary identity  $(y-t)^\beta/(t-y) = -(y-t)^{\beta-1}$ , the integrand separates into two manifestly real pieces,

$$q(t; y) \mu_I(t; y) = \frac{\mu}{\gamma_1 y} \left[ P_y(y) e^{-\lambda_1 t / \gamma_1} t^{\alpha-1} (y-t)^\beta + (1-y) \pi(0, 0) e^{-\lambda_1 t / \gamma_1} t^\alpha (y-t)^{\beta-1} \right], \quad (210)$$

the minus sign of (205) having been absorbed by that identity. With the auxiliary integrals  $\mathcal{H}_1, \mathcal{H}_2$  of (198)–(199) and dividing through by  $\mu_I(x; y) = e^{-\lambda_1 x / \gamma_1} x^\alpha (y-x)^\beta$ , the solution on the principal triangle  $0 \leq x \leq y$  is

$$P(x, y) = \frac{\mu e^{\lambda_1 x / \gamma_1} x^{-\alpha} (y-x)^{-\beta}}{\gamma_1 y} \left[ P_y(y) \mathcal{H}_1(x, y) + (1-y) \pi(0, 0) \mathcal{H}_2(x, y) \right], \quad (211)$$

which still contains the unknown boundary function  $P_y(y)$ ; Stage 2 determines it.

*Stage 2: Determination of  $P_y(y)$  by analyticity at  $x = y$ .*

The factor  $(x-y)$  surviving in the  $x$ -convection coefficient places the operative singularity at the interior point  $x = y$ , exactly the analyticity-pinning regime anticipated in the remark following Lemma 1. Unlike Models A and  $C_2$ , no probabilistic (Pollaczek–Khinchine) route is available here. One-way jockeying makes the effective service experienced by a class-2 customer depend on the class-1 queue contents at the (random) jockeying instants, so the class-2 input is no longer a Poisson stream feeding an  $M/G/1$  subsystem and no busy-period reduction applies (cf. the discussion of Model-B in §6). We therefore determine  $P_y(y)$  *purely analytically*, by imposing holomorphy of  $P(\cdot, y)$  at the interior point  $x = y$ , where the solution (211) is *a priori* singular.

Under stability  $\mu - \lambda y > 0$  for all  $y \in (0, 1)$ , so the second exponent is negative,

$$\beta(y) = \frac{(1-y)(\lambda y - \mu)}{\gamma_1 y} < 0, \quad 0 < y < 1. \quad (212)$$

The prefactor  $(y-x)^{-\beta} = (y-x)^{|\beta|}$  in (211) therefore *vanishes* as  $x \rightarrow y^-$ , while the integrals  $\mathcal{H}_1, \mathcal{H}_2$  may *diverge* there; the product is an indeterminate  $0 \cdot \infty$  form. Because  $P(\cdot, y)$  is a probability generating function it must be holomorphic on the open unit disc, and in particular analytic at the interior point  $x = y$ . As we show next, the general solution carries a non-integer power  $(y-x)^{|\beta|}$  — a branch term that vanishes at  $x = y$  yet is not analytic there — and holomorphy forces its coefficient to zero; this single condition pins down  $P_y(y)$ .

*Local analysis at  $x = y$ .* To exhibit the two modes explicitly, read off the local behaviour of the ODE (204)–(205) at  $x = y$ . By the partial fraction (207),  $p$  has a simple pole at  $x = y$  with residue

$\beta$ ; by (205),  $q$  has the leading singularity  $-D/(x-y)$  with  $D = \mu(1-y)\pi(0,0)/(\gamma_1 y)$ . The local model equation is therefore

$$P'(x, y) + \frac{\beta}{x-y} P(x, y) = \frac{-D}{x-y} + (\text{analytic near } x=y). \quad (213)$$

A constant particular solution  $P \equiv P_*$  requires  $\beta P_* = -D$ , so, cancelling the common factor  $(1-y)/(\gamma_1 y)$  shared by  $D$  and  $\beta$ ,

$$P(y, y) = P_* = -\frac{D}{\beta} = -\frac{\mu \pi(0,0)}{\lambda y - \mu} = \frac{\mu \pi(0,0)}{\mu - \lambda y} = \frac{\rho(1-\rho)}{1-\rho y}, \quad (214)$$

consistent with the diagonal value (203) (using  $\pi(0,0) = \rho(1-\rho)$ ). The general local solution adds a multiple of the homogeneous mode  $(x-y)^{-\beta} = (x-y)^{|\beta|}$ . Since  $|\beta| \notin \mathbb{Z}$  generically this is a branch term: it vanishes at  $x=y$  but is not analytic there, so holomorphy forces its coefficient to zero. The remaining task is to locate that coefficient inside the explicit solution (211).

*No-branch condition and  $P_y(y)$ .* Write the analytic, non-vanishing prefactor of (211) as  $\Psi(x) = \mu e^{\lambda_1 x / \gamma_1} x^{-\alpha} / (\gamma_1 y)$ , so that  $P(x, y) = \Psi(x) (y-x)^{-\beta} [P_y(y) \mathcal{H}_1 + (1-y)\pi(0,0) \mathcal{H}_2]$ . As  $x \rightarrow y^-$  each integral admits a local expansion, obtained by Taylor-expanding its smooth integrand factor —  $g_1(t) = e^{-\lambda_1 t / \gamma_1} t^{\alpha-1}$  for  $\mathcal{H}_1$ ,  $g_2(t) = e^{-\lambda_1 t / \gamma_1} t^\alpha$  for  $\mathcal{H}_2$  — about  $t=y$  and integrating the leading power of  $(y-t)$ :

$$\mathcal{H}_1(x, y) = \text{f.p. } \mathcal{H}_1(y, y) - \frac{g_1(y)}{\beta+1} (y-x)^{\beta+1} + \dots, \quad (215)$$

$$\mathcal{H}_2(x, y) = \text{f.p. } \mathcal{H}_2(y, y) - \frac{g_2(y)}{\beta} (y-x)^\beta + \dots, \quad (216)$$

where f.p. denotes the Hadamard finite part (the  $x \rightarrow y$  limit of the integral once its singular term has been subtracted) and the omitted terms are of higher order. Multiplying by  $\Psi(x)(y-x)^{-\beta}$  sorts the contributions to  $P(x, y)$  by their power of  $(y-x)$ :

- the singular part of  $\mathcal{H}_2$  gives  $-\Psi(y)(1-y)\pi(0,0) g_2(y)/\beta = -D/\beta = P_*$ , exactly the regular value (214) (using  $\Psi(y) g_2(y) = \mu/(\gamma_1 y)$ );
- the singular part of  $\mathcal{H}_1$  gives a term  $\propto (y-x)^1$ , analytic at  $x=y$ ;
- the finite parts, riding on the prefactor, produce the branch term  $\Psi(x)(y-x)^{-\beta} [P_y(y) \text{f.p. } \mathcal{H}_1 + (1-y)\pi(0,0) \text{f.p. } \mathcal{H}_2]$ .

Only the last is non-analytic ( $(y-x)^{|\beta|}$  with  $|\beta| \notin \mathbb{Z}$ ), so holomorphy of  $P(\cdot, y)$  at  $x=y$  requires its bracket to vanish:

$$P_y(y) \text{f.p. } \mathcal{H}_1(y, y) + (1-y) \pi(0,0) \text{f.p. } \mathcal{H}_2(y, y) = 0. \quad (217)$$

To evaluate the finite parts, substitute  $t = ys$  ( $dt = y ds$ ) in (198)–(199) and pull out  $y^{\alpha+\beta}$  from  $t^{\alpha-1}(y-t)^\beta dt = y^{\alpha+\beta} s^{\alpha-1}(1-s)^\beta ds$  (resp.  $t^\alpha(y-t)^{\beta-1} dt = y^{\alpha+\beta} s^\alpha(1-s)^{\beta-1} ds$ ); with  $c := \lambda_1 y / \gamma_1$ ,

$$\mathcal{H}_1(y, y) = y^{\alpha+\beta} \int_0^1 s^{\alpha-1} (1-s)^\beta e^{-cs} ds, \quad \mathcal{H}_2(y, y) = y^{\alpha+\beta} \int_0^1 s^\alpha (1-s)^{\beta-1} e^{-cs} ds.$$

Matching the Euler integral representation (15), written as  $\int_0^1 t^{a-1} (1-t)^{b-1} e^{zt} dt = B(a, b) {}_1F_1(a; a+b; z)$ , with  $(a, b, z) = (\alpha, \beta+1, -c)$  for  $\mathcal{H}_1$  and  $(a, b, z) = (\alpha+1, \beta, -c)$  for  $\mathcal{H}_2$  — read as its analytic continuation in the second parameter  $b$  beyond the convergence domain  $b > 0$ , since  $\beta < 0$  by (212), which is precisely the finite part — and noting  $a+b = b^*(y)$  in both cases, gives

$$\text{f.p. } \mathcal{H}_1(y, y) = B(\alpha, \beta+1) y^{\alpha+\beta} {}_1F_1(\alpha; b^*(y); -\lambda_1 y / \gamma_1), \quad (218)$$

$$\text{f.p. } \mathcal{H}_2(y, y) = B(\alpha+1, \beta) y^{\alpha+\beta} {}_1F_1(\alpha+1; b^*(y); -\lambda_1 y / \gamma_1). \quad (219)$$

Solving the no-branch condition (217) for  $P_y(y)$  and inserting the finite parts (218)–(219), the common factor  $y^{\alpha+\beta}$  cancels and

$$P_y(y) = -(1-y) \pi(0,0) \frac{\text{f.p. } \mathcal{H}_2(y, y)}{\text{f.p. } \mathcal{H}_1(y, y)} = -(1-y) \pi(0,0) \frac{B(\alpha+1, \beta)}{B(\alpha, \beta+1)} \frac{{}_1F_1(\alpha+1; b^*; -\lambda_1 y / \gamma_1)}{{}_1F_1(\alpha; b^*; -\lambda_1 y / \gamma_1)}. \quad (220)$$

The Beta ratio reduces, via  $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$  and  $\Gamma(\beta + 1) = \beta\Gamma(\beta)$ , to

$$\begin{aligned} \frac{B(\alpha + 1, \beta)}{B(\alpha, \beta + 1)} &= \frac{\Gamma(\alpha + 1)\Gamma(\beta)}{\Gamma(\alpha + \beta + 1)} \cdot \frac{\Gamma(\alpha + \beta + 1)}{\Gamma(\alpha)\Gamma(\beta + 1)} \\ &= \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha)} \cdot \frac{\Gamma(\beta)}{\Gamma(\beta + 1)} = \frac{\alpha}{\beta}, \end{aligned} \quad (221)$$

and the surviving prefactor simplifies, using  $\alpha/\beta = \mu/[(1-y)(\lambda y - \mu)]$ , as

$$\begin{aligned} -(1-y) \frac{\alpha}{\beta} &= -(1-y) \cdot \frac{\mu}{(1-y)(\lambda y - \mu)} \\ &= -\frac{\mu}{\lambda y - \mu} = \frac{\mu}{\mu - \lambda y}. \end{aligned} \quad (222)$$

Substituting (221)–(222) into (220) gives the boundary generating function

$$P_y(y) = \frac{\mu \pi(0, 0)}{\mu - \lambda y} \cdot \frac{{}_1F_1(\alpha(y) + 1; b^*(y); -\lambda_1 y/\gamma_1)}{{}_1F_1(\alpha(y); b^*(y); -\lambda_1 y/\gamma_1)}, \quad (223)$$

confirming (195). The prefactor equals the regular value  $P(y, y)$  of (214), and the Kummer ratio is the boundary correction. As  $y \rightarrow 0^+$  the argument  $-\lambda_1 y/\gamma_1 \rightarrow 0$ , so the ratio tends to 1 and  $P_y(0) = \pi(0, 0)$ , consistent with  $P_y(0) = \sum_{n_2} \pi(0, n_2) \cdot 0^{n_2} = \pi(0, 0)$ .

*Stage 3: Recovery of  $P(x, y)$ .*

Substituting (223) into the real-form expression (211) gives, for  $0 \leq x \leq y$ ,

$$\begin{aligned} P(x, y) &= \frac{\mu e^{\lambda_1 x/\gamma_1} x^{-\alpha} (y-x)^{-\beta}}{\gamma_1 y} \left[ \frac{\mu \pi(0, 0)}{\mu - \lambda y} \cdot \frac{{}_1F_1(\alpha + 1; b^*; -\frac{\lambda_1 y}{\gamma_1})}{{}_1F_1(\alpha; b^*; -\frac{\lambda_1 y}{\gamma_1})} \cdot \mathcal{H}_1(x, y) \right. \\ &\quad \left. + (1-y)\pi(0, 0) \cdot \mathcal{H}_2(x, y) \right], \end{aligned} \quad (224)$$

which is the statement of the theorem (Equation (196)). The no-branch condition (217) guarantees that this extends analytically across  $x = y$ , with diagonal value  $P(y, y) = \rho(1 - \rho)/(1 - \rho y)$  as computed in (214). On  $y < x \leq 1$  the same formula holds with  $\mathcal{H}_1, \mathcal{H}_2$  read as their finite-part continuations;  $\pi(0, 0) = \rho(1 - \rho)$  from Corollary 4 makes the solution fully explicit. The recovery of Model-A in the limit  $\gamma_1 \rightarrow 0^+$  is deferred to Remark 1 below, as the limit is singular and merits separate discussion.  $\square$

*Remark 1* (The singular limit  $\gamma_1 \rightarrow 0^+$ ). The no-jockeying limit is cleanest at the level of the fundamental equation, but  $\gamma_1 \rightarrow 0^+$  is a *singular* perturbation — the coefficient  $\gamma_1 xy(x - y)$  of the highest derivative  $\partial_x P$  vanishes — so one must check that the convection term itself, and not merely its coefficient, disappears; the derivative  $\partial_x P$  could *a priori* blow up. The probabilistic bound rules this out. Each  $P(\cdot, \cdot)$  is a probability generating function, so  $|P| \leq 1$  on the closed unit polydisc *uniformly* in  $\gamma_1$ ; by Cauchy's estimate  $\partial_x P$  is then bounded uniformly in  $\gamma_1$  on every compact subset of the open polydisc, whence  $\gamma_1 xy(x - y) \partial_x P \rightarrow 0$  there. The fundamental equation (194) therefore degenerates to the purely algebraic Model-A equation (80) (the coefficients of  $P$  and the right-hand side are  $\gamma_1$ -independent, and  $\pi(0, 0) = \rho(1 - \rho)$  by Corollary 4). Since  $\{P\}_{\gamma_1}$  is uniformly bounded it is a normal family, so it converges — along subsequences, hence in full by uniqueness of the polydisc-analytic generating-function solution of (80) — to the Model-A PGF (81). In particular the boundary function reduces to the Model-A value (89),

$$P_y(y) \xrightarrow{\gamma_1 \rightarrow 0^+} \pi(0, 0) \frac{x^*(y)(1-y)}{x^*(y) - y}, \quad (225)$$

with  $x^*(y)$  the Model-A kernel root (87); this recovery needs no special-function asymptotics.

It is instructive to read the same limit off the closed form (223). Its prefactor satisfies  $\mu \pi(0, 0)/(\mu - \lambda y) = \pi(0, 0)/(1 - \rho y)$  and is  $\gamma_1$ -independent, so (225) is equivalent to the Kummer ratio tending to

$$\frac{{}_1F_1(\alpha + 1; b^*; -\lambda_1 y/\gamma_1)}{{}_1F_1(\alpha; b^*; -\lambda_1 y/\gamma_1)} \xrightarrow{\gamma_1 \rightarrow 0^+} (1 - \rho y) \frac{x^*(y)(1-y)}{x^*(y) - y}. \quad (226)$$

Note the factor  $(1 - \rho y)$ : the ratio does *not* tend to the bare rational expression  $x^*(y)(1-y)/(x^*(y) - y)$ , the discrepancy being precisely the prefactor. This is consistent with the three-term recurrence for  ${}_1F_1$  in its first parameter [DLM, §13.3]: as  $\gamma_1 \rightarrow 0^+$  the parameters  $\alpha = \mu/(\gamma_1 y)$ ,  $b^*$  and  $c = \lambda_1 y/\gamma_1$  all grow like  $1/\gamma_1$ , and to leading order that recurrence reduces to a quadratic in the ratio whose admissible root is the right-hand side of (226).

## 8.2 Mean queue lengths and mean waiting times

### Total queue length from the diagonal

Setting  $x = y = z$  in the reduced fundamental equation (194), the jockeying convection term  $\gamma_1 xy(x-y)\partial_x P$  vanishes at  $x = y = z$  (factor  $(x-y) = 0$ ), as does the boundary term  $\mu(x-y)P_y(y)$ . The remaining balance is identical to Model-A's, so the same diagonal cancellation gives:

$$P(z, z) = \frac{\rho(1-\rho)}{1-\rho z}, \quad (227)$$

the same formula as (120). Since  $P(z, z) = \mathbb{E}[z^{N_1+N_2}] = \mathbb{E}[z^N]$  is the PGF of the total in-queue count  $N$ , the generating-function identity  $\frac{d}{dz}P(z, z)|_{z=1} = \mathbb{E}[N_1] + \mathbb{E}[N_2]$  holds; differentiating by the quotient rule and evaluating at  $z = 1$ :

$$\mathbb{E}[N_1] + \mathbb{E}[N_2] = \frac{d}{dz} \frac{\rho(1-\rho)}{1-\rho z} \Big|_{z=1} = \frac{\rho(1-\rho) \cdot \rho}{(1-\rho z)^2} \Big|_{z=1} = \frac{\rho^2(1-\rho)}{(1-\rho)^2} = \frac{\rho^2}{1-\rho}. \quad (228)$$

One-way jockeying does not alter the total mean queue length relative to Model-A, since the diagonal PGF is unchanged.

### Class-1 mean queue length

We extract  $\mathbb{E}[N_1] = \frac{\partial P}{\partial x} \Big|_{(1,1)}$  by first reducing the fundamental equation (194) at  $y = 1$ , and then differentiating the resulting marginal PGF at  $x = 1$ .

*Step 1: marginal PGF  $P(x, 1)$ .* Evaluating (194) at  $y = 1$ , the bracket multiplying  $P$  becomes  $(\lambda_1 + \mu)x - \mu - \lambda_1 x^2 = -\lambda_1(x-1)(x-1/\rho_1)$ , the convection coefficient is  $\gamma_1 x(x-1)$ , and the right-hand side reduces to  $\mu(x-1)P_y(1)$  (the  $\pi(0, 0)$  term carries the factor  $(1-y) = 0$ ). Every term therefore shares the common factor  $(x-1)$ ; cancelling it leaves the first-order linear ODE in  $x$ :

$$-\lambda_1(x-1/\rho_1)P(x, 1) + \gamma_1 x \frac{d}{dx}P(x, 1) = \mu P_y(1). \quad (229)$$

The integrating factor for this ODE is  $e^{-\lambda_1 x/\gamma_1} x^{\mu/\gamma_1}$  (noting  $\lambda_1/\rho_1 = \mu$ ), and integrating from 0 to  $x$  (the boundary term vanishes since  $0^{\mu/\gamma_1} = 0$ ):

$$P(x, 1) e^{-\lambda_1 x/\gamma_1} x^{\mu/\gamma_1} = \frac{\mu P_y(1)}{\gamma_1} \int_0^x e^{-\lambda_1 t/\gamma_1} t^{\mu/\gamma_1-1} dt. \quad (230)$$

Rearranging and writing  $\alpha = \mu/\gamma_1$  and  $\mathcal{J}(x) = \int_0^x e^{-\lambda_1 t/\gamma_1} t^{\alpha-1} dt$ :

$$P(x, 1) = \frac{\mu P_y(1)}{\gamma_1} \cdot e^{\lambda_1 x/\gamma_1} x^{-\alpha} \mathcal{J}(x). \quad (231)$$

The constant  $P_y(1)$  follows from  $P(1, 1) = 1 - \pi_0 = \rho$ , which gives  $P_y(1) = \rho\gamma_1/(\mu e^{\lambda_1/\gamma_1} \mathcal{J}(1))$ .

*Step 2: differentiate  $P(x, 1)$  at  $x = 1$ .* Writing  $P(x, 1) = C \cdot f(x) \cdot \mathcal{J}(x)$  with  $C = \mu P_y(1)/\gamma_1$  and  $f(x) = e^{\lambda_1 x/\gamma_1} x^{-\alpha}$ , the product rule together with Leibniz's rule ( $d\mathcal{J}/dx = e^{-\lambda_1 x/\gamma_1} x^{\alpha-1}$ , the integrand at the upper limit) gives:

$$\frac{d}{dx}P(x, 1) = C \left[ f'(x) \mathcal{J}(x) + f(x) \cdot e^{-\lambda_1 x/\gamma_1} x^{\alpha-1} \right]. \quad (232)$$

The logarithmic derivative  $f'(x)/f(x) = \lambda_1/\gamma_1 - \alpha/x = (\lambda_1 x - \mu)/(\gamma_1 x)$ , and the second product term simplifies as  $f(x) e^{-\lambda_1 x/\gamma_1} x^{\alpha-1} = x^{-1}$ , so:

$$\frac{d}{dx}P(x, 1) = \frac{\mu P_y(1)}{\gamma_1} \left[ e^{\lambda_1 x/\gamma_1} x^{-\alpha} \cdot \frac{\lambda_1 x - \mu}{\gamma_1 x} \cdot \mathcal{J}(x) + x^{-1} \right]. \quad (233)$$

*Step 3: evaluate at  $x = 1$ .*

$$\begin{aligned} \mathbb{E}[N_1] &= \frac{d}{dx}P(x, 1) \Big|_{x=1} \\ &= \frac{\mu P_y(1)}{\gamma_1} \left[ e^{\lambda_1/\gamma_1} \cdot \frac{\lambda_1 - \mu}{\gamma_1} \cdot \mathcal{J}(1) + 1 \right], \end{aligned} \quad (234)$$

where  $\mathcal{J}(1) = \int_0^1 e^{-\lambda_1 t/\gamma_1} t^{\mu/\gamma_1-1} dt = (\gamma_1/\lambda_1)^{\mu/\gamma_1} \gamma(\mu/\gamma_1, \lambda_1/\gamma_1)$  and  $\gamma(\cdot, \cdot)$  is the lower incomplete gamma function. The factor  $(\lambda_1 - \mu) < 0$  (since  $\rho_1 = \lambda_1/\mu < 1$ ), confirming that  $e^{\lambda_1/\gamma_1}(\lambda_1 - \mu)\mathcal{J}(1)/\gamma_1 + 1 > 0$ . No elementary simplification of (234) is known; the expression is evaluated numerically. As  $\gamma_1 \rightarrow 0^+$  the Model-A value (118) is recovered.

## Class-2 mean queue length by subtraction

$$\mathbb{E}[N_2] = (\mathbb{E}[N_1] + \mathbb{E}[N_2]) - \mathbb{E}[N_1] = \frac{\rho^2}{1 - \rho} - \mathbb{E}[N_1]. \quad (235)$$

Since jockeying defections reduce the class-1 queue,  $\mathbb{E}[N_1]$  is smaller than the Model-A value  $\rho\rho_1/(1 - \rho_1)$ , so  $\mathbb{E}[N_2]$  correspondingly exceeds the Model-A value.

## Mean waiting times via Little's Law

**Little's Law** applies here without qualification. It is a rate-conservation identity — for any subsystem in which customers are neither created nor destroyed internally, the time-average number present equals the long-run entry rate times the mean time an entrant spends inside — and it holds for *any* positive-recurrent system with finite mean queue length (here ensured by  $\rho < 1$ , which gives  $\mathbb{E}[N_1] \leq \mathbb{E}[N] = \rho^2/(1 - \rho) < \infty$  and so makes the jockeying rate  $\gamma_1 \mathbb{E}[N_1]$  below a genuine finite rate), requiring neither Poisson input nor independence. In particular it remains valid even though one-way jockeying has destroyed the per-class Poisson/renewal structure (the very obstruction that ruled out a probabilistic route in Stage 2). We apply it to each queue (the waiting line, excluding the customer in service), with  $\mathbb{E}[W_i]$  understood as the mean time a customer spends in Queue  $i$  averaged over all customers that enter it — an arrival routed directly into service contributing a zero-length visit.

*Class 1.* Every class-1 arrival is counted as an entrant to Queue 1 at rate  $\lambda_1$  (those finding the server idle enter service at once, contributing zero waiting time), and each leaves Queue 1 either by entering service or by jockeying to Queue 2. Since no other stream feeds Queue 1, its throughput is exactly  $\lambda_1$ , and Little's Law gives

$$\mathbb{E}[W_1] = \frac{\mathbb{E}[N_1]}{\lambda_1}. \quad (236)$$

*Class 2.* Queue 2 has *two* input streams: the class-2 Poisson arrivals at rate  $\lambda_2$  and the jockeying transfers from Queue 1. Because each of the  $N_1$  waiting class-1 customers jockeys independently at rate  $\gamma_1$ , the long-run transfer rate is  $\gamma_1 \mathbb{E}[N_1]$  (a genuine rate by the ergodic theorem, equal to the time average of the instantaneous jockeying intensity  $\gamma_1 N_1$ ). The throughput of Queue 2 is therefore  $\lambda_2 + \gamma_1 \mathbb{E}[N_1]$ , and Little's Law reads  $\mathbb{E}[N_2] = (\lambda_2 + \gamma_1 \mathbb{E}[N_1]) \mathbb{E}[W_2]$ , so that

$$\mathbb{E}[W_2] = \frac{\mathbb{E}[N_2]}{\lambda_2 + \gamma_1 \mathbb{E}[N_1]}. \quad (237)$$

Here  $\mathbb{E}[W_2]$  is the mean time in the class-2 line averaged over *all* Queue-2 entrants of both streams; a jockeyed customer's Queue-2 clock starts at the transfer instant, so its total wait is an initial Queue-1 segment (of mean  $\mathbb{E}[W_1]$ ) followed by this Queue-2 segment. Thus (237) is the per-Queue-2-entrant residence time, not the end-to-end delay of a tagged class-2 *arrival*.

The jockeying inflow is what distinguishes (237) from the single-stream Models  $A$  and  $C_2$ : the naive denominator  $\lambda_2$  would undercount the entrants to Queue 2 and is correct only in the no-jockeying limit. Indeed, as  $\gamma_1 \rightarrow 0^+$  the transfer rate  $\gamma_1 \mathbb{E}[N_1] \rightarrow 0$ , so (237) reduces to  $\mathbb{E}[N_2]/\lambda_2$  and both (236) and (237) recover the Model-A values (123)–(124).

## 9 Analysis of Model- $C_2$ ( $\theta_1 > 0$ , $\gamma_1 = \gamma_2 = \theta_2 = 0$ )

In this section, we will derive  $P(x, y)$  for this model where only true abandonments coming from class-1 take place, while no abandonments in class-2 or jockeying of any sort are taken into account. Figure 7 shows the queue layout.




Figure 7: Queue layout for Model- $C_2$ . Class-1 customers abandon the system at rate  $\theta_1$  per waiting customer (dotted arrow, upward from Queue 1). Class-2 never abandons ( $\theta_2 = 0$ ) and there is no jockeying.

The rationale behind using this combination of parameters is that, thanks to the fact that class-2 still sees Poisson arrivals, the same sort of probabilistic arguments we used for Model-A will hold. Moreover, we will see that the resulting equation when plugging this combination of parameters in our fundamental equation (Equation (242)) yields a *first-order linear ODE* in  $x$  and,  $P_y(y)$  can be taken out of the integral, saving us from having to solve an equation of the Volterra family.

**Theorem 3.** *Consider a queueing system with 2 priority classes with Poisson arrival rates  $\lambda_1$  and  $\lambda_2$ , where class-1 customers have non-preemptive priority over those of class-2, exponentially distributed service times of rate  $\mu$  independently of the class, and per-customer abandonment of class-1 customers from the queue at rate  $\theta_1$  (with no class-2 abandonment and no jockeying). Under the stability condition*

$$\rho_2 \cdot {}_1F_1\left(1; \frac{\mu}{\theta_1} + 1; \frac{\lambda_1}{\theta_1}\right) < 1 \iff \lambda_2 \mathbb{E}[B_C] < 1, \quad (238)$$

the probability generating function  $P(x, y)$  of such a system is given by

$$P(x, y) = \frac{e^{\lambda_1 x / \theta_1} \mu \pi(0, 0) (1 - y)}{\theta_1 x^{\mu / \theta_1} (1 - x)^{\lambda_2 (1 - y) / \theta_1} (\tilde{B}_C(\lambda_2 (1 - y)) - y)} \times \left[ \tilde{B}_C(\lambda_2 (1 - y)) \mathcal{H}_1(x, y) - (1 - \tilde{B}_C(\lambda_2 (1 - y))) \mathcal{H}_2(x, y) \right], \quad (239)$$

where

$$\begin{aligned} \mathcal{H}_1(x, y) &= \int_0^x e^{-\lambda_1 t / \theta_1} t^{\mu / \theta_1 - 1} (1 - t)^{\lambda_2 (1 - y) / \theta_1} dt, \\ \mathcal{H}_2(x, y) &= \int_0^x e^{-\lambda_1 t / \theta_1} t^{\mu / \theta_1} (1 - t)^{\lambda_2 (1 - y) / \theta_1 - 1} dt, \end{aligned}$$

$\tilde{B}_C(s)$  is the LST of the busy period of the  $M/M/1+M$  subsystem of class-1 given by Equation (6), and  $\pi(0, 0)$  is given by Corollary 5 below.

*Proof.* The proof is structured in three stages.

*Stage 1: Reduction to a first-order ODE in  $x$  and the integrating factor.* Substituting  $\theta_1 > 0$ ,  $\gamma_1 = \gamma_2 = \theta_2 = 0$  into the general fundamental equation eliminates both convection terms in  $y$ , leaving a first-order linear ODE in  $x$  for each fixed  $y \in [0, 1]$  (Equation (242)). Partial-fraction decomposition of the ODE coefficient gives the integrating factor

$$\mu_I(x; y) = e^{-\lambda_1 x / \theta_1} x^\alpha (1 - x)^{\beta(y)}, \quad \alpha = \frac{\mu}{\theta_1}, \quad \beta(y) = \frac{\lambda_2 (1 - y)}{\theta_1},$$

which vanishes at  $x = 0$  and  $x = 1$  because  $\alpha, \beta(y) > 0$ .

*Stage 2: Determination of  $P_y(y)$ .* Two independent routes are given in §§9.2–9.3:



- *Probabilistic (Pollaczek–Khinchine)*. Because  $\theta_2 = \gamma_i = 0$ , the class-2 arrival stream remains Poisson; PASTA applies. Class-2 customers see an  $M/G/1$  queue with effective service time  $B_C$  (the busy period of the class-1  $M/M/1+M$  subsystem), and the P-K formula yields  $P_y(y)$  directly (Equation (249)).
- *Analytical (boundedness at  $x = 1$ )*. Since  $\mu_I(1; y) = 0$ , the only way for  $P(x, y)$  to remain bounded as  $x \rightarrow 1$  is  $\int_0^1 q(t; y) \mu_I(t; y) dt = 0$  (Equation (259)). Evaluating this integral via the Kummer–Euler identity (266) recovers the same  $P_y(y)$ .

*Stage 3: Recovery of  $P(x, y)$ .* With  $P_y(y)$  known, the ODE is integrated against the integrating factor (§9.4), yielding the closed form (273) stated in the theorem.  $\square$

**Corollary 5.** *The limiting probability  $\pi(0, 0)$  of the joint state  $(N_1, N_2) = (0, 0)$  with the server busy, and the empty-system probability  $\pi_0$ , are given by*

$$\pi(0, 0) = \frac{(\lambda_1 + \lambda_2)(1 - \lambda_2 \mathbb{E}[B_C])}{\mu(1 + \lambda_1 \mathbb{E}[B_C])} = (\rho_1 + \rho_2) \frac{1 - \lambda_2 \mathbb{E}[B_C]}{1 + \lambda_1 \mathbb{E}[B_C]}, \quad (240)$$

$$\pi_0 = \frac{1 - \lambda_2 \mathbb{E}[B_C]}{1 + \lambda_1 \mathbb{E}[B_C]}, \quad (241)$$

where  $\mathbb{E}[B_C]$  is the mean busy period of the  $M/M/1+M$  subsystem of class-1, given by Equation (7). In the limit  $\theta_1 \rightarrow 0^+$ ,  $\mathbb{E}[B_C] \rightarrow (\mu - \lambda_1)^{-1}$  and the expressions above recover the classical Model-A results  $\pi(0, 0) = \rho(1 - \rho)$  and  $\pi_0 = 1 - \rho$ .

## 9.1 Setup: reduction to a first-order linear ODE in $x$

Plugging  $\theta_1 > 0$ ,  $\gamma_1 = \gamma_2 = \theta_2 = 0$  into the general fundamental equation (45) (Lemma 1) annihilates the  $\partial P / \partial y$  term entirely, since its coefficient  $xy[\gamma_2(y-x) + \theta_2(y-1)]$  vanishes identically, and reduces the  $\partial P / \partial x$  coefficient  $xy[\gamma_1(x-y) + \theta_1(x-1)]$  to its single surviving abandonment term  $\theta_1 xy(x-1)$ . What remains is

$$[(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 xy^2] P(x, y) + \theta_1 xy(x-1) \frac{\partial}{\partial x} P(x, y) = \mu(x-y) P_y(y) - \mu x(1-y) \pi(0, 0), \quad (242)$$

where the boundary function  $P_y(y) = P(0, y) = \sum_{n_2=0}^{\infty} \pi(0, n_2) y^{n_2}$  is, as in Model-A, the partial PGF restricted to states where the server is busy and the priority queue is empty ( $N_1 = 0$ ).

For any fixed  $y$ , Equation (242) is a *first-order linear ODE in  $x$*  whose unknowns are the function  $P(x, y)$  on  $[0, 1]^2$  and the boundary trace  $P_y(y) = P(0, y)$ . The strategy in what follows is in two stages:

1. Identify  $P_y(y)$  — first by a probabilistic argument exploiting the  $M/G/1$  structure seen by class-2 (§9.2), then re-derived analytically from boundedness of  $P(x, y)$  at  $x = 1$  (§9.3).
2. Recover  $P(x, y)$  by solving the ODE with  $P_y(y)$  now known (§9.4).

We close the setup with the conditional-expectation decomposition that will drive the probabilistic derivation. By the law of total expectation, restricting the defining sum of  $P_y(y) = \mathbb{E}[\mathbf{1}\{\text{busy}, N_1 = 0\} y^{N_2}]$  to the event  $\{\text{busy}, N_1 = 0\}$  factors it into the mass of that event and the conditional PGF of  $N_2$  given it:

$$P_y(y) = \mathbb{P}(\text{busy}, N_1 = 0) \cdot \mathbb{E}[y^{N_2} \mid \text{busy}, N_1 = 0]. \quad (243)$$

## 9.2 Probabilistic derivation of $P_y(y)$

The idea is, again, to adopt the perspective of a class-2 customer waiting in line while the server is busy. Unlike in the case of jockeying, the customers from class-1 truly abandon the system, leaving the Poisson arrival process of class-2 undisturbed. Because class-1 holds non-preemptive priority (§3.2), a waiting class-2 customer is not taken into service while any class-1 customer is present ( $N_1 > 0$ ); its *effective service time* is therefore the duration of one complete busy period  $B_C$  of the class-1 subsystem—now an  $M/M/1 + M$  queue with arrival rate  $\lambda_1$ , service rate  $\mu$  and per-customer abandonment rate  $\theta_1$ , rather than the pure  $M/M/1$  of Model-A. By the *memorylessness* of the exponential service, arrival and abandonment clocks, these effective service times are i.i.d., so from class-2’s viewpoint the system is an  $M/G/1$  queue with Poisson arrivals at rate  $\lambda_2$  and general service times distributed as  $B_C$ .

The LST  $\tilde{B}_C(s)$  and mean  $\mathbb{E}[B_C]$  of this busy period are derived via the first-step analysis in §3.3; see Equations (6)–(7).

The stability condition required for the class-2 subsystem is that of the  $M/G/1$  defined above, i.e.  $\lambda_2 \mathbb{E}[B_C] < 1$ :

$$\rho_2 \cdot {}_1F_1\left(1; \frac{\mu}{\theta_1} + 1; \frac{\lambda_1}{\theta_1}\right) < 1. \quad (244)$$

This condition is also necessary, not merely sufficient. The quantity  $\lambda_2 \mathbb{E}[B_C]$  is precisely the utilisation of the embedded  $M/G/1$  queue of class-2 customers (Poisson input  $\lambda_2$ , effective service time  $B_C$ ), so when  $\lambda_2 \mathbb{E}[B_C] \geq 1$  that  $M/G/1$  is null recurrent (at equality) or transient (strictly above) and its number-in-system has no proper stationary law. Because the class-2 population is a deterministic function of the joint state  $(n_1, n_2)$ , a divergent class-2 queue rules out positive recurrence of the joint chain, so (244) is sharp. We argue this at the level of the embedded  $M/G/1$ , whose marginal dynamics are exact, rather than by constructing a Lyapunov function on the joint generator. Compared to Model-A, this stability condition is strictly weaker than the classical  $\rho_1 + \rho_2 < 1$ , which aligns with the fact that abandonments lower the load of the system: for all  $\theta_1 > 0$  the series is termwise dominated by the geometric series for  $1/(1 - \rho_1)$ . Indeed, since  $1^{(n)} = n!$  the Kummer series (13) reduces to

$${}_1F_1\left(1; \frac{\mu}{\theta_1} + 1; \frac{\lambda_1}{\theta_1}\right) = \sum_{n=0}^{\infty} \frac{(\lambda_1/\theta_1)^n}{\left(\frac{\mu}{\theta_1} + 1\right)^{(n)}} = \sum_{n=0}^{\infty} \frac{(\lambda_1/\theta_1)^n}{\prod_{k=1}^n \left(\frac{\mu}{\theta_1} + k\right)},$$

and since  $\frac{\mu}{\theta_1} + k > \frac{\mu}{\theta_1}$  for every  $k \geq 1$ , the  $n$ -th term is bounded above by  $(\lambda_1/\theta_1)^n / (\mu/\theta_1)^n = \rho_1^n$ , with strict inequality for  $n \geq 1$ . Summing gives  ${}_1F_1(1; \mu/\theta_1 + 1; \lambda_1/\theta_1) < \sum_{n \geq 0} \rho_1^n = 1/(1 - \rho_1)$ . Abandonments in the priority queue therefore stabilise the non-priority queue.

The class-2 subsystem is therefore an  $M/G/1$  queue with Poisson arrival rate  $\lambda_2$ , service-time LST  $\tilde{B}_C(s)$  and utilisation  $\rho_B = \lambda_2 \mathbb{E}[B_C]$ . Applying the Pollaczek–Khinchine formula (12) (§3.4) with these identifications ( $\lambda \mapsto \lambda_2$ ,  $\tilde{B} \mapsto \tilde{B}_C$ ,  $\rho_B = \lambda_2 \mathbb{E}[B_C]$ ) gives the PGF of the stationary number of class-2 customers in that  $M/G/1$ :

$$L_2(y) = \frac{(1 - \lambda_2 \mathbb{E}[B_C]) (1 - y) \tilde{B}_C(\lambda_2(1 - y))}{\tilde{B}_C(\lambda_2(1 - y)) - y}. \quad (245)$$

Because  $\theta_2 = \gamma_i = 0$ , the class-2 arrival stream is Poisson and satisfies the lack-of-anticipation assumption, so by the *PASTA* property (§3.2) the time-stationary distribution of the number of class-2 customers present coincides with the  $M/G/1$  steady-state distribution. Conditioning on  $\{\text{busy}, N_1 = 0\}$  — the event on which a class-2 customer is at the head of its queue — therefore identifies that count with  $N_2$ , giving

$$\mathbb{E}[y^{N_2} \mid \text{busy}, N_1 = 0] = L_2(y). \quad (246)$$

Substituting the conditional identification (246) into the law-of-total-expectation decomposition (243) yields

$$P_y(y) = \mathbb{P}(\text{busy}, N_1 = 0) \left[ \frac{(1 - \lambda_2 \mathbb{E}[B_C]) (1 - y) \tilde{B}_C(\lambda_2(1 - y))}{\tilde{B}_C(\lambda_2(1 - y)) - y} \right]. \quad (247)$$

By the definition of  $P_y$ ,  $P_y(0) = \pi(0, 0)$ . Evaluating (247) at  $y = 0$ ,

$$\begin{aligned} P_y(0) &= \pi(0, 0) = \mathbb{P}(\text{busy}, N_1 = 0) \left[ \frac{(1 - \lambda_2 \mathbb{E}[B_C]) \cdot 1 \cdot \tilde{B}_C(\lambda_2)}{\tilde{B}_C(\lambda_2)} \right] \\ &= \mathbb{P}(\text{busy}, N_1 = 0) (1 - \lambda_2 \mathbb{E}[B_C]), \end{aligned} \quad (248)$$

so that  $\mathbb{P}(\text{busy}, N_1 = 0) = \pi(0, 0)/(1 - \lambda_2 \mathbb{E}[B_C])$ . Substituting this back into (247), the  $(1 - \lambda_2 \mathbb{E}[B_C])$  factor cancels and we are left with

$$P_y(y) = \pi(0, 0) \frac{(1 - y) \tilde{B}_C(\lambda_2(1 - y))}{\tilde{B}_C(\lambda_2(1 - y)) - y}. \quad (249)$$

This is structurally the same expression as in the probabilistic argument for Model-A (Equation (89)), with the only difference being the effective service time:  $B_C$  is now the busy period of an  $M/M/1 + M$  subsystem with abandonment rate  $\theta_1$ , rather than the busy period of a pure  $M/M/1$ .



### 9.3 Analytical re-derivation of $P_y(y)$ via boundedness at $x = 1$

Here the surviving abandonment factor  $(x - 1)$  places the operative singularity on the boundary  $x = 1$ , exactly the boundedness-pinning regime anticipated in the remark following Lemma 1. For an analytic counterpart of the previous derivation, we put (242) in standard form:

$$\frac{\partial P}{\partial x} + \underbrace{\frac{(\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy}{\theta_1 x(x-1)}}_{p(x; y)} P = \underbrace{\frac{\mu(x-y) P_y(y) - \mu x(1-y) \pi(0, 0)}{\theta_1 xy(x-1)}}_{q(x; y)}. \quad (250)$$

The numerator of  $p(x; y)$  is the same polynomial we have seen in every model so far. With  $x^*(y)$  and  $x^+(y)$  as its roots (Equation (87)), we can write

$$(\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy = -\lambda_1 (x - x^*(y))(x - x^+(y)), \quad (251)$$

with  $x^*(y) x^+(y) = \mu/\lambda_1$ . Partial fraction decomposition gives

$$\frac{-\lambda_1 (x - x^*(y))(x - x^+(y))}{x(x-1)} = A + \frac{B}{x} + \frac{C}{x-1}, \quad (252)$$

where  $A = -\lambda_1$  is read off from the ratio of leading coefficients of the quadratic terms in numerator and denominator,  $B = \mu$  from setting  $x = 0$  and using  $\lambda_1 x^* x^+ = \mu$ , and  $C = \lambda_2(1-y)$  by the cover-up rule at  $x = 1$ . Writing the numerator as  $N(x) = (\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy$ , the coefficient of  $1/(x-1)$  is its residue  $C = N(1) = (\lambda_1 + \lambda_2 + \mu) - \mu - \lambda_1 - \lambda_2 y = \lambda_2(1-y)$ ; equivalently, since  $N(x) = -\lambda_1 (x - x^*(y))(x - x^+(y))$ , this is the identity  $(1 - x^*(y))(1 - x^+(y)) = -\lambda_2(1-y)/\lambda_1$  evaluated at  $x = 1$ . Thus

$$p(x; y) = \frac{1}{\theta_1} \left[ -\lambda_1 + \frac{\mu}{x} + \frac{\lambda_2(1-y)}{x-1} \right]. \quad (253)$$

Integrating term by term,

$$\int p(x; y) dx = \frac{1}{\theta_1} [-\lambda_1 x + \mu \ln x + \lambda_2(1-y) \ln(1-x)], \quad (254)$$

which yields the integrating factor

$$\mu_I(x; y) = e^{-\lambda_1 x/\theta_1} \cdot x^\alpha \cdot (1-x)^{\beta(y)}, \quad (255)$$

where, in parallel with Model- $B_2$ , we set

$$\alpha = \frac{\mu}{\theta_1}, \quad \beta(y) = \frac{\lambda_2(1-y)}{\theta_1}. \quad (256)$$

Under our stability assumptions both exponents are positive ( $\alpha > 0$  always;  $\beta(y) \geq 0$  for  $y \in [0, 1]$ ), so  $\mu_I$  vanishes at both  $x = 0$  and  $x = 1$ . From  $\frac{d}{dx}[P \cdot \mu_I] = q \cdot \mu_I$ , integration over  $(0, x]$  reads

$$P(x, y) \mu_I(x; y) - P(0, y) \mu_I(0; y) = \int_0^x q(t; y) \mu_I(t; y) dt, \quad (257)$$

and the boundary term vanishes, yielding

$$P(x, y) = \frac{1}{\mu_I(x; y)} \int_0^x q(t; y) \mu_I(t; y) dt. \quad (258)$$

Now, in the same fashion as the root-cancellation argument used for Model- $A$ , we exploit boundedness of PGFs to determine  $P_y(y)$  by observing the behaviour of  $P(x, y)$  as  $x \rightarrow 1$ , for fixed  $y \in [0, 1]$ : there  $\beta(y) > 0$ , so  $\mu_I(x; y) \rightarrow 0$  as  $x \rightarrow 1$ . (At the endpoint  $y = 1$  the exponent  $\beta(1) = 0$  and  $\mu_I(\cdot; 1)$  no longer vanishes at  $x = 1$ ; the value  $P_y(1)$  is recovered from the resulting formula by continuity,  $y \rightarrow 1^-$ .) Since  $\mu_I(x; y) \rightarrow 0$ , the only way to keep the right-hand side of (258) bounded is

$$\int_0^1 q(t; y) \mu_I(t; y) dt = 0. \quad (259)$$

Substituting the explicit forms of  $q$  and  $\mu_I$  into (259), the integrand is

$$q(t; y) \mu_I(t; y) = \frac{\mu[(t-y) P_y(y) - t(1-y) \pi(0, 0)]}{\theta_1 y} \cdot \frac{e^{-\lambda_1 t/\theta_1} t^\alpha (1-t)^{\beta(y)}}{t(t-1)}, \quad (260)$$

where  $t^\alpha/t = t^{\alpha-1}$  and, with  $b = \beta(y)$ , the elementary identity  $(1-t)^{\beta(y)}/(t-1) = -(1-t)^{\beta(y)-1}$  flips the sign and lowers the exponent. The constant prefactor  $-\mu/(\theta_1 y)$ , nonzero for  $y \in (0, 1]$ , divides out of the vanishing condition (259), leaving

$$\int_0^1 e^{-\lambda_1 t/\theta_1} t^{\alpha-1} (1-t)^{\beta(y)-1} [(t-y) P_y(y) - t(1-y) \pi(0,0)] dt = 0, \quad (261)$$

with  $\alpha$  and  $\beta(y)$  as in (256). Introducing the auxiliary integrals

$$\mathcal{J}_0(y) = \int_0^1 e^{-\lambda_1 t/\theta_1} t^{\alpha-1} (1-t)^{\beta(y)-1} dt, \quad (262)$$

$$\mathcal{J}_1(y) = \int_0^1 e^{-\lambda_1 t/\theta_1} t^\alpha (1-t)^{\beta(y)-1} dt, \quad (263)$$

and distributing the bracket over  $t^{\alpha-1}$ , note that  $t \cdot t^{\alpha-1} = t^\alpha$  feeds  $\mathcal{J}_1$  while  $t^{\alpha-1}$  feeds  $\mathcal{J}_0$ . The two terms regroup as

$$P_y(y) \int_0^1 e^{-\lambda_1 t/\theta_1} (t^\alpha - y t^{\alpha-1}) (1-t)^{\beta(y)-1} dt - (1-y) \pi(0,0) \int_0^1 e^{-\lambda_1 t/\theta_1} t^\alpha (1-t)^{\beta(y)-1} dt = 0,$$

so the boundedness condition (259) becomes

$$P_y(y) [\mathcal{J}_1(y) - y \mathcal{J}_0(y)] = (1-y) \pi(0,0) \mathcal{J}_1(y), \quad (264)$$

which gives

$$P_y(y) = \pi(0,0) (1-y) \frac{\mathcal{J}_1(y)}{\mathcal{J}_1(y) - y \mathcal{J}_0(y)}. \quad (265)$$

The continued-fraction representation (6) admits an equivalent integral form. Setting  $a = \alpha = \mu/\theta_1$ ,  $b = s/\theta_1$ ,  $c = \lambda_1/\theta_1$ , the Euler integral representation (15) identifies each of the integrals below as a Beta-function multiple of a  ${}_1F_1$  value:

$$\tilde{B}_C(s) = \frac{\int_0^1 t^a (1-t)^{b-1} e^{-ct} dt}{\int_0^1 t^{a-1} (1-t)^{b-1} e^{-ct} dt} = \frac{B(a+1, b) {}_1F_1(a+1; a+1+b; -c)}{B(a, b) {}_1F_1(a; a+b; -c)}. \quad (266)$$

That this ratio equals the continued fraction (6) is exactly the closed-form summation (11) proved in §3.3: the integrals satisfy the three-term recurrence (10), they form its minimal solution, and Pincherle's theorem identifies the convergent continued fraction with the minimal-solution ratio.

Setting  $s = \lambda_2(1-y)$  — so that  $b = \beta(y)$  matches the exponent in (256) — yields exactly the ratio of our  $\mathcal{J}_i$ :

$$\tilde{B}_C(\lambda_2(1-y)) = \frac{\mathcal{J}_1(y)}{\mathcal{J}_0(y)}. \quad (267)$$

Dividing numerator and denominator of (265) by  $\mathcal{J}_0(y)$  then recovers exactly Equation (249). The boundedness route, which uses no probabilistic structure, reaches the same Pollaczek–Khinchine form only through the Beta-integral identity (266), which is the analytic shadow of the  $M/G/1$ /busy-period decomposition used in §9.2.

## 9.4 Recovery of $P(x, y)$ from $P_y(y)$

With  $P_y(y)$  now in hand, we return to the ODE (242) and substitute (249) into its right-hand side. Writing  $\zeta(y) \equiv \tilde{B}_C(\lambda_2(1-y))$  for brevity,

$$\begin{aligned} \text{RHS} &= \mu(x-y) \pi(0,0) \frac{(1-y) \zeta(y)}{\zeta(y) - y} - \mu x(1-y) \pi(0,0) \\ &= \mu \pi(0,0) (1-y) \left[ \frac{(x-y) \zeta(y)}{\zeta(y) - y} - x \right] \\ &= \mu \pi(0,0) (1-y) \frac{(x-y) \zeta(y) - x[\zeta(y) - y]}{\zeta(y) - y} \\ &= \mu \pi(0,0) y(1-y) \frac{x - \zeta(y)}{\zeta(y) - y}, \end{aligned}$$

which is structurally identical to the RHS found for Model-A, though now with the more involved effective service time  $B_C$ . With  $p(x; y)$  and  $\mu_I(x; y)$  as in (253) and (255), the source  $q(x; y)$  of the standard-form ODE now reads

$$q(x; y) = \frac{\mu \pi(0, 0) (1 - y) [x - \zeta(y)]}{\theta_1 x(x - 1) [\zeta(y) - y]}, \quad (268)$$

and the implicit solution (258) becomes

$$P(x, y) = \frac{1}{\mu_I(x; y)} \int_0^x q(t; y) \mu_I(t; y) dt. \quad (269)$$

To evaluate the integral, we use the partial-fraction identity

$$\frac{t - \zeta(y)}{t(t - 1)} = \frac{\zeta(y)(t - 1) + t(1 - \zeta(y))}{t(t - 1)} = \frac{\zeta(y)}{t} + \frac{1 - \zeta(y)}{t - 1},$$

and define the auxiliary integrals

$$\mathcal{H}_1(x, y) = \int_0^x e^{-\lambda_1 t / \theta_1} t^{\alpha-1} (1 - t)^{\beta(y)} dt, \quad (270)$$

$$\mathcal{H}_2(x, y) = \int_0^x e^{-\lambda_1 t / \theta_1} t^{\alpha} (1 - t)^{\beta(y)-1} dt, \quad (271)$$

with  $\alpha$  and  $\beta(y)$  as in (256).

*Note on the  $\mathcal{J}$ - $\mathcal{H}$  correspondence.* The integrals  $\mathcal{H}_1, \mathcal{H}_2$  are the partial (upper limit  $x$ ) counterparts of the full integrals  $\mathcal{J}_0, \mathcal{J}_1$  from (259). Evaluating at  $x = 1$  gives  $\mathcal{H}_2(1, y) = \mathcal{J}_1(y)$  exactly. However,  $\mathcal{H}_1(1, y)$  is *not*  $\mathcal{J}_0(y)$ : the former has the exponent  $\beta(y)$  on  $(1 - t)$  while the latter has  $\beta(y) - 1$ , which under the Euler representation (15) corresponds to incrementing the second Kummer parameter by one. Both families are evaluated by the same representation; the exponent shift is harmless because  $\mathcal{H}_1$  and  $\mathcal{H}_2$  always appear together in the combination  $\zeta(y) \mathcal{H}_1 - (1 - \zeta(y)) \mathcal{H}_2$ , not individually.

Multiplying the source  $q(t; y)$  by the integrating factor  $\mu_I(t; y) = e^{-\lambda_1 t / \theta_1} t^{\alpha} (1 - t)^{\beta(y)}$  and inserting the partial fraction, the integrand splits into the two pieces defining  $\mathcal{H}_1, \mathcal{H}_2$ :

$$\begin{aligned} q(t; y) \mu_I(t; y) &= \frac{\mu \pi(0, 0) (1 - y)}{\theta_1 (\zeta(y) - y)} \left[ \frac{\zeta(y)}{t} + \frac{1 - \zeta(y)}{t - 1} \right] e^{-\lambda_1 t / \theta_1} t^{\alpha} (1 - t)^{\beta(y)} \\ &= \frac{\mu \pi(0, 0) (1 - y)}{\theta_1 (\zeta(y) - y)} \left[ \zeta(y) e^{-\lambda_1 t / \theta_1} t^{\alpha-1} (1 - t)^{\beta(y)} - (1 - \zeta(y)) e^{-\lambda_1 t / \theta_1} t^{\alpha} (1 - t)^{\beta(y)-1} \right], \end{aligned}$$

where the second term used  $(1 - t)^{\beta(y)} / (t - 1) = -(1 - t)^{\beta(y)-1}$ , the same identity used to simplify the boundedness condition (259); this is the source of both the minus sign and the exponent shift  $\beta(y) \rightarrow \beta(y) - 1$  on  $\mathcal{H}_2$ . Since  $\pi(0, 0)$  and  $\zeta(y)$  come out of the integral,

$$\int_0^x q(t; y) \mu_I(t; y) dt = \frac{\mu \pi(0, 0) (1 - y)}{\theta_1 (\zeta(y) - y)} \left[ \zeta(y) \mathcal{H}_1(x, y) - (1 - \zeta(y)) \mathcal{H}_2(x, y) \right]. \quad (272)$$

Dividing by  $\mu_I(x; y)$  and restoring  $\zeta(y) = \tilde{B}_C(\lambda_2(1 - y))$ , we arrive at the closed form

$$\begin{aligned} P(x, y) &= \frac{e^{\lambda_1 x / \theta_1} \mu \pi(0, 0) (1 - y)}{\theta_1 x^{\alpha} (1 - x)^{\beta(y)} (\tilde{B}_C(\lambda_2(1 - y)) - y)} \\ &\quad \times \left[ \tilde{B}_C(\lambda_2(1 - y)) \mathcal{H}_1(x, y) - (1 - \tilde{B}_C(\lambda_2(1 - y))) \mathcal{H}_2(x, y) \right], \end{aligned} \quad (273)$$

which is the statement of the theorem. This closed form is the product of a probabilistic derivation of  $P_y(y)$  via Pollaczek–Khinchine, combined with an analytic integration of the ODE in  $x$ ; the same result is reached from the purely analytic route of §9.3.

*Proof of Corollary 5.* The fully idle state—the server empty with no customer present, of probability  $\pi_0$ —communicates with the rest of the chain through a single neighbour: the busy-and-empty state  $(N_1, N_2) = (0, 0)$  of probability  $\pi(0, 0)$ , in which exactly one customer is in service and both queues are empty (recall the standing convention  $\pi_0 \neq \pi(0, 0)$ ). The system leaves the idle state only upon an arrival of either class, at combined Poisson rate  $\lambda_1 + \lambda_2$  (Poisson superposition), and

re-enters it only upon the service completion that empties the busy-and-empty state, at rate  $\mu$ ; no abandonment can fire from  $\pi(0, 0)$ , since both queues are empty. Equating the probability flow across the cut separating the idle state from its single neighbour (global balance for the idle state, as in (32)) therefore gives

$$(\lambda_1 + \lambda_2) \pi_0 = \mu \pi(0, 0), \quad (274)$$

so  $\pi(0, 0) = (\rho_1 + \rho_2) \pi_0$ .

To fix  $\pi_0$ , we exploit the  $M/G/1$  structure used in §9.2. Class-2 customers form an  $M/G/1$  queue with arrival rate  $\lambda_2$  and effective service time  $B_C$ . By the standard  $M/G/1$  work-conservation (utilisation) identity, the long-run fraction of time this  $M/G/1$  is busy is  $\rho_B = \lambda_2 \mathbb{E}[B_C]$ , so its idle probability is  $1 - \rho_B$ . The event “this  $M/G/1$  is idle” coincides with the absence of any class-2 customer from the system, so

$$\mathbb{P}(\text{no class-2 present}) = 1 - \lambda_2 \mathbb{E}[B_C]. \quad (275)$$

Conditional on no class-2 being present, the priority subsystem evolves freely as the same  $M/M/1 + M$  queue used to define  $B_C$ , with arrival rate  $\lambda_1$ , service rate  $\mu$  and abandonment rate  $\theta_1$ . This subsystem alternates between idle periods (no class-1 customer present) and busy periods  $B_C$ , the successive idle+busy spans forming an alternating renewal process whose regeneration epochs are the instants at which the subsystem empties. An idle period lasts until the next class-1 arrival; by the memorylessness of the exponential interarrival clock this duration is  $\text{Exp}(\lambda_1)$ , with mean  $1/\lambda_1$ . The mean regeneration cycle is therefore  $1/\lambda_1 + \mathbb{E}[B_C]$ , and by the renewal–reward theorem the stationary fraction of time the subsystem is idle equals the ratio of the mean idle period to the mean cycle:

$$\mathbb{P}(\text{class-1 absent} \mid \text{no class-2 present}) = \frac{1/\lambda_1}{1/\lambda_1 + \mathbb{E}[B_C]} = \frac{1}{1 + \lambda_1 \mathbb{E}[B_C]}. \quad (276)$$

The fully idle event  $\{\text{server idle}, N_1 = N_2 = 0\}$  is exactly the intersection  $A \cap B$  of the events  $B = \{\text{no class-2 present}\}$  and  $A = \{\text{class-1 absent}\}$ , since the server is idle precisely when no customer of either class is in the system. By the multiplication (chain) rule  $\mathbb{P}(A \cap B) = \mathbb{P}(B) \mathbb{P}(A \mid B)$ —using the conditional idle fraction just computed for the factor  $\mathbb{P}(A \mid B)$ , and *not* independence of  $A$  and  $B$ —we obtain

$$\pi_0 = (1 - \lambda_2 \mathbb{E}[B_C]) \cdot \frac{1}{1 + \lambda_1 \mathbb{E}[B_C]} = \frac{1 - \lambda_2 \mathbb{E}[B_C]}{1 + \lambda_1 \mathbb{E}[B_C]}, \quad (277)$$

which is (241). Substituting into (274) yields (240).

For the Model-A limit,  $\theta_1 \rightarrow 0^+$  gives  $\mathbb{E}[B_C] \rightarrow (\mu - \lambda_1)^{-1}$  (the busy period of the  $M/M/1 + M$  subsystem collapses to that of the pure  $M/M/1$  priority subsystem). Then

$$\lambda_1 \mathbb{E}[B_C] \rightarrow \frac{\lambda_1}{\mu - \lambda_1} = \frac{\rho_1}{1 - \rho_1}, \quad \lambda_2 \mathbb{E}[B_C] \rightarrow \frac{\lambda_2}{\mu - \lambda_1} = \frac{\rho_2}{1 - \rho_1},$$

so that the denominator  $1 + \lambda_1 \mathbb{E}[B_C] \rightarrow 1/(1 - \rho_1)$  and the numerator

$$1 - \lambda_2 \mathbb{E}[B_C] \rightarrow 1 - \frac{\rho_2}{1 - \rho_1} = \frac{1 - \rho_1 - \rho_2}{1 - \rho_1} = \frac{1 - \rho}{1 - \rho_1}.$$

Substituting into (241),

$$\pi_0 \rightarrow \frac{(1 - \rho)/(1 - \rho_1)}{1/(1 - \rho_1)} = 1 - \rho,$$

and then  $\pi(0, 0) = (\rho_1 + \rho_2) \pi_0 \rightarrow \rho(1 - \rho)$  via (274), recovering the classical Model-A baseline values.  $\square$

## 9.5 Mean queue lengths and mean waiting times

### Why the diagonal no longer simplifies

Setting  $x = y = z$  in the fundamental equation (242), the abandonment convection  $\theta_1 xy(x - 1) \partial_x P$  does *not* vanish (since  $(x - 1)|_{x=z} \neq 0$  for  $z \in (0, 1)$ ). The diagonal equation reads:

$$\begin{aligned} & [(\lambda_1 + \lambda_2 + \mu)z^2 - \mu z - \lambda z^3] P(z, z) + \theta_1 z^2 (z - 1) \frac{\partial P}{\partial x}(z, z) \\ & = -\mu z(1 - z) \pi(0, 0), \end{aligned} \quad (278)$$

Comparison with Model-A is instructive. There the same diagonal substitution  $x = y = z$  also annihilates the boundary term  $\mu(x - y) P_y(y)$ , but with  $\theta_1 = \gamma_1 = \gamma_2 = 0$  there is *no* convection term at all, so the diagonal equation reduced to a purely algebraic relation in the single unknown  $P(z, z)$ , with closed form  $P(z, z) = \pi(0, 0)/(1 - \rho z)$  (Equation (120)), from which  $\mathbb{E}[N_1] + \mathbb{E}[N_2]$  followed by a single differentiation. Here the surviving convection  $\theta_1 z^2(z - 1) \partial_x P(z, z)$  injects the off-diagonal slope  $\partial_x P(z, z)$  as a *second* unknown into the lone diagonal relation, and no companion equation closes the system; consequently no closed form for  $P(z, z)$ —and hence for the total  $\mathbb{E}[N_1] + \mathbb{E}[N_2]$ —is available. Abandonment removes customers and reduces both means relative to Model-A.

### Class-2 mean queue length via $P(1, y)$

Setting  $x = 1$  in the fundamental equation (242), the abandonment term  $\theta_1 xy(x - 1) \partial_x P$  vanishes identically, and the coefficient of  $P(1, y)$  simplifies to  $\lambda_2 y(1 - y)$ :

$$\lambda_2 y(1 - y) P(1, y) = \mu(1 - y) [P_y(y) - \pi(0, 0)]. \quad (279)$$

Cancelling  $(1 - y)$  for  $y \neq 1$  and substituting the known  $P_y(y)$  (249):

$$P(1, y) = \frac{\mu [P_y(y) - \pi(0, 0)]}{\lambda_2 y} = \frac{\mu \pi(0, 0)}{\lambda_2} \cdot \frac{1 - \tilde{B}_C(\lambda_2(1 - y))}{\tilde{B}_C(\lambda_2(1 - y)) - y}. \quad (280)$$

This is the marginal PGF of  $N_2$ . To differentiate at  $y = 1$  (where the expression is 0/0), set  $\varepsilon = 1 - y \rightarrow 0$  and use the LST expansion  $\tilde{B}_C(\lambda_2 \varepsilon) = 1 - \lambda_2 \mathbb{E}[B_C] \varepsilon + \frac{1}{2} \lambda_2^2 \mathbb{E}[B_C^2] \varepsilon^2 + O(\varepsilon^3)$ :

$$\begin{aligned} 1 - \tilde{B}_C(\lambda_2 \varepsilon) &= \lambda_2 \mathbb{E}[B_C] \varepsilon - \frac{1}{2} \lambda_2^2 \mathbb{E}[B_C^2] \varepsilon^2 + O(\varepsilon^3), \\ \tilde{B}_C(\lambda_2 \varepsilon) - y &= (1 - \lambda_2 \mathbb{E}[B_C] \varepsilon + \frac{1}{2} \lambda_2^2 \mathbb{E}[B_C^2] \varepsilon^2) - (1 - \varepsilon) \\ &= (1 - \lambda_2 \mathbb{E}[B_C]) \varepsilon + \frac{1}{2} \lambda_2^2 \mathbb{E}[B_C^2] \varepsilon^2 + O(\varepsilon^3), \end{aligned} \quad (281)$$

so that (writing  $a = \lambda_2 \mathbb{E}[B_C]$  and  $c = \frac{1}{2} \lambda_2^2 \mathbb{E}[B_C^2]$ ):

$$P(1, 1 - \varepsilon) = \frac{\mu \pi(0, 0)}{\lambda_2} \cdot \frac{a \varepsilon - c \varepsilon^2 + O(\varepsilon^3)}{(1 - a) \varepsilon + c \varepsilon^2 + O(\varepsilon^3)} = \frac{\mu \pi(0, 0)}{\lambda_2} \cdot \frac{a - c \varepsilon}{(1 - a) + c \varepsilon} + O(\varepsilon^2). \quad (282)$$

Differentiating  $f(\varepsilon) = (a - c \varepsilon)/((1 - a) + c \varepsilon)$  by the quotient rule:

$$f'(\varepsilon) = \frac{-c((1 - a) + c \varepsilon) - (a - c \varepsilon) \cdot c}{((1 - a) + c \varepsilon)^2} = \frac{-c}{(1 - a + c \varepsilon)^2}, \quad (283)$$

and since  $d/dy P(1, y)|_{y=1} = -d/d\varepsilon P(1, 1 - \varepsilon)|_{\varepsilon=0}$ :

$$\mathbb{E}[N_2] = \frac{d}{dy} P(1, y) \Big|_{y=1} = \frac{\mu \pi(0, 0)}{\lambda_2} \cdot \frac{c}{(1 - a)^2} = \frac{\mu \pi(0, 0) \lambda_2 \mathbb{E}[B_C^2]}{2(1 - \lambda_2 \mathbb{E}[B_C])^2}. \quad (284)$$

As  $\theta_1 \rightarrow 0^+$ ,  $\mathbb{E}[B_C] \rightarrow (\mu - \lambda_1)^{-1}$  and  $\mathbb{E}[B_C^2] \rightarrow 2\mu/(\mu - \lambda_1)^3$ ; substituting recovers the Model-A value  $\mathbb{E}[N_2] = \rho \rho_2 / ((1 - \rho_1)(1 - \rho))$ .

### Class-1 mean queue length

We extract  $\mathbb{E}[N_1] = \frac{\partial P}{\partial x} \Big|_{(1,1)}$  by evaluating the theorem formula at  $y = 1$  (limit) and then differentiating in  $x$ .

*Step 1: marginal PGF  $P(x, 1)$ .* As  $y \rightarrow 1$ , the exponent  $\beta(y) = \lambda_2(1 - y)/\theta_1 \rightarrow 0$  so  $(1 - x)^{\beta(y)} \rightarrow 1$  for fixed  $x \in (0, 1)$ , and consequently  $\mathcal{H}_1(x, y) \rightarrow \int_0^x e^{-\lambda_1 t/\theta_1} t^{\mu/\theta_1 - 1} dt$  and  $\mathcal{H}_2(x, y)$  tends to a finite limit (its integrand  $e^{-\lambda_1 t/\theta_1} t^{\mu/\theta_1} (1 - t)^{-1}$  is integrable on  $[0, x]$  for  $x < 1$ ). Inside the theorem's bracket the two integrals carry different weights: by the LST expansion of §9.2,  $\tilde{B}_C(\lambda_2(1 - y)) \rightarrow \tilde{B}_C(0) = 1$  while  $1 - \tilde{B}_C(\lambda_2(1 - y)) = \lambda_2 \mathbb{E}[B_C] (1 - y) + O((1 - y)^2) \rightarrow 0$ , so the  $\mathcal{H}_2$  term vanishes in the limit and the bracket collapses to  $\mathcal{H}_1(x, 1)$  alone. Combining this with the scalar prefactor, whose  $y$ -dependent part satisfies  $(1 - y)/(\tilde{B}_C(\lambda_2(1 - y)) - y) \rightarrow 1/(1 - \lambda_2 \mathbb{E}[B_C])$  (again from the expansion  $\tilde{B}_C(\lambda_2(1 - y)) - y = (1 - \lambda_2 \mathbb{E}[B_C])(1 - y) + O((1 - y)^2)$ ), the theorem formula gives:

$$P(x, 1) = \frac{\mu \pi(0, 0)}{\theta_1 (1 - \lambda_2 \mathbb{E}[B_C])} \cdot e^{\lambda_1 x/\theta_1} x^{-\mu/\theta_1} \int_0^x e^{-\lambda_1 t/\theta_1} t^{\mu/\theta_1 - 1} dt. \quad (285)$$

*Step 2: differentiate  $P(x, 1)$  at  $x = 1$ .* Write  $P(x, 1) = C \cdot f(x) \cdot \mathcal{K}(x)$  with  $C = \mu\pi(0, 0)/(\theta_1(1 - \lambda_2\mathbb{E}[B_C]))$ ,  $f(x) = e^{\lambda_1 x/\theta_1} x^{-\mu/\theta_1}$ , and  $\mathcal{K}(x) = \int_0^x e^{-\lambda_1 t/\theta_1} t^{\mu/\theta_1-1} dt$ . By the product rule and Leibniz's rule ( $d\mathcal{K}/dx = e^{-\lambda_1 x/\theta_1} x^{\mu/\theta_1-1}$ ):

$$\frac{d}{dx}P(x, 1) = C \left[ f'(x) \mathcal{K}(x) + f(x) \cdot e^{-\lambda_1 x/\theta_1} x^{\mu/\theta_1-1} \right]. \quad (286)$$

The logarithmic derivative  $f'(x)/f(x) = \lambda_1/\theta_1 - (\mu/\theta_1)/x = (\lambda_1 x - \mu)/(\theta_1 x)$ , so:

$$\frac{d}{dx}P(x, 1) = C \left[ e^{\lambda_1 x/\theta_1} x^{-\mu/\theta_1} \cdot \frac{\lambda_1 x - \mu}{\theta_1 x} \cdot \mathcal{K}(x) + x^{-1} \right]. \quad (287)$$

*Step 3: evaluate at  $x = 1$ .*

$$\begin{aligned} \mathbb{E}[N_1] &= \left. \frac{d}{dx}P(x, 1) \right|_{x=1} \\ &= \frac{\mu\pi(0, 0)}{\theta_1(1 - \lambda_2\mathbb{E}[B_C])} \left[ e^{\lambda_1/\theta_1} \cdot \frac{\lambda_1 - \mu}{\theta_1} \cdot \mathcal{K}(1) + 1 \right], \end{aligned} \quad (288)$$

where  $\mathcal{K}(1) = \int_0^1 e^{-\lambda_1 t/\theta_1} t^{\mu/\theta_1-1} dt = (\theta_1/\lambda_1)^{\mu/\theta_1} \gamma(\mu/\theta_1, \lambda_1/\theta_1)$ , with  $\gamma(\cdot, \cdot)$  the lower incomplete gamma function. The factor  $(\lambda_1 - \mu) < 0$  (since  $\rho_1 < 1$ ), so the bracket is positive. No elementary simplification is known;  $\mathbb{E}[N_1]$  is evaluated numerically. As  $\theta_1 \rightarrow 0^+$ ,  $\mathbb{E}[N_1]$  recovers the Model-A value (118).

### Little's Law with abandonment

The universal relation  $\mathbb{E}[N_i] = \lambda_i \mathbb{E}[W_i]$  holds for any stable system. Here  $\lambda_i$  is the *arrival* rate of all class- $i$  customers, including those that will eventually abandon, so  $\mathbb{E}[W_i]$  is the mean time in queue averaged over both served and abandoned customers:

$$\mathbb{E}[W_1] = \frac{\mathbb{E}[N_1]}{\lambda_1}, \quad \mathbb{E}[W_2] = \frac{\mathbb{E}[N_2]}{\lambda_2}. \quad (289)$$

Using the throughput  $\mu(1 - \pi_0)$  instead would give the conditional waiting time of customers who complete service only — a different object. As  $\theta_1 \rightarrow 0^+$  both means recover the Model-A values (118)–(124).

## 10 Analysis of Model- $B_2^H$ ( $\gamma_1 > 0$ head-of-line, $\gamma_2 = \theta_1 = \theta_2 = 0$ )

Model- $B_2^H$  is a deliberately simplified variant of the one-way jockeying model Model- $B_2$ . In Model- $B_2$  *every* waiting class-1 customer may jockey to the class-2 queue, so the aggregate jockeying rate out of a state grows linearly with the queue length,  $\gamma_1 n_1$ . Here, by contrast, only the *customer at the head of Queue 1* — the one that would be the next class-1 customer to enter service — is allowed to jockey, at the fixed rate  $\gamma_1$  whenever  $n_1 \geq 1$ . The aggregate jockeying rate is therefore  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  rather than  $\gamma_1 n_1$ .

This single change is decisive for tractability. In Model- $B_2$  the linear coefficient  $\gamma_1 n_1$  produces, upon forming the joint PGF, a term proportional to  $xy(x - y) \partial P / \partial x$ , turning the fundamental equation into a first-order linear ODE in  $x$  that requires the integrating-factor machinery of §8. The flat rate  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$ , on the other hand, contributes only the *algebraic* factor  $\gamma_1 y(x - y)$  to the coefficient of  $P(x, y)$ ; *no derivative of  $P$  appears*. The fundamental equation is then purely algebraic and can be solved by the Kernel Method exactly as in Model-A (§5). The analysis below mirrors the analytical treatment of Model-A step by step, and the resulting PGF turns out to be *structurally identical* to that of Model-A, the only difference being a shift in the characteristic roots.

**Stability.** Jockeying relocates a waiting customer between the two queues but does not remove anyone from the system; the total number of customers is therefore conserved by the jockeying mechanism, exactly as in Model-A. Since there is no abandonment, the chain is positive recurrent if and only if the total traffic intensity satisfies

$$\rho = \frac{\lambda_1 + \lambda_2}{\mu} < 1, \quad (290)$$

with  $\rho_1 = \lambda_1/\mu$  governing the priority class.

## 10.1 Balance equations

Because the jockeying rate is now constant in  $n_1$  rather than linear, the balance equations are *not* obtained by zeroing parameters in the general Lemma 1 (whose jockeying coefficient is  $\gamma_1 n_1$ ); they must be stated directly. The jockeying transition  $(n_1, n_2) \rightarrow (n_1 - 1, n_2 + 1)$  occurs at the fixed rate  $\gamma_1$  whenever  $n_1 \geq 1$ . On the state space  $S$  the global balance equations read

$$\begin{aligned} & [\lambda_1 + \lambda_2 + \mu + \gamma_1] \pi(n_1, n_2) \\ &= \mu \pi(n_1 + 1, n_2) + \lambda_1 \pi(n_1 - 1, n_2) + \lambda_2 \pi(n_1, n_2 - 1) + \gamma_1 \pi(n_1 + 1, n_2 - 1) \end{aligned} \quad n_1 \geq 1, n_2 \geq 1, \quad (291)$$

$$[\lambda_1 + \lambda_2 + \mu + \gamma_1] \pi(n_1, 0) = \mu \pi(n_1 + 1, 0) + \lambda_1 \pi(n_1 - 1, 0) \quad n_1 \geq 1, n_2 = 0, \quad (292)$$

$$\begin{aligned} & [\lambda_1 + \lambda_2 + \mu] \pi(0, n_2) \\ &= \mu \pi(1, n_2) + \gamma_1 \pi(1, n_2 - 1) + \lambda_2 \pi(0, n_2 - 1) + \mu \pi(0, n_2 + 1) \end{aligned} \quad n_1 = 0, n_2 \geq 1, \quad (293)$$

$$[\lambda_1 + \lambda_2 + \mu] \pi(0, 0) = (\lambda_1 + \lambda_2) \pi_0 + \mu \pi(1, 0) + \mu \pi(0, 1), \quad (294)$$

$$(\lambda_1 + \lambda_2) \pi_0 = \mu \pi(0, 0). \quad (295)$$

Two structural remarks are in order. First, jockeying is an *out*-transition from any state with  $n_1 \geq 1$  (hence the  $+\gamma_1$  in the left-hand sides of (291) and (292)) and an *in*-transition into any state with  $n_2 \geq 1$  from the state  $(n_1 + 1, n_2 - 1)$  (hence the  $\gamma_1 \pi(n_1 + 1, n_2 - 1)$  in (291) and the  $\gamma_1 \pi(1, n_2 - 1)$  in (293)). Second, when  $n_2 = 0$  there is no jockeying in-flow, since it would have to originate from the non-existent state  $(n_1 + 1, -1)$ ; and when  $n_1 = 0$  there is no jockeying out-flow. The remaining transitions — Poisson arrivals at  $\lambda_1, \lambda_2$  and service completions at  $\mu$  — are exactly those of Model-A.

## 10.2 Fundamental PGF equation

**Lemma 4** (Fundamental equation for Model- $B_2^H$ ). *Assume positive recurrence, so that the stationary distribution exists and  $P(x, y) = \sum_{n_1, n_2 \geq 0} \pi(n_1, n_2) x^{n_1} y^{n_2}$  is analytic on the closed unit polydisc. Then  $P(x, y)$  and the boundary function  $P_y(y) = P(0, y)$  satisfy*

$$\begin{aligned} & [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2 + \gamma_1 y(x - y)] P(x, y) \\ &= [\mu(x - y) + \gamma_1 y(x - y)] P_y(y) - \mu x(1 - y) \pi(0, 0). \end{aligned} \quad (296)$$

*Proof.* Apart from the jockeying terms, equations (291)–(295) coincide with the balance equations of Model-A. Multiplying each balance equation by  $x^{n_1} y^{n_2}$ , summing over its domain and multiplying through by  $xy$ , the Model-A part reproduces exactly the left- and right-hand sides of the Model-A fundamental equation (80),

$$[(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2] P(x, y) = \mu(x - y) P_y(y) - \mu x(1 - y) \pi(0, 0),$$

so it remains only to collect the jockeying contribution. Its *out*-flow appears in every state with  $n_1 \geq 1$ :

$$\gamma_1 \sum_{n_1 \geq 1} \sum_{n_2 \geq 0} \pi(n_1, n_2) x^{n_1} y^{n_2} = \gamma_1 [P(x, y) - P_y(y)],$$

while its *in*-flow enters every state with  $n_2 \geq 1$  from  $(n_1 + 1, n_2 - 1)$ :

$$\gamma_1 \sum_{n_1 \geq 0} \sum_{n_2 \geq 1} \pi(n_1 + 1, n_2 - 1) x^{n_1} y^{n_2} = \gamma_1 \frac{y}{x} \sum_{m_1 \geq 1} \sum_{m_2 \geq 0} \pi(m_1, m_2) x^{m_1} y^{m_2} = \gamma_1 \frac{y}{x} [P(x, y) - P_y(y)],$$

where we substituted  $m_1 = n_1 + 1$ ,  $m_2 = n_2 - 1$ . The net jockeying contribution (out minus in) is therefore  $\gamma_1 (1 - \frac{y}{x}) [P(x, y) - P_y(y)]$ ; multiplying by  $xy$  gives  $\gamma_1 y(x - y) [P(x, y) - P_y(y)]$ . Adding this to the Model-A equation and moving the  $P_y$  part to the right-hand side yields (296).  $\square$

*Remark.* Equation (296) contains *no derivative* of  $P$ . The flat jockeying rate has contributed the algebraic factor  $\gamma_1 y(x - y)$  to the coefficient of  $P(x, y)$  where Model- $B_2$ , with its linear rate  $\gamma_1 n_1$ , contributes the convective term  $\gamma_1 x y(x - y) \partial P / \partial x$ . Setting  $\gamma_1 = 0$  recovers (80) verbatim.



### 10.3 Closed-form solution

The following theorem is the main result of this section; its proof follows the analytical (Kernel Method) treatment of Model-A in §5.1.1.

**Theorem 4** (Closed-form PGF for Model- $B_2^H$ ). *Consider the two-class non-preemptive priority queue with Poisson arrival rates  $\lambda_1, \lambda_2$ , common exponential service rate  $\mu$ , and head-of-line class-1 jockeying at rate  $\gamma_1 > 0$  (with  $\gamma_2 = \theta_1 = \theta_2 = 0$ ). Under the stability condition (290), the joint PGF  $P(x, y) = \mathbb{E}[x^{N_1} y^{N_2}]$  is the rational function*

$$P(x, y) = \frac{\mu \rho (1 - \rho) (1 - y)}{\lambda_1 (x^*(y) - y) (x^+(y) - x)}, \quad (297)$$

where  $x^*(y)$  and  $x^+(y)$  are the two roots of the kernel quadratic

$$\lambda_1 x^2 - [\lambda_1 + \lambda_2(1 - y) + \mu + \gamma_1]x + (\mu + \gamma_1 y) = 0, \quad (298)$$

with  $x^*(y)$  the root inside the unit disk. This is the same rational form as Model-A (Theorem 1); the two differ only in the kernel, where  $\mu$  has been replaced by  $\mu + \gamma_1 y$ .

**Corollary 6.** Under (290), the empty-state probabilities of Model- $B_2^H$  coincide with those of Model-A,

$$\pi_0 = 1 - \rho, \quad \pi(0, 0) = \rho(1 - \rho). \quad (299)$$

In particular  $P(1, 1) = 1 - \pi_0 = \rho$  and the diagonal generating function satisfies  $P(z, z) = \pi(0, 0)/(1 - \rho z)$ .

*Proof of Corollary 6.* Set  $x = y = z$  in the fundamental equation (296). The jockeying factor  $\gamma_1 y(x - y)$  vanishes on the diagonal — a direct consequence of the fact that jockeying conserves the total number of customers — and so does the term  $\mu(x - y)P_y(y)$ , leaving

$$[(\lambda_1 + \lambda_2 + \mu)z^2 - \mu z - (\lambda_1 + \lambda_2)z^3]P(z, z) = -\mu z(1 - z)\pi(0, 0).$$

Dividing by  $z$  and factoring the bracket as  $-(\lambda z - \mu)(z - 1)$ , with  $\lambda = \lambda_1 + \lambda_2$ ,

$$(\lambda z - \mu)(1 - z)P(z, z) = -\mu(1 - z)\pi(0, 0).$$

Cancelling  $(1 - z) \neq 0$  for  $z \in [0, 1)$  gives the diagonal identity

$$P(z, z) = \frac{\mu \pi(0, 0)}{\mu - \lambda z} = \frac{\pi(0, 0)}{1 - \rho z}. \quad (300)$$

Setting  $z = 1$  and using the normalisation  $P(1, 1) = 1 - \pi_0$  yields  $1 - \pi_0 = \pi(0, 0)/(1 - \rho)$ . The idle balance (295) gives  $\lambda \pi_0 = \mu \pi(0, 0)$ , i.e.  $\pi(0, 0) = \rho \pi_0$ ; substituting,  $(1 - \pi_0)(1 - \rho) = \rho \pi_0$ , hence  $\pi_0 = 1 - \rho$  and  $\pi(0, 0) = \rho(1 - \rho)$ .  $\square$

#### 10.3.1 Analytical approach for Model- $B_2^H$

*Proof of Theorem 4.* We apply the *Kernel Method* to the fundamental equation (296). Fix  $y \in (0, 1]$  and seek the values of  $x$  at which the coefficient of  $P(x, y)$  — the *kernel* — vanishes. Since  $P(x, y)$  is a PGF it is bounded on the closed unit disk, so at any such root the right-hand side of (296) must vanish as well. Dividing the kernel by  $y \neq 0$ ,

$$0 = (\lambda_1 + \lambda_2 + \mu)x - \mu - \lambda_1 x^2 - \lambda_2 xy + \gamma_1(x - y),$$

and rearranging into a quadratic in  $x$  gives exactly (298),

$$\lambda_1 x^2 - [\lambda_1 + \lambda_2(1 - y) + \mu + \gamma_1]x + (\mu + \gamma_1 y) = 0,$$

whose two roots are

$$x^\pm(y) = \frac{A(y) \pm \sqrt{A(y)^2 - 4\lambda_1(\mu + \gamma_1 y)}}{2\lambda_1}, \quad A(y) := \lambda_1 + \lambda_2(1 - y) + \mu + \gamma_1. \quad (301)$$

This is precisely the Model-A kernel quadratic (87) with  $\mu$  replaced by  $\mu + \gamma_1 y$ . We label the root carrying the negative sign  $x^*(y)$  and the other  $x^+(y)$ . By Vieta's formulas the product of the roots is  $x^*(y)x^+(y) = (\mu + \gamma_1 y)/\lambda_1$ .



To locate the roots relative to the unit disk we evaluate the kernel quadratic, written as  $f(x) := \lambda_1 x^2 - A(y)x + (\mu + \gamma_1 y)$ , at  $x = 1$ :

$$f(1) = \lambda_1 - [\lambda_1 + \lambda_2(1 - y) + \mu + \gamma_1] + (\mu + \gamma_1 y) = -(1 - y)(\lambda_2 + \gamma_1) < 0 \quad \text{for } y \in [0, 1).$$

Since  $f$  is an upward parabola ( $\lambda_1 > 0$ ) with  $f(1) < 0$ , the point  $x = 1$  lies strictly between its two roots, so  $x^*(y) < 1 < x^+(y)$  for all  $y \in [0, 1)$ . Hence  $x^*(y)$  is the unique root inside the unit disk, and at  $y = 1$  one checks  $f(1) = 0$  with  $x^*(1) = 1$  and  $x^+(1) = (\mu + \gamma_1)/\lambda_1 > 1$  (the latter exceeding 1 because  $\lambda_1 < \mu < \mu + \gamma_1$  under stability). For later use, we record  $\frac{d}{dy}x^*(1)$ . The cleanest route is implicit differentiation of  $f(x, y) := \lambda_1 x^2 - A(y)x + (\mu + \gamma_1 y) = 0$  at the point  $(x, y) = (1, 1)$ , where  $f_y = \lambda_2 x + \gamma_1$  and  $f_x = 2\lambda_1 x - A(y)$  evaluate to  $f_y(1, 1) = \lambda_2 + \gamma_1$  and  $f_x(1, 1) = \lambda_1 - \mu - \gamma_1$ , so  $\frac{d}{dy}x^*(1) = -f_y/f_x$ :

$$\frac{d}{dy}x^*(1) = \frac{\lambda_2 + \gamma_1}{\mu + \gamma_1 - \lambda_1}. \quad (302)$$

Unlike Model-A's (88), the numerator carries an extra  $\gamma_1$ : the constant term  $\mu + \gamma_1 y$  of the kernel (298) is itself  $y$ -dependent here, whereas in Model-A (and in Model- $C_2^H$  below) the constant term is independent of  $y$ .

Setting  $x = x^*(y)$  in (296) and equating the right-hand side to zero,

$$[\mu(x^*(y) - y) + \gamma_1 y(x^*(y) - y)]P_y(y) - \mu x^*(y)(1 - y)\pi(0, 0) = 0,$$

that is,  $(x^*(y) - y)(\mu + \gamma_1 y)P_y(y) = \mu x^*(y)(1 - y)\pi(0, 0)$ , whence the boundary function is

$$P_y(y) = \frac{\mu x^*(y)(1 - y)\pi(0, 0)}{(\mu + \gamma_1 y)(x^*(y) - y)}. \quad (303)$$

As a consistency check, set  $y = 0$  in (303): the factor  $\mu + \gamma_1 y$  becomes  $\mu$  and  $1 - y$  becomes 1, so  $P_y(0) = \mu x^*(0)\pi(0, 0)/(\mu x^*(0)) = \pi(0, 0)$  (using  $x^*(0) > 0$ , the smaller of the two positive roots), recovering the boundary value  $P_y(0) = \pi(0, 0)$ .

Finally, substitute (303) and  $\pi(0, 0) = \rho(1 - \rho)$  (Corollary 6) back into (296). Writing the kernel as  $-\lambda_1 y(x - x^*(y))(x - x^+(y))$  and simplifying the right-hand side,

$$\begin{aligned} [\mu(x - y) + \gamma_1 y(x - y)]P_y(y) - \mu x(1 - y)\pi(0, 0) &= \mu(1 - y)\pi(0, 0) \left[ \frac{(x - y)x^*(y)}{x^*(y) - y} - x \right] \\ &= \mu(1 - y)\pi(0, 0) \frac{y(x - x^*(y))}{x^*(y) - y}, \end{aligned}$$

where the bracket collapses because  $(x - y)x^* - x(x^* - y) = y(x - x^*)$ . Dividing by the factored kernel, the common factor  $(x - x^*(y))$  cancels and we obtain

$$P(x, y) = \frac{\mu(1 - y)\pi(0, 0)y(x - x^*(y))/(x^*(y) - y)}{-\lambda_1 y(x - x^*(y))(x - x^+(y))} = \frac{\mu\rho(1 - \rho)(1 - y)}{\lambda_1(x^*(y) - y)(x^+(y) - x)},$$

which is (297) and concludes the proof.  $\square$

### 10.3.2 Probabilistic approach for Model- $B_2^H$

For Model-A, the boundary function  $P_y(y)$  admits an independent derivation by the *Pollaczek-Khinchine* (PK) formula (§5.1.2): conditional on  $N_1 = 0$ , class 2 experiences an  $M/G/1$  queue whose service times are i.i.d. copies of a class-1 busy period, and the PK formula then delivers  $P_y(y)$ . *This route does not transfer to Model- $B_2^H$ .* Head-of-line jockeying injects class-1 customers into Queue 2 at the instants when  $N_1 \geq 1$ ; the resulting arrival stream into Queue 2 is the superposition of the exogenous Poisson stream  $\lambda_2$  and a state-dependent jockeying stream. This merged stream is *neither* Poisson *nor* independent of the system state — it is correlated with  $N_1$ . Since the PK formula requires PASTA, which in turn demands both a Poisson arrival process *and* the lack-of-anticipation property, and the merged input violates both, class 2 does not see a renewal  $M/G/1$  input and no PK representation of  $P_y(y)$  is available. The analytical derivation of §10.3.1 is consequently the only route, and we retain this subsection solely to record why the probabilistic one is unavailable — in the same spirit as the discussion of Models  $B$  and  $B_2$ .

## 10.4 Limits and sanity checks

As  $\gamma_1 \rightarrow 0^+$  the kernel quadratic (298) reduces to Model-A's quadratic (87), so  $x^*(y) \rightarrow x_A^*(y)$ , and formula (297) reduces to the factored form of Theorem 1; the empty-state probabilities already equal those of Model-A by Corollary 6. Setting  $x = y = 1$  in (297) through the diagonal identity (300) gives  $P(1, 1) = \rho = 1 - \pi_0$ , confirming normalisation, and the diagonal  $P(z, z) = \pi(0, 0)/(1 - \rho z)$  is exactly that of the combined  $M/M/1(\lambda, \mu)$  queue — as it must be, since jockeying conserves the total customer count.

## 10.5 Mean queue lengths and mean waiting times

### 10.5.1 Class-1 marginal PGF and mean queue length

Setting  $y = 1$  in the fundamental equation (296), the term  $\mu x(1 - y)\pi(0, 0)$  vanishes and the coefficient of  $P(x, 1)$  factors as  $-(x - 1)(\lambda_1 x - (\mu + \gamma_1))$ , while the right-hand side becomes  $(x - 1)(\mu + \gamma_1)P_y(1)$ . Cancelling  $(x - 1)$  for  $x \neq 1$ ,

$$P(x, 1) = \frac{(\mu + \gamma_1) P_y(1)}{\mu + \gamma_1 - \lambda_1 x} = \frac{P_y(1)}{1 - \rho_B x}, \quad \rho_B := \frac{\lambda_1}{\mu + \gamma_1}. \quad (304)$$

This is a geometric class-1 marginal: the priority queue behaves as an  $M/M/1$  queue whose effective *clearance rate* is  $\mu + \gamma_1$ , since a waiting class-1 customer at the head of Queue 1 departs that queue either by entering service (rate  $\mu$ ) or by jockeying (rate  $\gamma_1$ ). The constant  $P_y(1)$  is pinned by the normalisation  $P(1, 1) = 1 - \pi_0 = \rho$ , giving  $P_y(1) = \rho(1 - \rho_B)$ , so that  $P(x, 1) = \rho(1 - \rho_B)/(1 - \rho_B x)$ . Differentiating by the quotient rule and evaluating at  $x = 1$ ,

$$\mathbb{E}[N_1] = \left. \frac{d}{dx} P(x, 1) \right|_{x=1} = \frac{\rho(1 - \rho_B) \rho_B}{(1 - \rho_B)^2} = \frac{\rho \rho_B}{1 - \rho_B}. \quad (305)$$

This is Model-A's formula (118) with the priority load  $\rho_1$  replaced by the *jockeying-reduced* load  $\rho_B = \lambda_1/(\mu + \gamma_1) \leq \rho_1$ .

### 10.5.2 Total queue length from the diagonal

Because jockeying conserves the total number of customers, the combined process  $N_1 + N_2$  (together with the in-service customer) is the queue-length process of an ordinary  $M/M/1(\lambda, \mu)$  queue, and the diagonal generating function (300) is  $P(z, z) = \rho(1 - \rho)/(1 - \rho z)$ . Since  $\frac{d}{dz} P(z, z)|_{z=1} = \mathbb{E}[N_1] + \mathbb{E}[N_2]$ , differentiating gives

$$\mathbb{E}[N_1] + \mathbb{E}[N_2] = \left. \frac{d}{dz} \frac{\rho(1 - \rho)}{1 - \rho z} \right|_{z=1} = \frac{\rho^2(1 - \rho)}{(1 - \rho)^2} = \frac{\rho^2}{1 - \rho}, \quad (306)$$

identical to the combined  $M/M/1$  result of Model-A (122).

### 10.5.3 Class-2 mean queue length by subtraction

Subtracting (305) from (306),

$$\begin{aligned} \mathbb{E}[N_2] &= \frac{\rho^2}{1 - \rho} - \frac{\rho \rho_B}{1 - \rho_B} = \frac{\rho[\rho(1 - \rho_B) - \rho_B(1 - \rho)]}{(1 - \rho)(1 - \rho_B)} \\ &= \frac{\rho(\rho - \rho_B)}{(1 - \rho)(1 - \rho_B)}. \end{aligned} \quad (307)$$

The excess load admits a transparent decomposition,

$$\rho - \rho_B = \frac{\lambda_1 + \lambda_2}{\mu} - \frac{\lambda_1}{\mu + \gamma_1} = \frac{(\lambda_1 + \lambda_2)(\mu + \gamma_1) - \lambda_1 \mu}{\mu(\mu + \gamma_1)} = \frac{\lambda_2 \mu + \lambda_1 \gamma_1 + \lambda_2 \gamma_1}{\mu(\mu + \gamma_1)} = \rho_2 + \frac{\lambda_1 \gamma_1}{\mu(\mu + \gamma_1)},$$

so  $\rho - \rho_B$  exceeds the bare class-2 load  $\rho_2$  precisely by the load that jockeying transfers from Queue 1 into Queue 2. At  $\gamma_1 = 0$  it collapses to  $\rho_2$  and (307) recovers Model-A's  $\mathbb{E}[N_2] = \rho \rho_2 / [(1 - \rho_1)(1 - \rho)]$ .

### 10.5.4 Mean waiting times via Little's Law

We apply **Little's Law** to each queue separately, recalling that  $N_i$  counts only the customers *waiting*, so  $\mathbb{E}[N_i] = \Lambda_i \mathbb{E}[W_i]$  with  $\Lambda_i$  the long-run rate at which customers *enter* Queue  $i$  and  $\mathbb{E}[W_i]$  their mean residence time in that queue (from entry until they leave it, by service entry or by jockeying).

Every class-1 customer enters Queue 1 exactly once, by exogenous arrival, so  $\Lambda_1 = \lambda_1$  and

$$\mathbb{E}[W_1] = \frac{\mathbb{E}[N_1]}{\lambda_1} = \frac{\rho \rho_B}{\lambda_1(1 - \rho_B)} = \frac{\rho}{\mu + \gamma_1 - \lambda_1}. \quad (308)$$

Compared with Model-A's  $\mathbb{E}[W_1] = \rho/(\mu - \lambda_1)$  (123), the denominator is enlarged by  $\gamma_1$ : head-of-line jockeying drains the priority queue faster and shortens class-1 waits.

Queue 2 receives a *mixed* input: the exogenous Poisson stream at rate  $\lambda_2$  together with the jockeying stream. The latter fires at rate  $\gamma_1$  whenever  $N_1 \geq 1$ , so its long-run rate is  $\gamma_1 \mathbb{P}(N_1 \geq 1)$ . The probability that a class-1 customer is waiting follows from the geometric marginal (304) by subtracting the  $N_1 = 0$  mass from the busy mass,

$$\mathbb{P}(N_1 \geq 1) = P(1, 1) - P(0, 1) = \rho - \rho(1 - \rho_B) = \rho \rho_B,$$

since  $P(1, 1) = 1 - \pi_0 = \rho$  and  $P(0, 1) = P_y(1) = \rho(1 - \rho_B)$ . The total entrance rate into Queue 2 is therefore  $\Lambda_2 = \lambda_2 + \gamma_1 \rho \rho_B$ , and Little's Law gives the mean residence time per Queue-2 entrant,

$$\mathbb{E}[W_2] = \frac{\mathbb{E}[N_2]}{\Lambda_2} = \frac{\rho(\rho - \rho_B)}{(1 - \rho)(1 - \rho_B)(\lambda_2 + \gamma_1 \rho \rho_B)}. \quad (309)$$

As  $\gamma_1 \rightarrow 0^+$  the jockeying stream disappears,  $\Lambda_2 \rightarrow \lambda_2$  and  $\rho_B \rightarrow \rho_1$ , and (309) recovers Model-A's  $\mathbb{E}[W_2] = \rho/[\mu(1 - \rho_1)(1 - \rho)]$  (124).

## 11 Analysis of Model- $C_2^H$ ( $\theta_1 > 0$ head-of-line, $\gamma_1 = \gamma_2 = \theta_2 = 0$ )

Model- $C_2^H$  is the abandonment counterpart of Model- $B_2^H$ , and a simplified variant of the class-1 abandonment model Model- $C_2$ . In Model- $C_2$  *every* waiting class-1 customer is impatient and reneges at the linear rate  $\theta_1 n_1$ , turning the class-1 subsystem into an  $M/M/1+M$  (Erlang-A) queue and the fundamental equation into a first-order ODE. Here only the *customer at the head of Queue 1* may abandon, at the fixed rate  $\theta_1$  whenever  $n_1 \geq 1$ ; the aggregate abandonment rate is  $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$  rather than  $\theta_1 n_1$ . As with Model- $B_2^H$ , the flat rate contributes an *algebraic* term to the fundamental equation rather than a derivative, and the Kernel Method applies directly. Unlike jockeying, abandonment is a *true departure*: it removes a customer from the system and so breaks the conservation of the total customer count that Model- $B_2^H$  enjoyed. The empty-state probabilities and the class-2 mean queue length consequently require a separate argument.

**Stability.** Because only the head of Queue 1 abandons, the aggregate abandonment rate is bounded by  $\theta_1$  regardless of queue length, so — in contrast to the Erlang-A class-1 subsystem of Model- $C_2$ , which is positive recurrent for all loads — a genuine stability condition is required. As shown in Corollary 7 below, the chain is positive recurrent if and only if  $\pi_0 > 0$ , that is

$$(\mu + \theta_1)(\mu - \lambda) + \lambda_1 \theta_1 > 0, \quad \lambda = \lambda_1 + \lambda_2. \quad (310)$$

This is *weaker* than  $\rho < 1$ : if  $\rho < 1$  then  $\mu - \lambda > 0$  and the condition holds trivially, but it can also hold with  $\rho \geq 1$  provided  $\theta_1$  is large enough — head abandonment accelerates the clearance of the priority queue and can stabilise the system at offered loads exceeding the service capacity. Moreover (310) implies  $\rho_C := \lambda_1/(\mu + \theta_1) < 1$ , so the priority subqueue is itself stable whenever the condition holds.

### 11.1 Balance equations

The abandonment transition  $(n_1, n_2) \rightarrow (n_1 - 1, n_2)$  occurs at the fixed rate  $\theta_1$  whenever  $n_1 \geq 1$ . Since both a service completion and a head abandonment from  $(n_1 + 1, n_2)$  land in  $(n_1, n_2)$ , the two mechanisms combine into the coefficient  $\mu + \theta_1$  on the in-flows. On  $S$ ,

$$\begin{aligned} & [\lambda_1 + \lambda_2 + \mu + \theta_1] \pi(n_1, n_2) \\ &= (\mu + \theta_1) \pi(n_1 + 1, n_2) + \lambda_1 \pi(n_1 - 1, n_2) + \lambda_2 \pi(n_1, n_2 - 1) \end{aligned} \quad n_1 \geq 1, n_2 \geq 1, \quad (311)$$

$$[\lambda_1 + \lambda_2 + \mu + \theta_1] \pi(n_1, 0) = (\mu + \theta_1) \pi(n_1 + 1, 0) + \lambda_1 \pi(n_1 - 1, 0) \quad n_1 \geq 1, n_2 = 0, \quad (312)$$

$$\begin{aligned} & [\lambda_1 + \lambda_2 + \mu] \pi(0, n_2) \\ &= (\mu + \theta_1) \pi(1, n_2) + \lambda_2 \pi(0, n_2 - 1) + \mu \pi(0, n_2 + 1) \end{aligned} \quad n_1 = 0, n_2 \geq 1, \quad (313)$$

$$[\lambda_1 + \lambda_2 + \mu] \pi(0, 0) = (\lambda_1 + \lambda_2) \pi_0 + (\mu + \theta_1) \pi(1, 0) + \mu \pi(0, 1), \quad (314)$$

$$(\lambda_1 + \lambda_2) \pi_0 = \mu \pi(0, 0). \quad (315)$$

The contrast with Model- $C_2$  is the same as before: there the in-flow coefficients  $\mu + \theta_1(n_1 + 1)$  and the out-flow coefficient  $\theta_1 n_1$  grow with the queue length, whereas here they are the constants  $\mu + \theta_1$  and  $\theta_1$  throughout.

## 11.2 Fundamental PGF equation

**Lemma 5** (Fundamental equation for Model- $C_2^H$ ). *Assume positive recurrence. Then the joint PGF  $P(x, y)$  and the boundary function  $P_y(y) = P(0, y)$  satisfy*

$$\begin{aligned} & [(\lambda_1 + \lambda_2 + \mu)xy - \mu y - \lambda_1 x^2 y - \lambda_2 x y^2 + \theta_1 y(x - 1)] P(x, y) \\ &= [\mu(x - y) + \theta_1 y(x - 1)] P_y(y) - \mu x(1 - y) \pi(0, 0). \end{aligned} \quad (316)$$

*Proof.* As in Lemma 4, the non-abandonment terms of (311)–(315) reproduce the Model-A fundamental equation (80) after multiplication by  $x^{n_1} y^{n_2}$ , summation, and multiplication by  $xy$ . The abandonment out-flow appears in every state with  $n_1 \geq 1$ ,

$$\theta_1 \sum_{n_1 \geq 1} \sum_{n_2 \geq 0} \pi(n_1, n_2) x^{n_1} y^{n_2} = \theta_1 [P(x, y) - P_y(y)],$$

and its in-flow enters every state  $(n_1, n_2)$  from  $(n_1 + 1, n_2)$ ,

$$\theta_1 \sum_{n_1 \geq 0} \sum_{n_2 \geq 0} \pi(n_1 + 1, n_2) x^{n_1} y^{n_2} = \theta_1 x^{-1} \sum_{m_1 \geq 1} \sum_{n_2 \geq 0} \pi(m_1, n_2) x^{m_1} y^{n_2} = \theta_1 x^{-1} [P(x, y) - P_y(y)],$$

with  $m_1 = n_1 + 1$ . The net contribution is  $\theta_1(1 - x^{-1})[P(x, y) - P_y(y)]$ ; multiplying by  $xy$  gives  $\theta_1 y(x - 1)[P(x, y) - P_y(y)]$ . Adding to the Model-A equation and rearranging yields (316).  $\square$

*Remark.* Again no derivative of  $P$  appears: the algebraic factor  $\theta_1 y(x - 1)$  replaces the convective term  $\theta_1 x y(x - 1) \partial P / \partial x$  of Model- $C_2$ . Setting  $\theta_1 = 0$  recovers (80). Unlike the jockeying factor of Model- $B_2^H$ , the abandonment factor  $\theta_1 y(x - 1)$  does *not* vanish on the diagonal  $x = y$ , which is the analytic signature of the broken conservation. This algebraic collapse—no derivative term, and hence no singularity to resolve—is the head-of-line regime anticipated in the remark following Lemma 1.

## 11.3 Closed-form solution

**Theorem 5** (Closed-form PGF for Model- $C_2^H$ ). *Consider the two-class non-preemptive priority queue with Poisson arrival rates  $\lambda_1, \lambda_2$ , common exponential service rate  $\mu$ , and head-of-line class-1 abandonment at rate  $\theta_1 > 0$  (with  $\gamma_1 = \gamma_2 = \theta_2 = 0$ ). Under the stability condition (310), the joint PGF  $P(x, y) = \mathbb{E}[x^{N_1} y^{N_2}]$  is the rational function*

$$P(x, y) = \frac{\mu(\mu + \theta_1)(1 - y) \pi(0, 0)}{\lambda_1 (x^+(y) - x) [(\mu + \theta_1 y) x^*(y) - y(\mu + \theta_1)]}, \quad (317)$$

where  $x^*(y)$  and  $x^+(y)$  are the two roots of the kernel quadratic

$$\lambda_1 x^2 - [\lambda_1 + \lambda_2(1 - y) + \mu + \theta_1] x + (\mu + \theta_1) = 0, \quad (318)$$

with  $x^*(y)$  the root inside the unit disk, and  $\pi(0, 0)$  is given by Corollary 7. As  $\theta_1 \rightarrow 0^+$  the formula recovers Model-A (Theorem 1) exactly.

**Corollary 7.** *Under (310), the empty-state probabilities of Model- $C_2^H$  are*

$$\pi_0 = \frac{(\mu + \theta_1)(\mu - \lambda) + \lambda_1 \theta_1}{\mu(\mu + \theta_1) + \lambda_1 \theta_1}, \quad \pi(0, 0) = \frac{\lambda[(\mu + \theta_1)(\mu - \lambda) + \lambda_1 \theta_1]}{\mu[\mu(\mu + \theta_1) + \lambda_1 \theta_1]}, \quad (319)$$

with  $\lambda = \lambda_1 + \lambda_2$ . At  $\theta_1 \rightarrow 0^+$  these reduce to  $\pi_0 \rightarrow 1 - \rho$  and  $\pi(0, 0) \rightarrow \rho(1 - \rho)$ , recovering Model-A.

*Proof of Corollary 7.* Two evaluations of the fundamental equation (316) suffice. First set  $y = 1$ : the term  $\mu x(1 - y)\pi(0, 0)$  vanishes, the coefficient of  $P(x, 1)$  factors as  $-(x - 1)(\lambda_1 x - (\mu + \theta_1))$ , and the right-hand side becomes  $(x - 1)(\mu + \theta_1)P_y(1)$ . Cancelling  $(x - 1)$ ,

$$P(x, 1) = \frac{(\mu + \theta_1) P_y(1)}{\mu + \theta_1 - \lambda_1 x} = \frac{P_y(1)}{1 - \rho_C x}, \quad \rho_C := \frac{\lambda_1}{\mu + \theta_1}. \quad (320)$$

This geometric marginal shows that the priority queue clears at the effective rate  $\mu + \theta_1$  (service completions plus head abandonments). Evaluating at  $x = 1$  and using  $P(1, 1) = 1 - \pi_0$  gives

$$1 - \pi_0 = \frac{P_y(1)}{1 - \rho_C} \implies P_y(1) = (1 - \pi_0)(1 - \rho_C). \quad (321)$$

Second, set  $x = 1$  in (316). The coefficient of  $P(1, y)$  collapses to  $\lambda_2 y(1 - y)$  and the right-hand side to  $\mu(1 - y)[P_y(y) - \pi(0, 0)]$ , yielding the class-2 marginal relation

$$\lambda_2 y P(1, y) = \mu[P_y(y) - \pi(0, 0)]. \quad (322)$$

Letting  $y \rightarrow 1$  and using (321) and the idle balance (315),  $\pi(0, 0) = \lambda\pi_0/\mu$ ,

$$\lambda_2(1 - \pi_0) = \mu[P_y(1) - \pi(0, 0)] = \mu(1 - \pi_0)(1 - \rho_C) - \lambda\pi_0.$$

This is a linear equation in  $\pi_0$ . Expanding and collecting the  $\pi_0$  terms on the left,

$$[\mu(1 - \rho_C) + \lambda - \lambda_2]\pi_0 = \mu(1 - \rho_C) - \lambda_2, \quad \text{i.e.} \quad \pi_0 = \frac{\mu(1 - \rho_C) - \lambda_2}{\mu(1 - \rho_C) + \lambda_1},$$

where we used  $\lambda - \lambda_2 = \lambda_1$ . Substituting  $\mu(1 - \rho_C) = \mu(\mu + \theta_1 - \lambda_1)/(\mu + \theta_1)$  and multiplying numerator and denominator by  $\mu + \theta_1$ , the numerator becomes  $\mu(\mu + \theta_1 - \lambda_1) - \lambda_2(\mu + \theta_1) = (\mu + \theta_1)(\mu - \lambda) + \lambda_1\theta_1$  and the denominator  $\mu(\mu + \theta_1 - \lambda_1) + \lambda_1(\mu + \theta_1) = \mu(\mu + \theta_1) + \lambda_1\theta_1$ , yielding the first identity in (319); the second follows from  $\pi(0, 0) = \lambda\pi_0/\mu$ . The denominator  $\mu(\mu + \theta_1) + \lambda_1\theta_1$  is strictly positive, so  $\pi_0 > 0$  is equivalent to the stability condition (310).  $\square$

### 11.3.1 Analytical approach for Model- $C_2^H$

*Proof of Theorem 5.* We again apply the Kernel Method to (316). Dividing the coefficient of  $P(x, y)$  by  $y \neq 0$  and rearranging gives the kernel quadratic (318),

$$\lambda_1 x^2 - [\lambda_1 + \lambda_2(1 - y) + \mu + \theta_1]x + (\mu + \theta_1) = 0,$$

which is the Model-A kernel quadratic with  $\mu$  replaced by  $\mu + \theta_1$  both in the linear coefficient and in the constant term. Its roots are

$$x^\pm(y) = \frac{A(y) \pm \sqrt{A(y)^2 - 4\lambda_1(\mu + \theta_1)}}{2\lambda_1}, \quad A(y) := \lambda_1 + \lambda_2(1 - y) + \mu + \theta_1, \quad (323)$$

with product  $x^*(y)x^+(y) = (\mu + \theta_1)/\lambda_1$  by Vieta's formulas. Evaluating the quadratic  $f(x) := \lambda_1 x^2 - A(y)x + (\mu + \theta_1)$  at  $x = 1$ ,

$$f(1) = \lambda_1 - A(y) + (\mu + \theta_1) = -\lambda_2(1 - y) < 0 \quad \text{for } y \in [0, 1),$$

so  $x = 1$  lies between the two roots and  $x^*(y) < 1 < x^+(y)$  for  $y \in [0, 1)$ ; at  $y = 1$ ,  $x^*(1) = 1$  and  $x^+(1) = (\mu + \theta_1)/\lambda_1 > 1$  (the latter by  $\rho_C < 1$ ). Hence  $x^*(y)$  is the unique root inside the unit disk. Differentiating (323) at  $y = 1$ ,

$$\frac{d}{dy}x^*(1) = \frac{\lambda_2}{\mu + \theta_1 - \lambda_1}, \quad (324)$$

the analogue of (88) and (302).

Setting  $x = x^*(y)$  in (316) forces the right-hand side to vanish,

$$[\mu(x^*(y) - y) + \theta_1 y(x^*(y) - 1)]P_y(y) = \mu x^*(y)(1 - y)\pi(0, 0),$$

and, regrouping the bracket as  $\mu(x^* - y) + \theta_1 y(x^* - 1) = (\mu + \theta_1 y)x^* - y(\mu + \theta_1)$ , the boundary function is

$$P_y(y) = \frac{\mu x^*(y)(1 - y)\pi(0, 0)}{(\mu + \theta_1 y)x^*(y) - y(\mu + \theta_1)}. \quad (325)$$

At  $y = 0$  the denominator is  $\Delta(0) = \mu x^*(0)$ , so  $P_y(0) = \mu x^*(0)\pi(0,0)/(\mu x^*(0)) = \pi(0,0)$  (using  $x^*(0) > 0$ , the smaller positive root), as required.

Substituting (325) into (316), writing the kernel as  $-\lambda_1 y(x - x^*(y))(x - x^+(y))$ , and abbreviating the denominator of (325) by  $\Delta(y) := (\mu + \theta_1 y)x^*(y) - y(\mu + \theta_1)$ , the right-hand side becomes

$$\begin{aligned} [\mu(x - y) + \theta_1 y(x - 1)]P_y(y) - \mu x(1 - y)\pi(0,0) &= \frac{\mu(1 - y)\pi(0,0)}{\Delta(y)} \left\{ [\mu(x - y) + \theta_1 y(x - 1)]x^*(y) - x\Delta(y) \right\} \\ &= \frac{\mu(1 - y)\pi(0,0)}{\Delta(y)} (\mu + \theta_1)y(x - x^*(y)), \end{aligned}$$

where, after substituting  $\Delta(y) = (\mu + \theta_1 y)x^*(y) - y(\mu + \theta_1)$ , the curly bracket is *linear* in  $x^*(y)$  and collapses term-by-term to  $(\mu + \theta_1)y(x - x^*(y))$  (no use of the kernel relation is required). Dividing by the factored kernel, the factor  $(x - x^*(y))$  cancels and

$$P(x, y) = \frac{\mu(\mu + \theta_1)(1 - y)\pi(0,0)y(x - x^*(y))/\Delta(y)}{-\lambda_1 y(x - x^*(y))(x - x^+(y))} = \frac{\mu(\mu + \theta_1)(1 - y)\pi(0,0)}{\lambda_1(x^+(y) - x)\Delta(y)},$$

which is (317) and concludes the proof.  $\square$

### 11.3.2 Probabilistic approach for Model- $C_2^H$

As for Model- $B_2^H$ , the Pollaczek–Khinchine route used for Models  $A$  and  $C_2$  does not transfer. In Model- $C_2$  the class-1 subsystem seen by class 2 is a clean  $M/M/1+M$  (Erlang-A) queue whose busy period  $B_C$  is an i.i.d. renewal quantity, and class 2 sees an  $M/G/1$  queue with service  $B_C$ . Under head-only abandonment this clean renewal structure is lost: abandonment acts at the fixed rate  $\theta_1$  only while a class-1 customer waits ( $N_1 \geq 1$ ) but is inactive when the single class-1 present is in service, so the departure rate of the class-1 subsystem is state-dependent at its boundary and the class-1 busy period is not the busy period of any single  $M/M/1$  or  $M/M/1+M$  queue. Consequently class 2 does not see a renewal  $M/G/1$  input and no Pollaczek–Khinchine representation of  $P_y(y)$  is available. We retain this subsection, in parallel with the other models, only to record why the analytical route of §11.3.1 is the operative one.

## 11.4 Limits and sanity checks

As  $\theta_1 \rightarrow 0^+$  the kernel quadratic (318) reduces to Model- $A$ 's, so  $x^*(y) \rightarrow x_A^*(y)$  and  $\Delta(y) \rightarrow \mu(x_A^*(y) - y)$ ; together with  $\pi(0,0) \rightarrow \rho(1 - \rho)$  from Corollary 7, formula (317) reduces to the factored Model- $A$  PGF of Theorem 1. Setting  $x = 1$  in (317) and  $y \rightarrow 1$  reproduces  $P(1,1) = 1 - \pi_0$  via (321), confirming normalisation.

## 11.5 Mean queue lengths and mean waiting times

### 11.5.1 Class-1 marginal PGF and mean queue length

The class-1 marginal is the geometric law (320); pinning the constant by  $P(1,1) = 1 - \pi_0$  as in (321) gives  $P(x,1) = (1 - \pi_0)(1 - \rho_C)/(1 - \rho_C x)$ . Differentiating and evaluating at  $x = 1$ ,

$$\mathbb{E}[N_1] = \left. \frac{d}{dx} P(x,1) \right|_{x=1} = \frac{(1 - \pi_0)\rho_C}{1 - \rho_C}, \quad \rho_C = \frac{\lambda_1}{\mu + \theta_1}, \quad (326)$$

with  $1 - \pi_0 = \lambda(\mu + \theta_1)/[\mu(\mu + \theta_1) + \lambda_1\theta_1]$  from Corollary 7. As in Model- $B_2^H$ , the priority queue behaves as an  $M/M/1$  queue with effective clearance rate  $\mu + \theta_1$ , the only difference from Model- $B_2^H$ 's (305) being the prefactor  $1 - \pi_0$ , which here is no longer equal to  $\rho$  because abandonment alters the idle probability.

### 11.5.2 Class-2 mean queue length

Abandonment is a true departure and so breaks the *conservation of the total customer count* that underlies the diagonal argument: in Model- $B_2^H$  the jockeying factor  $\gamma_1 y(x - y)$  vanishes on the diagonal  $x = y$ , whereas here the abandonment factor  $\theta_1 y(x - 1)$  does not. Concretely, setting  $x = y = z$  in (316) leaves the term  $\theta_1 z(z - 1)P_y(z)$  on the right-hand side, so the diagonal equation no longer closes and the route that furnished  $\mathbb{E}[N_1] + \mathbb{E}[N_2]$  for Model- $B_2^H$  is unavailable. We

instead differentiate the class-2 marginal. From the marginal relation (322),  $P(1, y) = \mu [P_y(y) - \pi(0, 0)] / (\lambda_2 y)$ , and since  $\mathbb{E}[N_2] = \frac{d}{dy} P(1, y) \big|_{y=1}$ ,

$$\mathbb{E}[N_2] = \frac{\mu}{\lambda_2} P'_y(1) - (1 - \pi_0), \quad (327)$$

where we used  $\frac{\mu}{\lambda_2} [P_y(1) - \pi(0, 0)] = P(1, 1) = 1 - \pi_0$ . It remains to compute  $P'_y(1)$  from (325). Both the numerator  $\mu x^*(y)(1 - y)\pi(0, 0)$  and the denominator  $\Delta(y) = (\mu + \theta_1 y)x^*(y) - y(\mu + \theta_1)$  vanish at  $y = 1$  (since  $x^*(1) = 1$  gives  $\Delta(1) = (\mu + \theta_1) - (\mu + \theta_1) = 0$ ), so  $P_y(y)$  is a 0/0 form there. Writing  $N(y)$  and  $\Delta(y)$  for numerator and denominator, a second-order Taylor expansion about  $y = 1$  gives the value and the slope

$$P_y(1) = \frac{N'(1)}{\Delta'(1)} = (1 - \pi_0)(1 - \rho_C), \quad P'_y(1) = \frac{N''(1)\Delta'(1) - N'(1)\Delta''(1)}{2\Delta'(1)^2},$$

which is why both the first and the second root sensitivities of  $x^*$  at  $y = 1$  are needed, namely (324) together with

$$\frac{d^2}{dy^2} x^*(1) = \frac{2(\mu + \theta_1)\lambda_2^2}{(\mu + \theta_1 - \lambda_1)^3}$$

(the latter obtained by differentiating  $x^*(y)$  in (323) twice). Substituting these into  $P'_y(1)$ , then into (327), and simplifying yields the closed form

$$\mathbb{E}[N_2] = \frac{\lambda_2 \lambda (\mu + \theta_1) [(\mu + \theta_1)^2 - \lambda_1 \theta_1]}{(\mu + \theta_1 - \lambda_1) [\mu(\mu + \theta_1) + \lambda_1 \theta_1] [(\mu + \theta_1)(\mu - \lambda) + \lambda_1 \theta_1]}, \quad (328)$$

with  $\lambda = \lambda_1 + \lambda_2$ . The last bracket in the denominator is exactly the stability numerator of  $\pi_0$  from (319), so  $\mathbb{E}[N_2]$  is finite precisely under the stability condition (310) and diverges as  $\pi_0 \rightarrow 0^+$ . As  $\theta_1 \rightarrow 0^+$  the right-hand side reduces to  $\lambda_2 \lambda / [(\mu - \lambda_1)(\mu - \lambda)] = \rho \rho_2 / [(1 - \rho_1)(1 - \rho)]$ , recovering Model-A's (122).

### 11.5.3 Mean waiting times via Little's Law

We apply **Little's Law** to each queue. Every class-1 customer enters Queue 1 once by exogenous arrival and later leaves it either by entering service or by abandoning from the head; in both cases it leaves Queue 1, so the entrance rate is  $\Lambda_1 = \lambda_1$  and

$$\mathbb{E}[W_1] = \frac{\mathbb{E}[N_1]}{\lambda_1} = \frac{(1 - \pi_0)\rho_C}{\lambda_1(1 - \rho_C)}. \quad (329)$$

Here  $\mathbb{E}[W_1]$  is the mean time a class-1 customer spends *waiting* in Queue 1, averaged over all class-1 arrivals, whether they are eventually served or abandon. Queue 2, by contrast, receives no jockeying input and its customers never abandon ( $\theta_2 = 0$ ), so every class-2 arrival is eventually served and the entrance rate is simply  $\Lambda_2 = \lambda_2$ :

$$\mathbb{E}[W_2] = \frac{\mathbb{E}[N_2]}{\lambda_2} = \frac{\lambda(\mu + \theta_1) [(\mu + \theta_1)^2 - \lambda_1 \theta_1]}{(\mu + \theta_1 - \lambda_1) [\mu(\mu + \theta_1) + \lambda_1 \theta_1] [(\mu + \theta_1)(\mu - \lambda) + \lambda_1 \theta_1]}. \quad (330)$$

Both reduce to the corresponding Model-A expressions as  $\theta_1 \rightarrow 0^+$ .

## 11.6 Comparison of the head-of-line models

Table 4 summarises the structural differences between the two head-of-line variants and their full-rate counterparts. The common thread is that replacing a length-proportional mechanism rate ( $\gamma_1 n_1$  or  $\theta_1 n_1$ ) by a head-of-line rate ( $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  or  $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$ ) converts the governing ODE into an algebraic equation, so that  $P(x, y)$  becomes rational. For Model- $B_2^H$  the result is structurally identical to Model-A, the sole change being  $\mu \rightarrow \mu + \gamma_1 y$  in the kernel; for Model- $C_2^H$  the broken conservation introduces the extra factor  $(\mu + \theta_1)$  and the modified boundary denominator  $\Delta(y)$ , and shifts the stability boundary to (310).



Table 4: Structural comparison across the priority-queue models.

| Property              | A                | B <sub>2</sub>     | B <sub>2</sub> <sup>H</sup>            | C <sub>2</sub>                  | C <sub>2</sub> <sup>H</sup>            |
|-----------------------|------------------|--------------------|----------------------------------------|---------------------------------|----------------------------------------|
| Mechanism rate        | —                | $\gamma_1 n_1$     | $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$ | $\theta_1 n_1$                  | $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$ |
| PGF equation type     | algebraic        | ODE                | <b>algebraic</b>                       | ODE                             | <b>algebraic</b>                       |
| $P(x, y)$ form        | rational         | integral+ $_1 F_1$ | <b>rational</b>                        | integral+ $_1 F_1$              | <b>rational</b>                        |
| Same form as Model-A? | ✓                | ×                  | ✓                                      | ×                               | (similar)                              |
| $\pi(0, 0)$           | $\rho(1 - \rho)$ | $\rho(1 - \rho)$   | $\rho(1 - \rho)$                       | (240)                           | (319)                                  |
| $\pi_0$               | $1 - \rho$       | $1 - \rho$         | $1 - \rho$                             | (241)                           | (319)                                  |
| Conserves total $N$ ? | ✓                | ✓                  | ✓                                      | ×                               | ×                                      |
| Stability condition   | $\rho < 1$       | $\rho < 1$         | $\rho < 1$                             | $\lambda_2 \mathbb{E}[B_C] < 1$ | (310)                                  |

## 12 Comparison of the Solved Models

This section compares the five solved models: the baseline (A), the jockeying extension (B<sub>2</sub>), the abandonment extension (C<sub>2</sub>), and their head-of-line variants (B<sub>2</sub><sup>H</sup>, C<sub>2</sub><sup>H</sup>). The first three are the tractable specialisations of Model- $X$ , each obtained by zeroing all but one of the extra mechanisms; the head-of-line variants are the standalone simplifications of §§10–11. Each produces a closed-form or integral-form expression for  $P(x, y)$ . Table 5 collects the key structural facts for the three full-rate models (the head-of-line variants are compared in Table 4 and rejoin the numerical comparison below); the remainder of this section traces the consequences.

Table 5: Structural comparison of Models A, B<sub>2</sub>, and C<sub>2</sub>.

|                                     | Model-A                       | Model-B <sub>2</sub>                  | Model-C <sub>2</sub>                           |
|-------------------------------------|-------------------------------|---------------------------------------|------------------------------------------------|
| Extra mechanism                     | none                          | jockeying 1 → 2<br>( $\gamma_1 > 0$ ) | class-1 abandonment<br>( $\theta_1 > 0$ )      |
| Stability condition                 | $\rho < 1$                    | $\rho < 1$                            | $\rho_2 \cdot {}_1 F_1 < 1$                    |
| Fundamental equation                | algebraic PGF equation        | 1st-order PDE in $x, y$               | 1st-order ODE in $x$                           |
| Pinning of $P_y(y)$                 | zero of kernel: $x = x^*(y)$  | analyticity at $x = y$                | boundedness at $x = 1$                         |
| Probabilistic route                 | yes (P-K with $B$ )           | no (jockeying breaks Poisson)         | yes (P-K with $B_C$ )                          |
| $P_y(y)$                            | rational in $x^*(y)$          | Kummer ratio (223)                    | P-K / Kummer (249)                             |
| $P(x, y)$                           | closed form (80)              | integral form (196)                   | integral form (273)                            |
| $\pi(0, 0)$                         | $\rho(1 - \rho)$              | $\rho(1 - \rho)$                      | $(\rho_1 + \rho_2)\pi_0$ via $\mathbb{E}[B_C]$ |
| $\pi_0$                             | $1 - \rho$                    | $1 - \rho$                            | $> 1 - \rho$ (abandonment increases idle time) |
| $P(z, z)$                           | $\rho(1 - \rho)/(1 - \rho z)$ | $\rho(1 - \rho)/(1 - \rho z)$         | no closed form                                 |
| $\mathbb{E}[N_1] + \mathbb{E}[N_2]$ | $\rho^2/(1 - \rho)$           | $\rho^2/(1 - \rho)$                   | $< \rho^2/(1 - \rho)$                          |
| $\mathbb{E}[N_1]$ closed form       | $\rho\rho_1/(1 - \rho_1)$     | no                                    | no                                             |

### Model-A: the rational baseline

Model-A imposes no extra departure mechanism. Class-1 operates as an independent  $M/M/1(\lambda_1, \mu)$  queue; class-2 sees an  $M/G/1$  with service time equal to a class-1 busy period (by PASTA and the priority decomposition of §3.2). The Pollaczek–Khinchine formula (12) delivers  $P_y(y)$  as a rational function of  $x^*(y)$ , and  $P(x, y)$  is a ratio of polynomials in  $x, y$ , and  $x^*(y)$ . All performance measures —  $\pi_0, \pi(0, 0), \mathbb{E}[N_i], \mathbb{E}[W_i]$  — are in closed form. Model-A also furnishes the structural



identities that persist across the hierarchy:  $P(z, z) = \rho(1 - \rho)/(1 - \rho z)$  (total queue is  $M/M/1$ ),  $\pi_0 = 1 - \rho$ , and  $\pi(0, 0) = \rho(1 - \rho)$ .

### Effect of jockeying: Model-A $\rightarrow$ Model- $B_2$

Adding one-way jockeying ( $\gamma_1 > 0$ ,  $\gamma_2 = 0$ ) couples the two queues: a class-1 customer waiting in Queue 1 may defect to Queue 2 at rate  $\gamma_1$ . The consequences are asymmetric.

*What is preserved.* Jockeying transfers customers but does not remove them, so the total-customer process remains a standard  $M/M/1(\lambda, \mu)$  birth-death chain. Consequently  $P(z, z)$ ,  $\pi_0$ ,  $\pi(0, 0)$ , and  $\mathbb{E}[N]$  are identical to Model-A for all  $\gamma_1 > 0$ .

*What is destroyed.* Jockeying breaks the per-class Poisson/renewal structure: a class-2 customer can no longer identify its effective service time with a class-1 busy period, because class-1 customers now arrive at Queue 2 from inside the system as well as from outside. The P-K route is therefore unavailable.

*What is new.* The fundamental PGF equation gains the convection term  $\gamma_1 xy(x - y)\partial_x P$ , which reduces it to a first-order ODE in  $x$  for each fixed  $y$  (since  $\gamma_2 = 0$  kills the  $\partial_y$  term). The solution closes via an integrating factor with a branch singularity at the *interior* point  $x = y$ ; the branch coefficient must vanish for  $P$  to be holomorphic, and this analyticity condition pins  $P_y(y)$  as a Kummer ratio. The individual means  $\mathbb{E}[N_1]$  and  $\mathbb{E}[N_2]$  change with  $\gamma_1$  (class-1 becomes shorter, class-2 longer), with no closed form for the split.

### Effect of abandonment: Model-A $\rightarrow$ Model- $C_2$

Adding class-1 abandonment ( $\theta_1 > 0$ ,  $\theta_2 = 0$ ,  $\gamma_i = 0$ ) allows each waiting class-1 customer to leave the system at rate  $\theta_1$ .

*What is preserved.* Class-2 customers still see Poisson arrivals (abandonment only affects class-1; the class-2 arrival process is undisturbed). PASTA therefore still applies, and the  $M/G/1$  subsystem for class-2 is intact — now with effective service time  $B_C$  (the busy period of the  $M/M/1 + M$  queue of §3.3) replacing  $B$ . The P-K formula (12) applies with  $B_C$  and delivers  $P_y(y)$ .

*What is destroyed.* Abandonment is a true departure: it reduces the total number of customers and breaks the conservation property. Consequently  $P(z, z)$  no longer has the simple  $M/M/1$  form (120),  $\pi_0 > 1 - \rho$ ,  $\pi(0, 0)$  requires  $\mathbb{E}[B_C]$ , and  $\mathbb{E}[N] < \rho^2/(1 - \rho)$ . The stability condition is also generalised:  $\rho < 1$  is no longer necessary; the relevant condition is  $\lambda_2 \mathbb{E}[B_C] < 1$ , which is strictly weaker for all  $\theta_1 > 0$ .

*What is new.* The fundamental equation gains the convection term  $\theta_1 xy(x - 1)\partial_x P$ , which already reduces it to a first-order ODE in  $x$  for each fixed  $y$  (with  $\theta_2 = 0$  killing the  $\partial_y$  term). The integrating factor now has its singularity at the *boundary*  $x = 1$  with a positive exponent, so  $\mu_I$  vanishes there and mere boundedness of  $P$  forces  $\int_0^1 q \mu_I dt = 0$ . This determines  $P_y(y)$  without any reference to holomorphy. Both  $\mathbb{E}[N_1]$  and  $\mathbb{E}[N_2]$  decrease relative to Model-A as  $\theta_1$  increases: class-2 benefits indirectly because class-1 abandonments free the server sooner.

### Structural comparison of Models $B_2$ and $C_2$

Both models reduce the fundamental equation to a first-order linear ODE in  $x$  and both close in confluent hypergeometric functions; yet the mechanism that pins  $P_y(y)$  is structurally different. The contrast is as follows.

|                          | Model- $C_2$ ( $\theta_1 > 0$ )                                         | Model- $B_2$ ( $\gamma_1 > 0$ )                                                          |
|--------------------------|-------------------------------------------------------------------------|------------------------------------------------------------------------------------------|
| Convection coefficient   | $\theta_1 x(x - 1)$                                                     | $\gamma_1 x(x - y)$                                                                      |
| Operative singular point | boundary $x = 1$                                                        | interior $x = y$                                                                         |
| Exponent at singularity  | $\beta(y) = \lambda_2(1 - y)/\theta_1 > 0$<br>( $\mu_I \rightarrow 0$ ) | $\beta(y) = (1 - y)(\lambda y - \mu)/(\gamma_1 y) < 0$<br>( $\mu_I \rightarrow \infty$ ) |
| Pinning principle        | <i>boundedness</i> : $\int_0^1 q \mu_I dt = 0$                          | <i>analyticity</i> : branch coefficient = 0                                              |
| $\pi(0, 0)$              | $M/G/1$ +renewal (conservation broken)                                  | $\rho(1 - \rho)$ exactly (conservation holds)                                            |
| Probabilistic backbone   | class-2 is $M/G/1$ ; P-K with $B_C$                                     | class-1 is $M/M/1+M$ (reneging = $\gamma_1$ ); no P-K                                    |

In words: in Model- $C_2$  the abandonment convection  $\theta_1 x(x-1)$  places the singularity at the boundary  $x=1$  with a positive integrating-factor exponent, so  $\mu_I$  vanishes there and mere boundedness forces the vanishing integral that determines  $P_y(y)$ . The Poisson character of the class-2 stream survives abandonment, so P-K delivers  $P_y(y)$  probabilistically and an  $M/G/1$  renewal argument delivers  $\pi(0,0)$ . In Model- $B_2$ , the jockeying convection  $\gamma_1 x(x-y)$  places the singularity at the interior point  $x=y$  with a negative exponent, so  $\mu_I$  blows up, the homogeneous mode vanishes, and only the stronger requirement of holomorphy — the absence of the non-integer branch  $(x-y)^{|\beta|}$  — pins  $P_y(y)$ . Jockeying destroys the per-class Poisson/renewal structure, so no P-K identity is available; what remains probabilistic is the total-customer  $M/M/1$  behaviour and the  $M/M/1+M$  reading of the class-1 marginal, which is the origin of the  ${}_1F_1$  factors.

## Common Erlang-A structure

A common object appears in both  $B_2$  and  $C_2$ :

$${}_1F_1\left(1; \frac{\mu}{r} + 1; \frac{\lambda_1}{r}\right), \quad (331)$$

with  $r = \gamma_1$  (Model- $B_2$ ) or  $r = \theta_1$  (Model- $C_2$ ). This is the Erlang-A utilisation factor: in  $C_2$  it equals  $\mu \mathbb{E}[B_C]$  (the mean class-1 busy period scaled by  $\mu$ ), while in  $B_2$  it controls the Kummer ratio in  $P_y(y)$  through the exponent  $\alpha(y) = \mu/(\gamma_1 y)$ . In both cases the function evaluates at a negative argument  $(-\lambda_1/r)$  in  $C_2$  and  $(-\lambda_1 y/\gamma_1)$  in  $B_2$ , and the ratio  ${}_1F_1(\alpha+1; \cdot; \cdot)/{}_1F_1(\alpha; \cdot; \cdot)$  acts as a boundary correction to the Model- $A$  rational formula.

Model- $B_2$  degenerates to Model- $A$  only in the singular limit  $\gamma_1 \rightarrow 0^+$ , in which  $\alpha, \beta \rightarrow \infty$  and the Kummer ratio collapses to the rational expression  $x^*(y)(1-y)/(x^*(y)-y)$  of Model- $A$ ; this limit is not uniform, consistent with  $\gamma_1$  multiplying the highest-order term of the ODE. Model- $C_2$  degenerates to Model- $A$  uniformly as  $\theta_1 \rightarrow 0^+$ :  $\mathbb{E}[B_C] \rightarrow (\mu - \lambda_1)^{-1}$ , and all  $C_2$  formulas recover their Model- $A$  counterparts.

### 12.1 Traffic-intensity sweep

The structural comparison above is now made quantitative. Holding the service rate at  $\mu = 1$  and the arrival split fixed at  $\lambda_1:\lambda_2 = 3:4$ , the offered load  $\rho = \lambda_1 + \lambda_2$  is swept across  $[0.10, 0.95]$  and the stationary performance of the three models is solved exactly on the truncated CTMC. Figure 8 collects the mean queue length  $\mathbb{E}[N]$ , the priority-class queue  $\mathbb{E}[N_1]$ , the priority waiting time  $\mathbb{E}[W_1] = \mathbb{E}[N_1]/\lambda_1$  (Little's Law), and the idle probability  $\pi_0$ ; Figure 9 isolates  $\pi(0,0)$  against the Model- $A$  parabola  $\rho(1-\rho)$ .




Figure 8: Performance versus traffic intensity  $\rho$  for the five solved models — the baseline  $A$ , the jockeying pair  $B_2$ ,  $B_2^H$  ( $\gamma_1 = 0.5$ ), and the abandonment pair  $C_2$ ,  $C_2^H$  ( $\theta_1 = 0.5$ ) — at  $\mu = 1$  and  $\lambda_1 : \lambda_2 = 3 : 4$ ; the vertical line marks the canonical  $\rho = 0.70$ . *Top-left*: total queue  $\mathbb{E}[N]$  —  $A$ ,  $B_2$  and  $B_2^H$  coincide because jockeying conserves  $N$  (Corollaries 1, 6), while  $C_2$  and  $C_2^H$  lie strictly below because abandonment removes customers ( $C_2^H$  by less). *Top-right*: priority queue  $\mathbb{E}[N_1]$  — the head-of-line variants drain it less than their full-rate cousins. *Bottom-left*: priority waiting time  $\mathbb{E}[W_1]$ . *Bottom-right*: idle probability  $\pi_0$ , with the  $1 - \rho$  law (Models  $A$ ,  $B_2$ ,  $B_2^H$ ) shown dotted;  $C_2$  and  $C_2^H$  sit above it for every  $\rho$ , as predicted by Corollaries 5, 7.

Two mechanisms produce two qualitatively different responses. Jockeying is internal: the  $1 \rightarrow 2$  transition relabels a waiting customer without changing the total count, so the  $B_2$  and  $A$  curves for  $\mathbb{E}[N]$  and  $\pi_0$  are indistinguishable across the entire sweep — the server sees the same workload. Abandonment is a true departure at rate  $\theta_1 n_1$ , so  $C_2$  carries a strictly lighter queue and a strictly larger idle fraction: the priority backlog is bled off before it can be served. At  $\rho = 0.70$  (Table 6) this gives  $\pi_0 = 0.3696$  for  $C_2$  against 0.3000 for  $A$  and  $B_2$ , and  $\mathbb{E}[N] = 0.8358$  against 1.6333.

The divergence widens with load. At  $\rho = 0.90$  the baseline class-2 queue reaches  $\mathbb{E}[N_2] = 7.5346$ , whereas  $C_2$  holds it to 1.9245 — a factor of nearly four — because the self-regulating valve  $\theta_1 n_1$  acts most strongly precisely when the queue is long. The priority queue  $\mathbb{E}[N_1]$  is bounded for  $C_2$  for the same reason and never exceeds 0.2292 over the sweep, while the  $A$  value grows monotonically toward the  $\rho \rightarrow 1$  singularity. We note for completeness that near criticality the dense CTMC truncation leaves a boundary tail mass of order  $6 \times 10^{-4}$  for  $A$  and  $B_2$  at  $\rho = 0.95$  (against  $1.5 \times 10^{-7}$  for  $C_2$ , whose tail is light); the  $\rho \geq 0.92$  points for the non-abandoning models therefore carry a small downward truncation bias and should be read as lower bounds.




Figure 9:  $\pi(0,0)$  versus  $\rho$ . The three customer-conserving models  $A$ ,  $B_2$  and  $B_2^H$  lie exactly on the parabola  $\rho(1 - \rho)$  (Corollaries 1, 4, 6: jockeying preserves  $N$ , hence the busy-but-empty probability), whereas the abandonment models  $C_2$  and  $C_2^H$  rise strictly above it — abandonment inflates  $\pi(0,0)$  through the enlarged idle probability  $\pi(0,0) = \rho \pi_0$  (Corollaries 5, 7), the full rate ( $C_2$ ) lifting it more than the head-of-line rate ( $C_2^H$ ).

The invariance of  $\pi(0,0) = \rho(1 - \rho)$  along the  $A$  and  $B_2$  curves of Figure 9 is not a numerical coincidence but a direct reading of Corollary 4: the event “server busy, both queues empty” depends only on the total occupancy  $N$ , and jockeying leaves the law of  $N$  unchanged. For  $C_2$  the identity  $\pi(0,0) = \rho \pi_0$  still holds, but  $\pi_0 > 1 - \rho$ , so the curve is lifted — at  $\rho = 0.70$ ,  $\pi(0,0) = 0.2587$  against the baseline 0.2100 (Table 6). The lift is also asymmetric: whereas the conserving parabola  $\rho(1 - \rho)$  is symmetric about  $\rho = \frac{1}{2}$ , the  $C_2$  curve in Figure 9 attains its maximum to the *right* of that point, the signature of abandonment renormalising the effective load — the busy-but-empty probability keeps climbing past the symmetric-load peak because the self-regulating valve blunts congestion fastest exactly where the baseline is heaviest, before the shrinking busy fraction finally pulls it down.

The two head-of-line variants  $B_2^H$  and  $C_2^H$  added to Figures 8–9 confirm that the qualitative split survives the head-of-line simplification but in a muted form. Model- $B_2^H$  conserves  $N$  exactly like  $B_2$  (Corollary 6), so its  $\mathbb{E}[N]$  and  $\pi_0$  curves are indistinguishable from the  $A/B_2$  pair and lie on the same  $\rho(1 - \rho)$  parabola in Figure 9; it departs only in the split of  $\mathbb{E}[N_1]$  (at  $\rho = 0.70$ ,  $\mathbb{E}[N_1] = 0.1750$  against 0.1545 for  $B_2$ , Table 6). Model- $C_2^H$  tracks  $C_2$  but bleeds the priority queue more gently — at  $\rho = 0.70$  it gives  $\mathbb{E}[N] = 0.9015$  and  $\pi_0 = 0.3636$ , both lying between the baseline and the full-rate  $C_2$  values (0.8358 and 0.3696). The structural reason for this systematic “weaker-mechanism” behaviour is isolated in §12.5.

Table 6 reads off the full set of stationary descriptors at three loads; the class-1 loss accounting — throughput, abandonment flow, and the loss fraction  $P_{\text{loss}}^{(1)}$  — is carried in the compact companion Table 7 so the main table stays within the margins. Two columns deserve a reviewer’s caution. First, the priority waiting time  $\mathbb{E}[W_1]$  is smallest for  $C_2$  at every load (Table 8: 0.5941 at  $\rho = 0.90$  against 0.6808 for  $B_2$  and 1.4651 for  $A$ ), but  $\mathbb{E}[W_i] = \mathbb{E}[N_i]/\lambda_i$  is the mean time a customer is *counted in queue*  $i$ , and that residence is shortened by every departure mode — service, *and* attrition. In  $C_2$  a class-1 customer leaves the count by being served *or* by abandoning; in  $B_2$  by being served *or* by jockeying into class 2. A smaller  $\mathbb{E}[W_1]$  therefore does not by itself certify faster service, and cross-model  $\mathbb{E}[W_1]$  comparisons conflate latency reduction with customer loss. The magnitude of that confound is exactly the class-1 loss fraction  $L_1 = \theta_1 \mathbb{E}[N_1]/\lambda_1$  derived in §12.4 (eq. (334)): in  $C_2$  at  $\rho = 0.70$  a fraction  $L_1 = 0.232$  of class-1 arrivals leaves by abandonment rather than by service, so the reported  $\mathbb{E}[W_1] = 0.4639$  averages the queue residence over a population of which only a fraction 0.768 ever reaches the server. Tagging served class-1 customers in simulation (§13) isolates the fair latency: the conditional mean time-to-service of served class-1 customers is 0.3863 for  $C_2$ , against the count-based unconditional 0.4639 and against 0.9944 for Model- $A$  (where the conditional latency coincides with the unconditional 1.0000 up to simulation error, since no class-1 customer is lost). The genuine service-latency advantage over Model- $A$  is therefore real, but the part of the apparent  $\mathbb{E}[W_1]$  gap that is *attrition* rather than faster service is

Table 6: Comparison of Models A,  $B_2$  and  $B_2^H$  ( $\gamma_1 = 0.5$ ), and  $C_2$  and  $C_2^H$  ( $\theta_1 = 0.5$ ) at three traffic intensities with  $\lambda_1:\lambda_2 = 3:4$ ,  $\mu = 1$ . The primed models use the head-of-line rate  $\mathbf{1}_{\{n_1 \geq 1\}}$ ; the unprimed use the full rate  $n_1$ .

| $\rho$ | Model   | $E[N_1]$ | $E[N_2]$ | $E[N]$ | $E[W_1]$ | $E[W_2]$ | $\pi_0$ | $\pi(0,0)$ |
|--------|---------|----------|----------|--------|----------|----------|---------|------------|
| 0.50   | A       | 0.1364   | 0.3636   | 0.5000 | 0.6364   | 1.2727   | 0.5000  | 0.2500     |
|        | $B_2$   | 0.0767   | 0.4233   | 0.5000 | 0.3578   | 1.4817   | 0.5000  | 0.2500     |
|        | $B_2^H$ | 0.0833   | 0.4167   | 0.5000 | 0.3889   | 1.4583   | 0.5000  | 0.2500     |
|        | $C_2$   | 0.0712   | 0.2521   | 0.3233 | 0.3323   | 0.8823   | 0.5356  | 0.2678     |
|        | $C_2^H$ | 0.0778   | 0.2593   | 0.3370 | 0.3630   | 0.9074   | 0.5333  | 0.2667     |
| 0.70   | A       | 0.3000   | 1.3333   | 1.6333 | 1.0000   | 3.3333   | 0.3000  | 0.2100     |
|        | $B_2$   | 0.1545   | 1.4788   | 1.6333 | 0.5151   | 3.6970   | 0.3000  | 0.2100     |
|        | $B_2^H$ | 0.1750   | 1.4583   | 1.6333 | 0.5833   | 3.6458   | 0.3000  | 0.2100     |
|        | $C_2$   | 0.1392   | 0.6967   | 0.8358 | 0.4639   | 1.7417   | 0.3696  | 0.2587     |
|        | $C_2^H$ | 0.1591   | 0.7424   | 0.9015 | 0.5303   | 1.8561   | 0.3636  | 0.2545     |
| 0.90   | A       | 0.5651   | 7.5346   | 8.0997 | 1.4651   | 14.6506  | 0.1000  | 0.0900     |
|        | $B_2$   | 0.2626   | 7.8371   | 8.0997 | 0.6808   | 15.2388  | 0.1000  | 0.0900     |
|        | $B_2^H$ | 0.3115   | 7.7882   | 8.0997 | 0.8077   | 15.1437  | 0.1000  | 0.0900     |
|        | $C_2$   | 0.2292   | 1.9245   | 2.1537 | 0.5941   | 3.7421   | 0.2146  | 0.1931     |
|        | $C_2^H$ | 0.2760   | 2.2084   | 2.4844 | 0.7157   | 4.2941   | 0.2025  | 0.1823     |

exactly the fraction  $L_1 = 0.232$  of class-1 demand that  $C_2$  sheds outright. The same caveat applies to  $B_2$  for a different reason — only a fraction 0.742 of its class-1 customers are served directly from queue 1, the rest jockeying into class 2, which pulls its conditional time-to-service 0.4433 below the count-based 0.5151 — so only Model-A has a genuinely vacuous distinction.

Second, the  $E[W_2]$  column shows that jockeying actively *harms* class 2: at  $\rho = 0.70$ ,  $B_2$  raises  $E[W_2]$  to 3.6970 from the baseline 3.3333 (+10.9%), and at  $\rho = 0.90$  to 15.2388 from 14.6506. This is the direct cost of the  $1 \rightarrow 2$  transition — displaced class-1 customers arrive at the tail of the class-2 queue and lengthen it ( $E[N_2]$  rises from 1.3333 to 1.4788 at  $\rho = 0.70$ ). Abandonment, by contrast, lightens both classes, since it reduces the offered work rather than redistributing it.

## 12.2 The jockeying mechanism (Model $B_2$ )

Figure 10 isolates the action of  $\gamma_1$  at fixed load  $\rho = 0.70$ , sweeping  $\gamma_1 \in [10^{-3}, 50]$ .




Figure 10: Model- $B_2$  at  $\rho = 0.70$  ( $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1$ ). *Left*:  $E[N_1]$  decreases and  $E[N_2]$  increases monotonically in  $\gamma_1$  (Model-A values dotted). *Centre*:  $E[N]$  is flat — jockeying conserves the total count. *Right*:  $\pi(0,0) \equiv \rho(1-\rho) = 0.21$  for all  $\gamma_1$ , the numerical signature of Corollary 4.

The monotone decrease of  $E[N_1]$  in  $\gamma_1$  follows directly from the rate structure: each waiting class-1 customer generates a jockeying event at rate  $\gamma_1$ , so the expected outflow from the class-1 queue is  $\gamma_1 E[N_1]$ , an additional drain that acts independently of the service process and grows with  $\gamma_1$ . At  $\gamma_1 = 0.5$  this halves the priority queue, from  $E[N_1] = 0.3000$  to 0.1545 (−48.5%, Table 6). The mass is not destroyed but transferred:  $E[N_2]$  rises correspondingly from 1.3333 to 1.4788, and the centre panel confirms  $E[N] = 1.6333$  is held constant to plotting precision. The right panel is the cleanest empirical check of Corollary 4: because every jockeying transition preserves  $N$ , the load on the server — and therefore  $\pi_0 = 1 - \rho$  and  $\pi(0,0) = \rho(1 - \rho) = 0.21$  — is exactly

Table 7: Class-1 loss accounting accompanying Table 6, at the same three loads and parameters ( $\lambda_1:\lambda_2 = 3:4$ ,  $\mu = 1$ ; mechanism parameter 0.5). *Throughput* is the carried rate  $\mu(1 - \pi_0)$ ; *abandonment rate* is the class-1 loss flow ( $\theta_1 \mathbb{E}[N_1]$  for the full rate,  $\theta_1 \mathbb{P}(N_1 \geq 1)$  for head-of-line); and  $P_{\text{loss}}^{(1)} = \theta_1 \mathbb{E}[N_1]/\lambda_1$  is the fraction of class-1 arrivals shed before service (eq. (334)). The conserving models  $A$ ,  $B_2$ ,  $B_2^H$  lose nothing; only  $C_2$  and  $C_2^H$  have  $P_{\text{loss}}^{(1)} > 0$ , and at every row throughput + abandonment rate = offered load  $\rho$ .

| $\rho$ | Model   | Throughput | Aband. rate | $P_{\text{loss}}^{(1)}$ |
|--------|---------|------------|-------------|-------------------------|
| 0.50   | A       | 0.5000     | 0           | 0                       |
|        | $B_2$   | 0.5000     | 0           | 0                       |
|        | $B_2^H$ | 0.5000     | 0           | 0                       |
|        | $C_2$   | 0.4644     | 0.0356      | 0.166                   |
|        | $C_2^H$ | 0.4667     | 0.0333      | 0.156                   |
| 0.70   | A       | 0.7000     | 0           | 0                       |
|        | $B_2$   | 0.7000     | 0           | 0                       |
|        | $B_2^H$ | 0.7000     | 0           | 0                       |
|        | $C_2$   | 0.6304     | 0.0696      | 0.232                   |
|        | $C_2^H$ | 0.6364     | 0.0636      | 0.212                   |
| 0.90   | A       | 0.9000     | 0           | 0                       |
|        | $B_2$   | 0.9000     | 0           | 0                       |
|        | $B_2^H$ | 0.9000     | 0           | 0                       |
|        | $C_2$   | 0.7854     | 0.1146      | 0.297                   |
|        | $C_2^H$ | 0.7975     | 0.1025      | 0.266                   |

Table 8: Class-1 mean queue length  $E[N_1]$  and waiting time  $E[W_1]$  under Models A,  $B_2$  ( $\gamma_1 = 0.5$ ) and  $C_2$  ( $\theta_1 = 0.5$ ), with  $\mu = 1$  and  $\lambda_1:\lambda_2 = 3:4$ .

| $\rho$ | $E[N_1]$ |        |        | $E[W_1]$ |        |        |
|--------|----------|--------|--------|----------|--------|--------|
|        | A        | $B_2$  | $C_2$  | A        | $B_2$  | $C_2$  |
| 0.50   | 0.1364   | 0.0767 | 0.0712 | 0.6364   | 0.3578 | 0.3323 |
| 0.70   | 0.3000   | 0.1545 | 0.1392 | 1.0000   | 0.5151 | 0.4639 |
| 0.90   | 0.5651   | 0.2626 | 0.2292 | 1.4651   | 0.6808 | 0.5941 |

$\gamma_1$ -invariant.




Figure 11: Class-1 queue-residence ratio  $\mathbb{E}[W_1]/\mathbb{E}[W_2]$  versus  $\gamma_1$  at  $\rho = 0.70$ , with the Model-A value dotted. The ratio decreases toward zero because  $\mathbb{E}[W_1] = \mathbb{E}[N_1]/\lambda_1$  measures time spent in the class-1 queue, which the jockeying drain  $\gamma_1 \mathbb{E}[N_1]$  collapses; it does not measure time to service.

Figure 11 must be read with the residence-time caveat of §12.1 firmly in mind. As  $\gamma_1$  grows the ratio  $\mathbb{E}[W_1]/\mathbb{E}[W_2]$  falls monotonically from the Model-A value of 0.300 toward zero, which would naively suggest an ever-improving class-1 service. The mechanism is the opposite: a large  $\gamma_1$  ejects class-1 customers from their own queue almost immediately, so the *count*  $\mathbb{E}[N_1]$  and hence  $\mathbb{E}[W_1]$  vanish, but the ejected customers simply wait out their delay at the back of the class-2 queue. The falling ratio records the dissolution of the class-1 queue, not an acceleration of class-1 service. This is the precise sense in which strong one-way jockeying *erodes the priority structure*: in the limit there is effectively a single class.

### 12.3 Class asymmetry

Fixing  $\rho = 0.70$  and varying the split  $\alpha = \lambda_1/(\lambda_1 + \lambda_2)$  exposes how the mechanisms interact with the class mix. Figure 12 compares the three models; Figure 13 resolves the jockeying case at three rates.




Figure 12: Class-asymmetry sweep at  $\rho = 0.70$ :  $\mathbb{E}[N_1]$ ,  $\mathbb{E}[N_2]$ , the priority ratio  $\mathbb{E}[W_1]/\mathbb{E}[W_2]$  (vertical dotted lines mark each model's ratio-extremising split over the swept range  $[0.1, 0.9]$ ), and the absolute priority benefit  $\mathbb{E}[W_2] - \mathbb{E}[W_1]$ , for Models A,  $B_2$  and  $B_2^H$  ( $\gamma_1 = 0.5$ ), and  $C_2$  and  $C_2^H$  ( $\theta_1 = 0.5$ ).

As  $\alpha \rightarrow 0$  the priority class is vanishingly rare and its mechanism barely engages: all models converge in  $\mathbb{E}[N_1]$ . As  $\alpha \rightarrow 1$  the priority class dominates the offered load and the non-preemptive discipline gives it near-exclusive access to the server, driving  $\mathbb{E}[N_1] \rightarrow 0$  in every model. The interesting behaviour is interior. The absolute priority benefit  $\mathbb{E}[W_2] - \mathbb{E}[W_1]$  (bottom-right) is largest where class 2 bears the most displaced load; jockeying narrows this gap relative to A by exporting class-1 delay into the class-2 statistic, while abandonment reshapes it by removing class-1 work outright.

The priority ratio  $\mathbb{E}[W_1]/\mathbb{E}[W_2]$  (bottom-left) carries the sharpest structural signal. For Model-A it is *exactly*  $\alpha$ -invariant: writing the ratio as  $\mathbb{E}[W_1]/\mathbb{E}[W_2] = (\mathbb{E}[N_1] \lambda_2)/(\mathbb{E}[N_2] \lambda_1)$  and substituting the closed forms  $\mathbb{E}[N_1] = \rho \rho_1/(1 - \rho_1)$  and  $\mathbb{E}[N] = \rho^2/(1 - \rho)$  collapses the split dependence entirely,

$$\left. \frac{\mathbb{E}[W_1]}{\mathbb{E}[W_2]} \right|_A = 1 - \rho \quad \text{for every } \alpha \in (0, 1), \quad (332)$$

the flat line at  $0.300 = 1 - \rho$  in the panel. The two mechanisms break this invariance in *opposite, monotone* directions. Jockeying makes the ratio strictly *decreasing* in  $\alpha$ , so it peaks at 0.188 at the left endpoint  $\alpha^* = 0.10$  (a larger priority share feeds the  $1 \rightarrow 2$  drain harder); abandonment makes



it strictly *increasing*, peaking at 0.357 at the right endpoint  $\alpha^* = 0.90$  (a larger priority share is exactly where the abandonment valve sheds the most class-1 work). The head-of-line variants track their parents —  $B_2^H$  peaks at 0.193 ( $\alpha^* = 0.10$ ) and  $C_2^H$  at 0.395 ( $\alpha^* = 0.90$ ) — and are likewise monotone. No curve turns over: across all five models the maximiser  $\alpha^*$  sits at a *boundary* of the swept range  $[0.1, 0.9]$ , never at an interior point. An interior  $\alpha^*$  would require mechanism engagement (which strengthens as the priority class grows) to balance class dominance at some intermediate split; here the two effects pull the ratio the same way throughout, so the extremum is always pushed to an endpoint.




Figure 13: Model- $B_2$ :  $\mathbb{E}[N_1]$  and  $\mathbb{E}[N_2]$  versus the split  $\alpha$  for three jockeying rates  $\gamma_1 \in \{0.1, 1.0, 10.0\}$  at  $\rho = 0.70$ . Stronger jockeying flattens  $\mathbb{E}[N_1]$  toward zero across the whole range of  $\alpha$  and loads the difference onto  $\mathbb{E}[N_2]$ .

The family in Figure 13 shows that the jockeying drain dominates the class mix: at  $\gamma_1 = 10$  the priority queue is essentially empty for every  $\alpha$ , because the per-customer ejection rate  $\gamma_1$  overwhelms the arrival rate  $\lambda_1$  regardless of the split. At  $\gamma_1 = 0.1$  the curves still track the Model- $A$  shape, the drain being a small perturbation. The crossover between these regimes occurs where  $\gamma_1$  is comparable to the effective class-1 service rate, consistent with the singular role of  $\gamma_1$  in the  $B_2$  ODE noted above.

## 12.4 The abandonment mechanism and the throughput–quality tradeoff (Model $C_2$ )

Figure 14 sweeps  $\theta_1 \in [10^{-3}, 50]$  at  $\rho = 0.70$ .






Figure 14: Model- $C_2$  at  $\rho = 0.70$ . *Top-left:*  $\mathbb{E}[N_1] \rightarrow 0$  as the abandonment valve  $\theta_1 n_1$  strengthens. *Top-right:*  $\mathbb{E}[N_2]$  decreases monotonically (no rebound at these parameters). *Bottom-left:*  $\pi_0 > 1 - \rho$  for all  $\theta_1 > 0$  (Corollary 5). *Bottom-right:* the throughput-quality tradeoff — throughput falls and the abandonment rate rises with  $\theta_1$ .

The priority queue  $\mathbb{E}[N_1]$  collapses toward zero as  $\theta_1$  grows: the abandonment outflow  $\theta_1 \mathbb{E}[N_1]$  is a self-regulating valve that opens wider the longer the queue, so no class-1 backlog can persist. The idle-probability panel is the empirical face of Corollary 5: every positive  $\theta_1$  lifts  $\pi_0$  strictly above the Model-A value  $1 - \rho = 0.30$ , because customers who abandon never reach the server and the busy fraction shrinks accordingly. This sweep also settles, as a *negative finding*, a question flagged in the simulation brief. Two effects compete in  $\mathbb{E}[N_2]$  as  $\theta_1$  rises: abandonment is class-1-specific, so clearing the priority backlog lets the server reach class 2 sooner — an effect that could in principle make  $\mathbb{E}[N_2]$  rise transiently before falling — set against the plain reduction in the total work the class-2 queue inherits once class 1 is bled off faster. At  $\rho = 0.70$  the second effect dominates throughout:  $\mathbb{E}[N_2]$  is *monotonically decreasing* in  $\theta_1$  with no initial hump, contradicting the conjectured non-monotonicity. We observed no rebound at any load tested and conjecture that  $\mathbb{E}[N_2]$  is monotone decreasing in  $\theta_1$  for every  $\rho < 1$ ; a proof, or a counterexample at extreme class asymmetry, is left to future work (§14.5). The bottom-right panel quantifies the price and, in doing so, supplies an independent numerical check. Throughput falls from the Model-A value  $\mu\rho = 0.70$  to 0.6304 at  $\theta_1 = 0.5$ , a deficit of 0.0696; computed *separately* as the abandonment flow  $\theta_1 \mathbb{E}[N_1]$ , the lost rate reproduces that same 0.0696, so carried plus lost returns the offered 0.70. Because the throughput deficit  $\mu\rho - \mu(1 - \pi_0)$  and the abandonment integral  $\theta_1 \mathbb{E}[N_1]$  are distinct functionals of the stationary law — one read from the idle probability of Corollary 5, the other from the mean queue length — their agreement is not a restatement of flow balance but an independent validation of both.

Written as a fraction, this balance defines the model's *loss*. The server delivers carried work at rate  $\mu(1 - \pi_0)$  whenever it is busy, and the offered rate is  $\lambda_1 + \lambda_2 = \mu\rho$ ; since in steady state every arrival is either served or abandons, flow conservation forces the total abandonment rate to equal the *idle excess* that abandonment creates over the no-attribution baseline,

$$\Lambda_{\text{loss}} = \mu\rho - \mu(1 - \pi_0) = \mu(\pi_0 - (1 - \rho)), \quad (333)$$

the strictly positive gap  $\pi_0 - (1 - \rho) > 0$  of Corollary 5. The identity (333) holds for either abandonment rate — the full-rate mechanism realises  $\Lambda_{\text{loss}} = \theta_1 \mathbb{E}[N_1]$  and the head-of-line mechanism  $\Lambda_{\text{loss}} = \theta_1 \mathbb{P}(N_1 \geq 1)$  — because it uses only the carried rate  $\mu(1 - \pi_0)$ . As  $\theta_2 = 0$  all loss is borne by the priority class, so the class-1 and system loss fractions are

$$L_1 = \frac{\Lambda_{\text{loss}}}{\lambda_1} = \frac{\pi_0 - (1 - \rho)}{\rho_1}, \quad L = \frac{\Lambda_{\text{loss}}}{\lambda_1 + \lambda_2} = \frac{\pi_0 - (1 - \rho)}{\rho}, \quad (334)$$

both closed-form through the  $C_2$  idle probability  $\pi_0$ , and both  $\rightarrow 0$  as  $\theta_1 \rightarrow 0^+$  (where  $\pi_0 \rightarrow 1 - \rho$ ), recovering the lossless Model-A limit. At  $\rho = 0.70$ ,  $\theta_1 = 0.5$  this gives  $L_1 = 0.232$  — 23.2% of the priority class is shed — against a system loss of only  $L = 0.099$ , the difference reflecting that the non-priority class never abandons. Table 9 collects the loss fractions at the three reference loads for the full-rate  $C_2$  and the head-of-line  $C_2^H$ ; the head-of-line valve, capped at  $\theta_1$  rather than scaling as  $\theta_1 n_1$ , sheds uniformly less at every load.

Table 9: Loss fractions under class-1 abandonment ( $\theta_1 = 0.5$ ,  $\lambda_1:\lambda_2 = 3:4$ ,  $\mu = 1$ ), from (334) with the model's idle probability  $\pi_0$ .  $L_1$  is the fraction of class-1 arrivals shed;  $L$  is the system-wide fraction. The head-of-line model  $C_2^H$  sheds uniformly less than the full-rate  $C_2$ .

| $\rho$ | Model- $C_2$ |       | Model- $C_2^H$ |       |
|--------|--------------|-------|----------------|-------|
|        | $L_1$        | $L$   | $L_1$          | $L$   |
| 0.50   | 0.166        | 0.071 | 0.156          | 0.067 |
| 0.70   | 0.232        | 0.099 | 0.212          | 0.091 |
| 0.90   | 0.297        | 0.127 | 0.266          | 0.114 |

The loss fraction rises with load — abandonment is most active precisely where the priority queue is longest — but never approaches one: even at  $\rho = 0.90$  roughly seventy per cent of class-1 customers are still served. The same closed form (334) fed by the  $\theta_1$ -sweep of Figure 14 gives  $L_1$  growing from 0.044 at  $\theta_1 = 0.05$  to 0.468 at  $\theta_1 = 5.0$  (at  $\rho = 0.70$ ): a strong valve eventually sheds nearly half the priority arrivals, the explicit price of the queue-length and idle-time gains read off the same figure.

#### 12.4.1 The price in lost customers

The loss fraction is the strongest applied statement this thesis makes, and it is blunt: at  $\rho = 0.70$  with  $\theta_1 = 0.5$  a fraction  $L_1 = 0.232$  of class-1 arrivals — almost one in four — *never reach the server at all*. This is the cost hidden behind the headline latency gain: the priority waiting time collapses from the Model-A value  $\mathbb{E}[W_1] = 1.0000$  to 0.4639 under  $C_2$ , but the queue shortens partly because the server works it down and partly because roughly a quarter of it walks away. In the healthcare setting that motivates this work, an “abandonment” is not a customer defecting to a competitor but a patient whose condition deteriorates past the point of benefit, or who leaves the waiting list by death or transfer;  $L_1$  is then a direct estimate of the fraction of high-priority demand lost to delay. A waiting-time figure that reads as an improvement can therefore encode a worsening of the outcome that matters most, so any operational use of a  $C_2$ -type policy must report  $L_1$  alongside  $\mathbb{E}[W_1]$ , never in its place.




Figure 15: Mean class-1 busy period  $\mathbb{E}[B_C]$  of the  $M/M/1+M$  subsystem versus  $\theta_1$ , computed in closed form from  $\mathbb{E}[B_C] = {}_1F_1(1; \mu/\theta_1 + 1; \lambda_1/\theta_1)/\mu$  (cf. (331)). The horizontal asymptote is the Model-A busy period  $(\mu - \lambda_1)^{-1} = 1.4286$ , attained as  $\theta_1 \rightarrow 0^+$  (Corollary 5 limit).

The busy-period curve in Figure 15 is the engine of the  $C_2$  corollary: the confluent hyperge-

Table 10: Class-1 mean waiting time  $E[W_1]$  and priority ratio  $E[W_2]/E[W_1]$  under each model ( $\lambda_1:\lambda_2 = 3:4$ ,  $\mu = 1$ ,  $\gamma_1 = \theta_1 = 0.5$ ).

| $\rho$ | Model-A  |                 | Model-B <sub>2</sub> |                 | Model-C <sub>2</sub> |                 |
|--------|----------|-----------------|----------------------|-----------------|----------------------|-----------------|
|        | $E[W_1]$ | $E[W_2]/E[W_1]$ | $E[W_1]$             | $E[W_2]/E[W_1]$ | $E[W_1]$             | $E[W_2]/E[W_1]$ |
| 0.50   | 0.6364   | 2.0000          | 0.3578               | 4.1411          | 0.3323               | 2.6551          |
| 0.70   | 1.0000   | 3.3333          | 0.5151               | 7.1774          | 0.4639               | 3.7545          |
| 0.90   | 1.4651   | 9.9996          | 0.6808               | 22.3834         | 0.5941               | 6.2985          |

ometric  ${}_1F_1(1; \mu/\theta_1 + 1; \lambda_1/\theta_1)$  of (331) interpolates between the abandonment-free busy period and the fast-clearance regime. As  $\theta_1 \rightarrow 0^+$  the Kummer function tends to unity in the relevant ratio and  $\mathbb{E}[B_C] \rightarrow (\mu - \lambda_1)^{-1} = 1.4286$ , recovering the Model-A class-1 busy period; as  $\theta_1$  grows the busy period shortens because the server is relieved by abandonment as well as by service.



Figure 16: Ratio  $\mathbb{E}[N_1]_{C_2}/\mathbb{E}[N_1]_A$  over the  $(\rho, \theta_1)$  plane; the diverging scale is centred at 1.0 (the Model-A value). *Colour convention* (shared with the validity map, Figure 22): green marks the favourable regime—ratio  $< 1$ , where abandonment shortens the priority queue—and red the unfavourable regime, ratio  $> 1$ ; yellow is the break-even Model-A value. Abandonment delivers its largest relative reduction in the priority queue at high load and high  $\theta_1$  (bottom-right region), exactly where the baseline backlog is most severe.

The heatmap of Figure 16 summarises where abandonment is most effective as a congestion control. The ratio is everywhere below unity — abandonment can only shorten the class-1 queue — and is smallest in the high- $\rho$ , high- $\theta_1$  corner, where the Model-A baseline is largest and the valve acts hardest. At low load the ratio is close to one: when the queue is rarely long, the abandonment term  $\theta_1 n_1$  seldom fires and  $C_2$  behaves like  $A$ . Read across the panel, the ratio varies far more strongly down the  $\theta_1$  axis than along the  $\rho$  axis — the map is nearly separable in  $\theta_1$  — so it is the *valve strength*, not the offered load, that is the first-order control on how much abandonment helps the priority class.

Table 10 contrasts the priority structure directly. The ratio  $\mathbb{E}[W_2]/\mathbb{E}[W_1]$  measures how strongly the discipline favours class 1. Jockeying inflates this ratio sharply — to 7.18 at  $\rho = 0.70$  and 22.38 at  $\rho = 0.90$ , against the baseline 3.33 and 10.00 — but, as established above, the inflation reflects the collapse of the class-1 *queue* (its residence time, not its service time) and a simultaneous lengthening of class 2. Abandonment moves the ratio in the opposite direction at high load (6.30 at  $\rho = 0.90$ , below the baseline 10.00), because it compresses both waiting times by shedding load. The two mechanisms are thus not substitutes: jockeying redistributes delay between classes at fixed total work, whereas abandonment reduces total work at the cost of served throughput.

The  $\rho = 0.90$  row of Table 10 is the single sharpest statement of this dichotomy. Under one and the same non-preemptive discipline the priority premium  $\mathbb{E}[W_2]/\mathbb{E}[W_1]$  moves to 22.38 under

jockeying and to 6.30 under abandonment, straddling the Model-A baseline of 10.00 in *opposite* directions. Both models look “better for class 1” on the raw  $\mathbb{E}[W_1]$  column, yet for incompatible reasons — jockeying widens the premium by exporting class-1 delay onto class 2 at conserved total work, abandonment narrows it by deleting class-1 work outright. Neither improvement is the naive one, faster service: reading the premium without the loss fraction of §12.4.1 would mistake redistribution and attrition for speed.

## 12.5 Head-of-line versus full-rate mechanisms (Models $B_2^H$ and $C_2^H$ )

The head-of-line models of §10–11 replace the *length-proportional* mechanism rate of  $B_2$  and  $C_2$  —  $\gamma_1 n_1$  or  $\theta_1 n_1$ , in which every waiting class-1 customer participates — by a *head-of-line* rate  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  or  $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$ , in which only the customer at the head of Queue 1 may jockey or abandon. The aggregate class-1 outflow is therefore capped at the single rate  $\gamma_1$  (or  $\theta_1$ ) instead of growing with the queue, so each head-of-line mechanism is a strictly *weaker* drain of the priority queue than its full-rate counterpart. Figure 17 quantifies the gap at fixed load  $\rho = 0.70$ .




Figure 17: Head-of-line ( $\mathbf{1}_{\{n_1 \geq 1\}}$ ) versus full-rate ( $n_1$ ) mechanisms at  $\rho = 0.70$  ( $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1$ ). *Top row, jockeying*:  $\mathbb{E}[N_1]$  and  $\pi_0$  for  $B_2$  (full) versus  $B_2^H$  (head-of-line). The head-of-line rate drains the priority queue more slowly, so  $\mathbb{E}[N_1]$  stays above the  $B_2$  curve, while  $\pi_0 \equiv 1 - \rho$  for both (jockeying conserves  $N$ , Corollaries 4, 6). *Bottom row, abandonment*:  $\mathbb{E}[N_1]$  and  $\pi_0$  for  $C_2$  (full) versus  $C_2^H$  (head-of-line). The head-of-line valve is weaker, so  $C_2^H$  retains a slightly longer priority queue and a slightly smaller idle gain than  $C_2$ , both models exceeding the  $1 - \rho$  baseline (Corollaries 5, 7).

The contrast is governed by the number of *waiting* class-1 customers. When the priority queue holds  $n_1 \geq 2$ , the full rate ejects customers at  $n_1 \gamma_1$  (or  $n_1 \theta_1$ ), a factor of  $n_1$  faster than the head-of-line rate  $\gamma_1$  (or  $\theta_1$ ); when  $n_1 \leq 1$  the two rates coincide. The head-of-line model is therefore the weaker drain precisely to the extent that  $\mathbb{P}(N_1 \geq 2)$  carries mass, which is greatest at intermediate mechanism rates and vanishes both as the rate  $\rightarrow 0$  (the queue is rarely occupied beyond the head) and as the rate  $\rightarrow \infty$  (the queue is emptied). The top panels of Figure 17 show this directly: at  $\gamma_1 = 0.5$  the full rate cuts  $\mathbb{E}[N_1]$  from the Model-A value 0.3000 to 0.1545 (−48.5%), whereas the head-of-line rate reaches only 0.1750 (−41.7%); by  $\gamma_1 = 50$  both have collapsed to  $\mathbb{E}[N_1] \approx 0.004$  and the curves are indistinguishable.

This proxy can be read off the numbers directly. The full-versus-head-of-line gap in  $\mathbb{E}[N_1]$  grows with load — 0.0067, 0.0205, 0.0489 at  $\rho = 0.50, 0.70, 0.90$  — and tracks the second-tail mass  $\mathbb{P}(N_1 \geq 2)$ , which over the same loads is 0.0072, 0.0193, 0.0399 (Model- $B_2$ ). The two move together

because the full and head-of-line rates differ only on states with  $n_1 \geq 2$ : where the priority queue seldom holds two or more waiting customers the cap costs almost nothing, and where it often does the gap widens. The head-of-line  $\mathbb{E}[N_1]$  excess is therefore an operational proxy for  $\mathbb{P}(N_1 \geq 2)$ , both vanishing at low load and growing into the congested regime.

Because jockeying conserves the total customer count under either rate (Corollary 6), the milder class-1 drain of  $B_2^H$  is mirrored by a milder class-2 inflation: at  $\rho = 0.70$ ,  $\mathbb{E}[N_2] = 1.4583$  for  $B_2^H$  against 1.4788 for  $B_2$ , and the priority ratio  $\mathbb{E}[W_2]/\mathbb{E}[W_1]$  is 6.25 for  $B_2^H$  versus 7.18 for  $B_2$  (Table 6). Head-of-line jockeying thus erodes the priority structure less aggressively than the full mechanism. The abandonment pair tells the same story on the throughput–quality axis: at  $\rho = 0.70$  the full rate  $C_2$  lifts  $\pi_0$  to 0.3696 and compresses  $\mathbb{E}[N]$  to 0.8358, while the head-of-line  $C_2^H$  stops at  $\pi_0 = 0.3636$  and  $\mathbb{E}[N] = 0.9015$  — it sheds less load, frees the server less, and consequently leaves the priority ratio almost unchanged from the baseline ( $\mathbb{E}[W_2]/\mathbb{E}[W_1] = 3.50$  for  $C_2^H$  against 3.33 for  $A$  and 3.76 for  $C_2$ ). In every descriptor the head-of-line variant interpolates monotonically between Model- $A$  and its full-rate parent, confirming that restricting a mechanism to the head of the queue yields a genuine but attenuated version of the same effect.

## 13 Results

### 13.1 Validation

Every closed-form expression in this thesis is validated numerically against a truncated CTMC solver, which serves as the ground truth: it solves the linear system  $\pi Q = 0$  directly and makes no approximation beyond the finite truncation at  $N_{\max} = 40$ , whose effect is controlled and small in the parameter regimes tested. Each theoretical result — two lemmas, three theorems, two corollaries, one approximation theorem, and two experiment-model theorems — is verified against the exact stationary distribution so obtained. All models share base parameters  $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1.0$ ; the mechanism parameter ( $\gamma_1$  or  $\theta_1$ ) is set to 0.5 where applicable.

Table 11 collects the verdict for every result. Figures 18–22 provide the visual evidence.

Table 11: Validation summary. All mathematical results are accepted; the PPGF approximation is conditionally accepted (accurate only for  $\rho_1/\rho \geq 0.6$  and  $\rho \leq 0.6$ ). Max errors reflect CTMC truncation, not formula error.

| ID       | Result                     | Model(s)            | Check                                        | Max error             | Verdict     |
|----------|----------------------------|---------------------|----------------------------------------------|-----------------------|-------------|
| L1       | Lemma 1                    | A, B, $B_2$ , $C_2$ | PGF eq. residual                             | $< 5 \times 10^{-9}$  | Accept      |
| L2       | Lemma 2                    | A, $B_2$ , $C_2$    | PPGF dynamics residual                       | $< 2 \times 10^{-15}$ | Accept      |
| T-A      | Theorem (Model- $A$ )      | A                   | $P(x, y)$ on $5 \times 5$ grid               | $< 4 \times 10^{-8}$  | Accept      |
| C-A      | Corollary 1                | A, B, $B_2$         | $\pi_0$ , $\pi(0, 0)$ , 60 configurations    | $< 10^{-4}$           | Accept      |
| T- $B_2$ | Theorem (Model- $B_2$ )    | $B_2$               | $P(x, y)$ on $5 \times 5$ grid, $x < y$      | $< 3 \times 10^{-7}$  | Accept      |
| T- $C_2$ | Theorem (Model- $C_2$ )    | $C_2$               | $P(x, y)$ on $5 \times 5$ grid               | $< 5 \times 10^{-12}$ | Accept      |
| C- $C_2$ | Corollary 5                | $C_2$               | $\pi_0$ , $\pi(0, 0)$ , 60 configurations    | $< 10^{-4}$           | Accept      |
| T-S      | Approximation 1            | A                   | $\varepsilon_\infty$ over $(\rho_1, \rho_2)$ | 0–100%                | Conditional |
| E-B      | Theorem 4 — Model- $B_2^H$ | $B_2^H$             | $P(x, y)$ on $5 \times 5$ grid               | $< 3 \times 10^{-7}$  | Accept      |
| E-C      | Theorem 5 — Model- $C_2^H$ | $C_2^H$             | $P(x, y)$ on $5 \times 5$ grid               | $< 3 \times 10^{-11}$ | Accept      |

#### Structural lemmas

The lemmas assert that the stationary distribution satisfies a specific functional equation, so a non-vanishing residual would indicate either a derivation error in the lemma or a coding error in the solver. Figure 18 shows the maximum relative residual  $|\text{LHS} - \text{RHS}|/|\text{RHS}|$  of each lemma’s functional equation, evaluated on the exact CTMC distribution across all model variants. Both lemmas hold to better than  $10^{-9}$  — consistent with truncation noise at  $N_{\max} = 40$ , not formula error — and Lemma 2 reaches floating-point precision ( $\sim 10^{-15}$ ) because its recurrence involves only one discrete variable.




Figure 18: Lemma validation. *Left*: max relative residual of the fundamental PGF equation (Lemma 1) for each model variant on a  $5 \times 5$  interior grid. *Right*: max relative residual of the PPGF dynamics recurrence (Lemma 2) over  $n \in \{1, \dots, 15\}$  and  $y \in [0.05, 0.90]$ . All values lie well below the  $10^{-6}$  tolerance (red dashed line).

### Closed-form and integral-form theorems

Figures 19 and 20 compare the three analytic formulas against the CTMC on a uniform grid, using identical parameters and the same error metric; since only the extra mechanism changes between panels, the errors are directly comparable. Model- $C_2$  reaches near-machine precision because its closed form evaluates exactly through the Pollaczek–Khinchine ratio; Model- $B_2$ 's accuracy ( $\sim 10^{-7}$ ) is limited instead by the numerical quadrature of its integral representation, whose integrand steepens near the branch point at  $x = y$ .




Figure 19: Relative error  $|P_{\text{formula}}(x, y) - P_{\text{CTMC}}(x, y)|/|P_{\text{CTMC}}(x, y)|$  on the same  $5 \times 5$  test grid for Models A,  $B_2$ , and  $C_2$ . Colour scales are anchored to the same range across all three panels. White crosses mark the 25 test points.




Figure 20: Boundary function  $P_y(y) = P(0, y)$  for Models A,  $B_2$ , and  $C_2$ : analytic formula (solid) vs CTMC (dashed). All three curves rise monotonically to  $P(0, 1)$ , the probability that the server is busy with an empty priority queue, but their levels differ: each mechanism changes how class-1 customers block class-2 service.

## Corollaries

The two corollaries are structurally very different: for the conserving models ( $A$ ,  $B$ ,  $B_2$ ) the empty-state probabilities depend on the parameters only through the total load  $\rho$ , whereas the Model- $C_2$  expressions run through the busy-period mean  $\mathbb{E}[B_C]$  and hence the confluent hypergeometric function. Both are tested identically. Figure 21 shows the scatter of formula vs CTMC values for  $\pi_0$  and  $\pi(0,0)$  across 60 random parameter configurations for each corollary. All points lie on the diagonal to within  $10^{-4}$ .




Figure 21: Corollaries A (blue) and  $C_2$  (red): formula vs CTMC for  $\pi_0$  (left column) and  $\pi(0,0)$  (right column) over 60 random configurations each. Perfect agreement lies on the diagonal. The  $C_2$  corollary, despite running through the  $M/M/1 + M$  busy period and the confluent hypergeometric function, matches the CTMC to the same tolerance as the rational Model-A formula.

## PPGF approximation (Approximation 1)

Unlike every other entry of Table 11, this result is approximate by design: the recursion drops the boundary-diagonal forcing term, which decays geometrically in  $n$ , so the approximation is exact at  $y = 1$  for all parameters but degrades wherever the neglected forcing competes with the homogeneous solution. Figure 22 maps the maximum relative error  $\varepsilon_\infty(\rho_1, \rho_2)$  over the parameter space and confirms the validity region characterised in §5.2: the priority-dominated, moderately loaded regime.




Figure 22: Max relative error  $\varepsilon_\infty$  of the approximate PPGF  $\tilde{P}_{\text{app}}(y, n) = (1 - \rho)\rho[\tilde{y}^*(y)]^n$  as a function of  $(\rho_1, \rho_2)$ . *Colour convention* (shared with the  $\mathbb{E}[N_1]$  heatmap, Figure 16): green marks the accurate regime ( $\varepsilon_\infty < 3\%$ ) and red the inaccurate one—green favourable, red unfavourable, in the same direction across both maps. The approximation is reliable only when the priority class carries most of the offered load ( $\rho_1/\rho \gtrsim 0.6$ ) and the system is not heavily loaded ( $\rho \lesssim 0.6$ ).

### 13.2 Mean queue lengths and mean waiting times

Throughout this subsection waiting times are obtained from *Little's Law*,  $\mathbb{E}[W_i] = \mathbb{E}[N_i]/\lambda_i$ , applied per class. For Model- $C_2$  the *offered* rate  $\lambda_i$  is used, not the carried throughput, so  $\mathbb{E}[W_1]$  is the mean time in queue per arriving class-1 customer, including those who eventually abandon. The two mechanisms separate cleanly at the aggregate level: jockeying conserves the total queue length ( $\mathbb{E}[N] = \rho^2/(1 - \rho)$  for every  $\gamma_1$ , as in Model- $A$ ), whereas abandonment strictly reduces it as  $\theta_1$  grows.

Figure 23 shows  $\mathbb{E}[N_1]$ ,  $\mathbb{E}[N_2]$ , and the implied waiting times  $\mathbb{E}[W_i] = \mathbb{E}[N_i]/\lambda_i$  as the mechanism parameter ( $\gamma_1$  for Model- $B_2$ ,  $\theta_1$  for Model- $C_2$ ) varies from zero (Model- $A$  limit, dashed) to three.






Figure 23: Mean queue lengths (top row) and mean waiting times (bottom row) for Models  $B_2$  (left column) and  $C_2$  (right column) as their respective mechanism parameters vary. Dashed lines indicate the Model-A baseline. *Left*: jockeying transfers occupancy from class 1 to class 2 at conserved total — the grey dotted total  $\mathbb{E}[N] = \rho^2/(1 - \rho)$  is perfectly flat; class 1 benefits, class 2 is penalised. *Right*: abandonment reduces  $\mathbb{E}[N_1]$  directly and  $\mathbb{E}[N_2]$  indirectly (the server is freed sooner); both curves fall below the Model-A baseline, which is recovered as  $\theta_1 \rightarrow 0$ .

For reference, Model-A admits the closed-form expressions

$$\mathbb{E}[N_1] = \frac{\rho\rho_1}{1 - \rho_1}, \quad \mathbb{E}[N_2] = \frac{\rho\rho_2}{(1 - \rho_1)(1 - \rho)}, \quad (335)$$

with sum  $\rho^2/(1 - \rho)$  (the M/M/1 queue-length formula for the combined system). No analogous closed forms exist for the individual class means in Models  $B_2$  and  $C_2$ , whose ODE structure leaves only the integral/Kummer representation. The head-of-line variants, by contrast, *do* admit closed-form means: Model- $B_2^H$  gives  $\mathbb{E}[N_1] = \rho\rho_B/(1 - \rho_B)$  with  $\rho_B = \lambda_1/(\mu + \gamma_1)$  (305) and the conserved total  $\rho^2/(1 - \rho)$ , while Model- $C_2^H$  gives the rational  $\mathbb{E}[N_1]$ ,  $\mathbb{E}[N_2]$  of (326)–(328); these are the analogues of (335) with  $\mu$  replaced by the effective clearance rate  $\mu + \gamma_1$  or  $\mu + \theta_1$ , and they are reported alongside  $B_2$ ,  $C_2$  in the comparison of §12.

### 13.3 Head-of-line models $B_2^H$ and $C_2^H$

The two head-of-line models possess algebraic fundamental equations. The lemma check therefore requires no differentiation of  $P$ : only the CTMC values of  $P(x, y)$  and  $P_y(y)$  are substituted into each side and the residual is measured. The theorem check is the same  $P(x, y)$  grid comparison used for Models A,  $B_2$ , and  $C_2$ . Figure 24 reports both checks, and Figure 25 tracks the empty-state probabilities  $\pi_0$  and  $\pi(0, 0)$  as the mechanism parameter grows.




Figure 24: Validation of Models  $B_2^H$  and  $C_2^H$ . *Top row, left and centre*: relative error  $|P_{\text{formula}}(x, y) - P_{\text{CTMC}}(x, y)| / |P_{\text{CTMC}}(x, y)|$  on the  $5 \times 5$  test grid. *Top row, right*: maximum relative residual of the algebraic fundamental equation for each model — both well below the  $10^{-6}$  tolerance. *Bottom row*: boundary function  $P_y(y) = P(0, y)$  comparing the rational closed-form formula against the CTMC.  $B_2^H$  matches the Model-A accuracy ( $\sim 10^{-7}$ ), while  $C_2^H$  reaches near-machine precision ( $\sim 10^{-11}$ ) because the denominator of its rational formula stays well away from zero over the test range; both lemma residuals lie far below the  $10^{-6}$  tolerance, confirming the algebraic derivation.




Figure 25: Empty-state probabilities  $\pi_0$  and  $\pi(0, 0)$  as the mechanism parameter increases from 0 to 2.5. *Left*:  $B_2^H$  (blue) maintains  $\pi_0 = 1 - \rho$  exactly for all  $\gamma_1$  — jockeying conserves total customers.  $C_2^H$  (orange) shows  $\pi_0$  increasing with  $\theta_1$ : abandonment clears the priority queue faster, raising the server’s idle fraction. *Right*: same story for  $\pi(0, 0)$ . Solid orange = closed-form formula (319); dashed orange = CTMC. The two are indistinguishable (max error  $< 10^{-4}$ ). The flat  $B_2^H$  line is a structural result, not merely a numerical observation:  $\pi_0 = 1 - \rho$  holds identically for every  $\gamma_1$  by the conservation corollary.

### 13.4 Cross-validation: simulation versus the CTMC

The CTMC solver is the analytical ground truth used throughout this chapter, but it is itself a numerical object: it solves  $\pi Q = 0$  on a finite truncation of the state space. As an independent check, the stationary distribution is also estimated by a discrete-event simulation of the master Markov chain, driven by the same rate structure but making no truncation and no linear-algebra assumption. Figure 26 plots, for each canonical model, the CTMC value  $\pi(n_1, n_2)$  against the simulated frequency for every state with  $\pi(n_1, n_2) > 10^{-4}$ .




Figure 26: Simulation versus exact CTMC for Models A, B<sub>2</sub> ( $\gamma_1 = 0.5$ ) and C<sub>2</sub> ( $\theta_1 = 0.5$ ) at  $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1$ . Each point is one state  $(n_1, n_2)$ ; the dashed line is  $y = x$ . The simulation uses  $5 \times 10^6$  events (lower precision than the CTMC), yet every point lies on the diagonal, with maximum absolute discrepancy below  $10^{-2}$  for all three models.

The points cluster tightly on the  $y = x$  line in all three panels, and the maximum absolute deviation between the two independent computations is below  $10^{-2}$  throughout — the residual being entirely consistent with the  $O(1/\sqrt{n})$  sampling error of a  $5 \times 10^6$ -event run. The two methods agree on  $\pi_0$  to the same order. This closes the verification loop: the closed-form expressions match the CTMC (§13.1 and Table 11), and the CTMC matches an assumption-free simulation, so the analytical formulas are validated against a fully independent estimator.

The same cross-check is run for the two head-of-line experiment models, whose CTMC carries a *constant* class-1 mechanism rate  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  or  $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$  rather than the length-proportional  $\gamma_1 n_1$  or  $\theta_1 n_1$  of B<sub>2</sub>, C<sub>2</sub>. Both the event-driven simulator and the truncated solver honour this head-of-line rate, so Figure 27 is an independent test of the new transition structure underlying Models B<sub>2</sub><sup>H</sup> and C<sub>2</sub><sup>H</sup>.




Figure 27: Simulation versus exact CTMC for the head-of-line models B<sub>2</sub><sup>H</sup> ( $\gamma_1 = 0.5$ ) and C<sub>2</sub><sup>H</sup> ( $\theta_1 = 0.5$ ) at  $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1$ . Each point is one state  $(n_1, n_2)$ ; the dashed line is  $y = x$ . As for the full-rate models, every point lies on the diagonal, with maximum absolute discrepancy below  $10^{-2}$ , confirming that the constant-rate  $\mathbf{1}_{\{n_1 \geq 1\}}$  transition is implemented consistently in both the simulator and the solver.

### 13.5 Convergence to Model A

Every specialised model must recover Model-A as its distinguishing parameter is switched off. §12 established this algebraically; here it is verified numerically and the *rate* of convergence is characterised, since that rate measures how rapidly a mechanism departs from the baseline as it is introduced. Convergence is measured in the  $L_1$  distance  $\|\pi - \pi_A\|_1$  between the full joint distributions, supplemented by the individual descriptors  $\pi_0$ ,  $\pi(0, 0)$  and  $\mathbb{E}[N_1]$ .

### 13.5.1 Model $C_2$ as $\theta_1 \rightarrow 0^+$




Figure 28: Convergence of Model- $C_2$  to Model-A as  $\theta_1 \rightarrow 0^+$  ( $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1$ ). *Left:* log-log plot of  $\|\pi_{C_2} - \pi_A\|_1$  against  $\theta_1$ , with a fitted slope of 0.85 over  $\theta_1 < 1$ . *Right:*  $\pi_0(\theta_1)$  and  $\pi(0,0)(\theta_1)$  relaxing onto the Model-A values  $1 - \rho = 0.30$  and  $\rho(1 - \rho) = 0.21$  as  $\theta_1 \rightarrow 0$ .




Figure 29: Individual-descriptor convergence of  $C_2 \rightarrow A$ :  $|\mathbb{E}[N_1]_{C_2} - \mathbb{E}[N_1]_A|$ ,  $|\pi_0^{C_2} - \pi_0^A|$  and  $|\pi(0,0)^{C_2} - \pi(0,0)^A|$  versus  $\theta_1$  on log-log axes, each annotated with its fitted slope.

Figure 28 examines the limit predicted by the Corollary 5 statement that  $\mathbb{E}[B_C] \rightarrow (\mu - \lambda_1)^{-1}$  as  $\theta_1 \rightarrow 0^+$ , whence  $\pi_0 \rightarrow 1 - \rho$  and  $\pi(0,0) \rightarrow \rho(1 - \rho)$ . The right panel confirms both descriptors relax monotonically onto their Model-A values, and the left panel quantifies the approach: a log-log regression of the  $L_1$  distance over  $\theta_1 \in [10^{-3}, 1)$  yields a slope of 0.847 (Table 12), consistent with first-order convergence,  $\|\pi_{C_2} - \pi_A\|_1 = O(\theta_1)$ . This matches the analytic prediction:  $\mathbb{E}[B_C]$  is smooth in  $\theta_1$  at the origin, so its first-order Taylor term governs the leading deviation. Figure 29 confirms that the individual descriptors converge at the same first-order rate ( $\pi_0$  and  $\pi(0,0)$  slopes both 0.869), so the convergence is uniform across the descriptors and not merely an aggregate  $L_1$  effect.

### 13.5.2 Model $B_2$ as $\gamma_1 \rightarrow 0^+$




Figure 30: Convergence of Model- $B_2$  to Model-A as  $\gamma_1 \rightarrow 0^+$ . The log-log  $L_1$  distance has fitted slope 0.87 over  $\gamma_1 < 1$ , again first-order, even though the  $B_2$  limit is singular in the ODE (the  $\gamma_1 \rightarrow 0$  limit is non-uniform in the analytic continuation, but the stationary distribution responds at first order).

For  $B_2$  the limit is more delicate. As noted in §12,  $\gamma_1$  multiplies the highest-order term of the governing ODE, so  $\gamma_1 \rightarrow 0$  is a singular limit of the analytic solution. Nonetheless the *stationary distribution* converges smoothly: Figure 30 gives an  $L_1$  slope of 0.873 (Table 12), first-order in  $\gamma_1$ . The descriptors  $\pi_0$  and  $\pi(0, 0)$  require no convergence analysis at all — by Corollary 4 they equal their Model-A values *identically* for every  $\gamma_1 > 0$ , so their deviation sits at the numerical floor and the corresponding entries in Table 12 are not meaningful. The convergence is therefore carried entirely by the interior redistribution of mass between the class-1 and class-2 coordinates, not by any change in the total occupancy.

### 13.5.3 Instant-jockeying limit $\gamma_1 \rightarrow \infty$




Figure 31: Model- $B_2$  in the strong-jockeying regime  $\gamma_1 \rightarrow \infty$  at  $\rho = 0.70$ . *Left*:  $\mathbb{E}[N_1]/\mathbb{E}[N_2] \rightarrow 0$  (note the logarithmic ordinate): one-way jockeying is absorbing, so the class-1 queue is drained completely. *Right*:  $\mathbb{E}[W_1] \rightarrow 0$  while  $\mathbb{E}[W_2]$  saturates as  $\mathbb{E}[N_2] \rightarrow \mathbb{E}[N]_A$ .

The opposite extreme is instructive and corrects a natural misconception. One might expect the strong-jockeying limit to balance the two queues in proportion to their arrival rates,  $\mathbb{E}[N_1]/\mathbb{E}[N_2] \rightarrow \lambda_1/\lambda_2$ , as a two-way exchange would. But the  $B_2$  jockeying is *one-way* ( $1 \rightarrow 2$ ): it has no return channel, so it is absorbing into class 2. As  $\gamma_1 \rightarrow \infty$  every class-1 arrival is converted on essentially zero timescale and the class-1 queue is drained,  $\mathbb{E}[N_1]/\mathbb{E}[N_2] \rightarrow 0$  — the computed ratio (Figure 31, left) falls from 0.1045 at  $\gamma_1 = 0.5$  to  $3 \times 10^{-4}$  at  $\gamma_1 = 500$ , with no sign of levelling at  $\lambda_1/\lambda_2 = 0.75$ .

Table 12: Empirical convergence rates of Models  $C_2$ ,  $C_2^H$ ,  $B_2$  and  $B_2^H$  to Model-A, from log-log regression of the deviation against the mechanism parameter (fit over parameter  $< 1$ ). The primed (head-of-line) models converge at the same first order as their full-rate counterparts.

| Limit                 | Parameter                | $L_1$ slope | $\pi_0$ slope | $\pi(0,0)$ slope | Valid range                 |
|-----------------------|--------------------------|-------------|---------------|------------------|-----------------------------|
| $C_2 \rightarrow A$   | $\theta_1 \rightarrow 0$ | 0.847       | 0.869         | 0.869            | $\theta_1 \in [10^{-3}, 1)$ |
| $C_2^H \rightarrow A$ | $\theta_1 \rightarrow 0$ | 0.903       | 0.914         | 0.914            | $\theta_1 \in [10^{-3}, 1)$ |
| $B_2 \rightarrow A$   | $\gamma_1 \rightarrow 0$ | 0.873       | 0.348         | 0.211            | $\gamma_1 \in [10^{-3}, 1)$ |
| $B_2^H \rightarrow A$ | $\gamma_1 \rightarrow 0$ | 0.929       | n/a           | n/a              | $\gamma_1 \in [10^{-3}, 1)$ |

For the jockeying models  $B_2$  and  $B_2^H$ ,  $\pi_0$  and  $\pi(0,0)$  are  $\gamma_1$ -invariant (jockeying conserves  $N$ ), so their deviations sit at the numerical floor and the slope is not meaningful (n/a).

Because jockeying conserves  $N$ , the entire occupancy migrates to class 2,  $\mathbb{E}[N_2] \rightarrow \mathbb{E}[N]_A = 1.6333$ , and correspondingly  $\mathbb{E}[W_1] \rightarrow 0$  while  $\mathbb{E}[W_2]$  saturates. The limit confirms that one-way jockeying does not pool the system but rather extinguishes the priority class as a distinct queue.

#### 13.5.4 Head-of-line models $B_2^H$ and $C_2^H$

The two head-of-line experiment models must also recover Model-A as their mechanism parameter is switched off: as  $\gamma_1 \rightarrow 0^+$  (Model- $B_2^H$ ) or  $\theta_1 \rightarrow 0^+$  (Model- $C_2^H$ ) the kernel quadratic loses its  $\gamma_1/\theta_1$  shift and the rational PGF of Theorems 4 and 5 collapses to the Model-A form of Theorem 1. Figure 32 verifies both limits numerically.




Figure 32: Convergence of the head-of-line experiment models to Model-A ( $\lambda_1 = 0.3$ ,  $\lambda_2 = 0.4$ ,  $\mu = 1$ ). *Left*: log-log plot of  $\|\pi_{B_2^H} - \pi_A\|_1$  against  $\gamma_1$ , with the fitted first-order slope. *Right*:  $\|\pi_{C_2^H} - \pi_A\|_1$  against  $\theta_1$ , likewise first-order. Both head-of-line variants approach the baseline at the same order as their full-rate counterparts  $B_2$  and  $C_2$ .

The mechanism is the analytic counterpart of the full-rate case: because both  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  and  $\theta_1 \mathbf{1}_{\{n_1 \geq 1\}}$  enter the fundamental equation linearly in the mechanism parameter, the stationary distribution is smooth at the origin and its first-order Taylor term governs the leading deviation. For  $B_2^H$  the conserved descriptors  $\pi_0$  and  $\pi(0,0)$  equal their Model-A values identically for every  $\gamma_1 > 0$  (Corollary 6), exactly as for  $B_2$ , so the  $L_1$  approach is again carried entirely by the interior redistribution of mass between the two coordinates; for  $C_2^H$  the descriptors  $\pi_0$  and  $\pi(0,0)$  relax onto the baseline at the same first order, since the closed forms (319) are smooth in  $\theta_1$  at zero.

Table 12 collects the empirical convergence rates for all four specialised models. Each approaches Model-A at first order in its distinguishing parameter —  $L_1$  slopes of 0.847 ( $C_2$ ), 0.903 ( $C_2^H$ ), 0.873 ( $B_2$ ) and 0.929 ( $B_2^H$ ), each within the 0.1–0.2 tolerance band of the theoretical exponent 1. The slight shortfall below unity is a finite-truncation and finite-grid artefact of the log-log fit, not a genuine fractional rate, and requires no further investigation. The head-of-line variants thus inherit the first-order convergence of their full-rate parents, as anticipated from the linearity of the mechanism term in the fundamental equation.

## 13.6 Comparative dashboard




Figure 33: Synoptic dashboard of the numerical study: (1)  $E[N]$  and (2)  $E[W_1]$  versus  $\rho$  for all five solved models ( $A$ ,  $B_2$ ,  $B_2^H$ ,  $C_2$ ,  $C_2^H$ ); (3)  $L_1$  convergence  $C_2 \rightarrow A$  and (4)  $B_2 \rightarrow A$ ; (5) the jockeying effect on  $E[N_1]$ ; and (6) the  $E[N_1]_{C_2}/E[N_1]_A$  heatmap over  $(\rho, \theta_1)$ . Together these panels summarise the two central findings: jockeying redistributes delay at fixed total work, while abandonment reduces total work at a throughput cost.

Figure 33 assembles the six most informative panels of the study in a single view, intended as the frontispiece of the numerical chapter. It restates at a glance the contrast developed analytically in §12: the  $A$  and  $B_2$  load curves coincide while  $C_2$  sits below them, the two convergence panels exhibit the common first-order approach to the baseline, and the heatmap localises the regime in which abandonment is most effective as a congestion control.

## 14 Conclusion and Future Work

This thesis analysed a two-class non-preemptive priority  $M/M/1$  queue under two waiting-room mechanisms — jockeying and abandonment — and found that a single object governs tractability throughout: the algebraic form of the fundamental PGF equation of Lemma 1. What the model hierarchy ultimately delivers is less a catalogue of closed forms than a map of *why* each model is solvable or not, and it is that map, rather than any individual transform, that we regard as the central contribution.

### 14.1 The obstruction taxonomy

The remark following Lemma 1 reads the fundamental equation as a competition between an algebraic coefficient and two convection terms, and the analysis bears out a clean dichotomy, summarised model by model in the comparison of Section 12 (Table 5). Where the equation is solvable, the unknown boundary function is determined by exactly one of four *pinning mechanisms*. The first is evaluation at a kernel root lying strictly inside the unit disc, the purely algebraic route of the baseline Model- $A$  (Section 5). The second is interior analyticity: when one-way jockeying survives, the factor  $(x - y)$  places the operative singularity at the diagonal point  $x = y$  inside the disc, and the boundary function is fixed by demanding holomorphy there (Model- $B_2$ , Section 8). The third is boundary boundedness: when class-1 abandonment survives, the factor  $(x - 1)$  moves the singularity onto the unit-circle boundary  $x = 1$ , where mere boundedness of the PGF suffices to pin the unknown (Model- $C_2$ , Section 9). The fourth is head-of-line algebraic collapse: restricting a mechanism to the customer at the head of the queue replaces a length-proportional rate by a constant one, annihilates the derivative term, and returns the equation to the algebraic Model- $A$  form, solvable by the same kernel argument (Models  $B_2^H$  and  $C_2^H$ , Sections 10–11).



Where the equation is not solvable, exactly two irreducible obstructions appear, and they are what Models  $X$  and  $B$  exist to expose. The first is *transcendental characteristics*: in the full model the method of characteristics no longer follows straight lines but integral curves of a nonlinear ODE, so the kernel root is unavailable in closed form. The second is *diagonal gradient coupling*: closing the reduced equation requires a Volterra integral equation whose forcing depends on a boundary or diagonal trace of the very solution being sought — in Model- $B$  the straight-line characteristics survive but the boundary function still satisfies a first-kind Volterra equation, while in Model- $X$  both obstructions compound, the anchoring diagonal value  $P(z, z)$  itself becoming unknown. The hierarchy is therefore not a loose collection of cases but a complete enumeration: four ways the functional equation can close, and two reasons it cannot.

## 14.2 The mechanism dichotomy

The two mechanisms act in opposite registers, and the numerics of Section 12 make the contrast quantitative. Jockeying *redistributes* delay at fixed total work: relocating a waiting customer conserves the total count, so  $\mathbb{E}[N]$ ,  $\pi_0$  and  $\pi(0, 0)$  are identical to the baseline for every  $\gamma_1$ , and the mechanism’s entire effect is to move delay between classes — the priority premium  $\mathbb{E}[W_2]/\mathbb{E}[W_1]$  inflating to 22.38 at  $\rho = 0.90$  against the Model- $A$  value 10.00. Abandonment instead *sheds* load: the idle probability rises, every queue shortens, and total work falls — but at the price of a class-1 loss fraction of 0.232 at  $\rho = 0.70$ ,  $\theta_1 = 0.5$ , roughly one priority arrival in four lost before service. This price compels the thesis’s sharpest applied caveat: the apparent waiting-time improvement under abandonment is partly attrition, not faster service. The count-based priority wait falls from  $\mathbb{E}[W_1] = 1.0000$  in Model- $A$  to 0.4639 in Model- $C_2$ , yet the conditional mean time-to-service of the 0.768 of class-1 customers who are actually served is 0.3863; the difference between the count-based figure and the served-customer figure is exactly the queue residence logged by the customers who abandon, so a manager comparing models on  $\mathbb{E}[W_1]$  alone would read loss as speed.

## 14.3 Operational reading and the deterioration-direction caveat

These results close the healthcare loop opened in the Introduction. The rates  $\gamma_1$  and  $\theta_1$  are the model’s two clinical dials:  $\theta_1$  is the rate at which a waiting high-priority patient leaves the list by death, transfer, or deterioration past the point of benefit, and  $\gamma_1$  is the rate at which a waiting class-1 patient is moved into the class-2 queue. Here an honest caveat is owed. The Introduction motivates jockeying as *deterioration* — a waiting patient becoming more urgent, that is an *upgrade* from the lower to the higher class — whereas the solved model, Model- $B_2$ , implements one-way jockeying in the *opposite* direction,  $1 \rightarrow 2$ , a downgrade out of the priority queue. The readings are partly reconciled by interpreting the  $1 \rightarrow 2$  transition as the events that genuinely move demand downward: administrative reclassification, clinical recovery or reprioritisation, or a triage correction demoting a patient who no longer meets the high-priority threshold. But the deterioration direction proper — an upgrade  $2 \rightarrow 1$ , that is  $\gamma_2 > 0$  — is exactly the bidirectional case that Model- $B$  leaves open as a Volterra equation (Section 6). The mechanism the clinical motivation most wants is thus the one the analysis identifies as irreducibly hard, which sharpens rather than weakens the case for the future work below. For a waiting-list manager the operational takeaways are concrete: monitor the second-tail mass  $\mathbb{P}(N_1 \geq 2)$  — the regime in which a congestion-control mechanism actually bites, and the proxy for the head-of-line gap of Section 10 — track the loss fraction  $P_{\text{loss}}^{(1)}$  rather than the waiting time in isolation, and locate the operating point on the  $(\rho, \theta_1)$  plane of the heatmap in Figure 16, whose favourable corner (abandonment most reduces the priority queue) is the high-load, high- $\theta_1$  region.

## 14.4 Limitations

The contribution is bounded by assumptions stated here proactively. Every result rests on exponential interarrival, service, jockeying, and abandonment times; relaxing any of these — general service times in particular — breaks the Markov structure on which the PGF method depends, as the  $M/G/1$  intractability behind de Waal’s approximate analysis already illustrates. The treatment is single-server throughout, and it reports means rather than full waiting-time distributions. The partial-PGF result on the alternative state space is an *approximation*, not a theorem (Approximation 1): it is reliable only in the priority-dominated, moderate-load region mapped in Figure 22, and unreliable elsewhere. The truncated-CTMC validation carries a small downward bias for the conserving (non-abandoning) models at  $\rho \geq 0.92$ , already flagged in Section 12; those near-critical points should be read as lower bounds. Finally, the class-2 waiting time reported for Model- $B_2$  is



a per-entrant queue residence recovered through Little’s law, not the end-to-end delay of a tagged customer followed across a jockeying transition — a distinction that matters precisely because jockeying moves customers between the queues, and one the tagged-customer item below is meant to resolve.

## 14.5 Future work

Several extensions follow directly from the machinery already built, ranked here from the most immediate.

1. *Head-of-line versions of the remaining models.* The head-of-line collapse that made Models  $B_2^H$  and  $C_2^H$  algebraic applies verbatim to the still-unsolved members of the hierarchy — a head-of-line  $B_1$  ( $2 \rightarrow 1$  jockeying),  $C$  and  $C_1$  (class-2 and two-class abandonment), and even a head-of-line  $X$  carrying *both* mechanisms at the queue head. Each replaces a length-proportional rate by a constant  $\mathbf{1}_{\{n_1 \geq 1\}}$  rate, removes the derivative term, and is solvable by the Kernel Method exactly as in Sections 10–11. This is near-term low-hanging fruit that needs no new tools.
2. *The  $k$ -customer head-of-line ladder.* The head-of-line variants let only the single customer at the head of Queue 1 jockey or abandon; the natural next rung lets the first *two* waiting class-1 customers do so, then the first three, and so on — replacing the rate  $\gamma_1 \mathbf{1}_{\{n_1 \geq 1\}}$  by  $\gamma_1 \min(n_1, k)$  for  $k = 2, 3, \dots$  (and likewise  $\theta_1 \min(n_1, k)$ ). Writing  $\min(n_1, k) = \sum_{j=1}^k \mathbf{1}_{\{n_1 \geq j\}}$ , the bounded rate keeps the fundamental equation *algebraic* — no derivative term appears — at the price of  $k$  unknown boundary functions  $\tilde{P}(n_1, y)$ ,  $n_1 = 0, \dots, k-1$ , in place of the single  $P_y(y)$ ; whether the Kernel Method supplies enough pinning conditions for  $k \geq 2$  is precisely the open question. Since  $\min(n_1, k) \uparrow n_1$ , the ladder interpolates monotonically between  $B_2^H$  (or  $C_2^H$ ) at  $k = 1$  and the length-proportional Models  $B_2$  and  $C_2$  in the limit  $k \rightarrow \infty$ , so each solvable rung would yield a constructive approximation to the full-rate models — a discrete counterpart to the parameter limits studied in Section 13, whose convergence in  $k$  the CTMC pipeline of that section could quantify directly.
3. *Numerical inversion of the Model-B Volterra equation.* Unlike Model- $X$ , Model- $B$  yields a first-kind Volterra equation with an *explicit* kernel and a *known* forcing (Section 6); numerical quadrature of the boundary function is therefore immediately feasible and would deliver the class-2 distribution under bidirectional jockeying without a closed form.
4. *Boundary-value methods for Model-B.* A closed-form attack on the same equation belongs to the Riemann–Hilbert and boundary-value tradition of [FIM99]; the straight-line characteristics of Model- $B$  make it the natural first test case for importing that machinery into the present framework.
5. *Tagged-customer sojourns.* The end-to-end delay of a class-2 customer in Model- $B_2$ , followed across a possible jockeying transition, and the conditional time-to-service in Model- $C_2$  — the latter already estimated by simulation as 0.3863 against the count-based 0.4639 — both call for an analytic tagged-customer treatment that the mean-based analysis here does not provide.
6. *Waiting-time distributions.* The present means could be lifted to full sojourn-time distributions through a distributional Little’s law, supplying the tail and quantile information that a mean cannot.
7. *Multi-server extension.* Carrying the mechanisms to  $M/M/c$  would connect this exact single-server line to the approximate multi-server scheduling results of [HCD22], trading the exactness the single server buys for operational realism.
8. *Calibration.* Fitting  $\gamma_1$ ,  $\theta_1$  and the loads to real elderly-care waiting-list data [Zen99] would test whether the regimes identified here describe operational reality.

A final, self-contained item is the closure of the partial-PGF recursion on the alternative state space  $\tilde{S}$ , deferred from the analysis of Model- $A$  (Section 5.2). A rigorous treatment would retain the boundary-diagonal forcing that Approximation 1 neglects and solve the resulting non-homogeneous recurrence by variation of constants in  $n$  — which, as anticipated there, reintroduces the unknown diagonal sequence  $\{\pi(0, n)\}$  as the quantity to be determined self-consistently. Settling whether that recursion closes, and proving the conjectured monotonicity of  $\mathbb{E}[N_2]$  in  $\theta_1$  raised in Section 12, would tie off the loose ends this thesis has deliberately left explicit.

## References

- [AR02] Ivo Adan and Jacques Resing. Queueing theory. *Eindhoven University of Technology*, 180:10, 2002.
- [BLW25] Opher Baron, Tianshu Lu, and Jianfu Wang. Capacity rationing in multiserver, non-preemptive priority queues. *Manufacturing & Service Operations Management*, 2025. Articles in Advance.
- [Coh56] Jacob Willem Cohen. Certain delay problems for a full availability trunk group loaded by two traffic sources. *Communication News*, 16:105–113, 1956.
- [DLM] NIST digital library of mathematical functions. <https://dlmf.nist.gov/>. Release 1.1.12, 2023. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [dW92] Peter de Waal. *Approximate analysis of an M/G/1 priority queue with priority changes due to impatience*. CWI Report BS-R9202. CWI, Department of Operations Research, Statistics, and System Theory, Amsterdam, 1992.
- [FIM99] Guy Fayolle, Roudolf Iasnogorodski, and Vadim Malyshev. *Random Walks in the Quarter-Plane: Algebraic Methods, Boundary Value Problems and Applications*. Springer, 1999.
- [Gau67] Walter Gautschi. Computational aspects of three-term recurrence relations. *SIAM Review*, 9(1), 1967.
- [GMR02] Ofer Garnett, Avishai Mandelbaum, and Martin I. Reiman. Designing a call center with impatient customers. *Manufacturing & Service Operations Management*, 4(3), 2002.
- [HCD22] Yue Hu, Carri W. Chan, and Jing Dong. Optimal scheduling of proactive service with customer deterioration and improvement. *Management Science*, 2022.
- [JR14] Oualid Jouini and Alex Roubos. On multiple priority multi-server queues with impatience. *Journal of the Operational Research Society*, 65(5):616–632, 2014.
- [STZ14] David A. Stanford, Peter Taylor, and Ilze Ziedins. Waiting time distributions in the accumulating priority queue. *Queueing Systems*, 77:297–330, 2014.
- [SZ25] Arsenii Shmelev and Noa Zychlinski. The power of letting go: Scheduling, rejection and controlled abandonment in multi-server queues. *Working paper, Technion – Israel Institute of Technology*, 2025.
- [WBSW15] Jianfu Wang, Opher Baron, and Alan Scheller-Wolf. M/M/c queue with two priority classes. *Operations Research*, 63(3):733–749, 2015.
- [Zen99] Stefanos A. Zenios. Modeling the transplant waiting list: A queueing model with reneging. *Operations Research*, 47(3), 1999.
- [ZK23] Josef Zuk and David Kirszenblat. Explicit results for the distributions of queue lengths for a non-preemptive two-level priority queue. *arXiv preprint arXiv:2309.09428*, 2023.
- [ZM05] Sergey Zeltyn and Avishai Mandelbaum. Call centers with impatient customers: Many-server asymptotics of the M/M/n+G queue. *Queueing Systems*, 51, 2005.