

Case Enteral Feeding (2)

The patient, Hanan, is 22 years old. She was admitted to the hospital for kidney failure in the last stage, for unknown reasons. It weight 62 kg, and is height 160, the patient has some clinical signs, pallor of the face, dry mouth and nausea slightly, and she is unable to eat food through the mouth due to her critical condition, difficulty breathing and the possibility of fluids in the lower parts, as shown by taking the patient's dietary history, increased intake of calories by 112%, as well as an increase in protein intake of 1.9 grams / kg, and the consumption of sodium in eating was also high, for the fluid intake. 500 ml/250-250 milliliters of juice and 480 milliliters of tea with lemon, some laboratory tests were conducted and the results were shown as follows:

Test	Result	Normal Rang	Interpretation
Hemoglobin	6.1	F: 11.5 -16.5 g/dl	
Creatinine	5.2	F: 0.6-0.9 mg/dl	
Sodium	135.0	135-145 mEq/L	
Potassium	4.5	3.6-5.0 mEq/L	
Calcium	7.5	8.5-10.5 mg/dl	
phosphate	3	0.8-1.5 mEq/L	
Glucose	81	65-100 mg/dl	

1. Determine the type of tubular nutritional administration?

.....

2. What is the appropriate nutritional intervention for this case, using the steps of the (NCP)?

Nutrition Care Process

1- nutrition assessment:

A- anthropometric measurement:

- ∂ Name:
- ∂ Age:
- ∂ Sex:
- ∂ Weight:
- ∂ Height:
- ∂ BMI:
- ∂ S.F:

B- biochemical test:

C- clinical features:

D – dietary history:

2- Nutritional diagnosis:

- Problem:
- Etiology:
- S|S:

3- Nutritional intervention:

- Goals:
- Estimated energy requirement:
- Nutrient requirement:

- Food exchange list:

Food item	Serving	CHO	protein	Fat	Calories
Milk					
Vegetable					
Fruit					
Bread					
Meat					
Fat					
Total					
Requirement					

Planning diet:

4- Monitoring and evaluation: