

Nutrients – Drug Interaction Table (<i>Diuretics</i>)					
Uses and Interactions	Furosemide	Indapamide	Triamterene	Budesonide	Metolazone
	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Absorption is decreased with meals or tube feeding - Monitor Potassium levels and intake - Used as diuretic and Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium & potassium excretion. - Used as diuretic and Antihypertensive agent.	- Avoid grapefruits - Avoid high fat food. - Used as diuretic.	- Monitor Potassium & Sodium levels and intake - Inhibits reabsorption of sodium & potassium. - Used as diuretic and Antihypertensive agent.
Interactions	Bumetanide	Hydrochlorothiazide	Chlorothiazide	Spironolactone	
	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium - Used as diuretic and Antihypertensive agent.	- Monitor levels and intake - Inhibits reabsorption of sodium - Used as diuretic and Antihypertensive agent.	- Avoid Potassium containing diet - Used as Potassium sparing diuretic.	

Nutrients – Drug Interaction Table (*Antihypertension drugs*)

Uses and Interactions	Valsartan	Nifedipine	Chlorothiazide	Chlorothiazide	Hydrochlorothiazide
	- Avoid salt substitutes containing K - Used as Antihypertensive agent.	- Avoid grapefruits. - Used as Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.
	Bumetanide	Captopril	Hydralazine	Felodipine	Altace
	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Absorption is decreased with meals or tube feeding - Avoid potassium sources. - Used as Antihypertensive agent.	- Monitor Vitamin B6 levels and intake, supplementation is recommended - Used as Antihypertensive agent.	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.	- Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.
	Fosinopril	Benazepril	Clonidine	Nicardipine	Diltiazem
	- Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.	- Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.	- Monitor glucose levels (Hyperglycemia) - Used as Antihypertensive agent.	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.

Nutrients – Drug Interaction Table (Antihypertension drugs)					
Uses and Interactions	Amlodipine	Triamterene	Furosemide	Indapamide	Quinapril
	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - & the potassium excretion. - Used as diuretic and Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Diuretic and Antihypertensive agent.	- Absorption is decreased with meals or tube feeding - Monitor Potassium levels and intake - Used as diuretic and Antihypertensive agent.	- Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.
	Enalapril	Moexipril	Ramipril	Nisoldipine	Nitrendipine
	- Avoid grapefruits - ACE inhibitors used as Antihypertensive agent.	- Avoid taking with meals or tube feeding - Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.	- Absorption is decreased with meals or tube feeding - Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.	- Avoid grapefruit - Avoid high fat meal or tube feeding - Calcium Channel Blockers Used as Antihypertensive agent.	- Avoid grapefruit - Calcium Channel Blockers Used as Antihypertensive agent.

Nutrients – Drug Interaction Table (sedatives)					
Uses and Interactions	Fentanyl	Midazolam	Phenobarbital	Primidone	Eszopiclone
	- Avoid grapefruits. - Used as sedative agents.	- Avoid grapefruits. - Used as sedative agents.	- Monitor Calcium, D, K intake & levels - Used as sedative agent & Anticonvulsant.	- Monitor Calcium, D, K intake & levels - Used as sedative agent & Anticonvulsant.	- Avoid high fat meal or with tube feeding. - Used as sedative & hypnotic (sleep medicine).
	Phenytoin	Carbamazepine	Triazolam		
	- Tube feeding decreases absorption (1-2 hr gap between TF & This medication & flush by 30 mL before & after drug administration) + Monitor Calcium, D, K intake & levels - Anti-convulsant, sedative used to treat Epilepsy.	- Avoid grapefruits - Monitor Iron & Sodium intake & Levels - Used as Anticonvulsant.	- Avoid grapefruits - Used as sedative agent & Antidepressant.		

Nutrients – Drug Interaction Table (Tube feeding)

Uses and Interactions	Phenytoin	Indapamide	Levothyroxine	Moexipril	Ramipril
	- Tube feeding decreases absorption (1-2 hr gap between TF & This medication & flush by 30 mL before & after drug administration) + Monitor Calcium, D, K intake & levels - Anticonvulsant, sedative used to treat Epilepsy.	-Absorption is decreased with meals or tube feeding -Avoid potassium sources. -Used as diuretic Antihypertensive agent.	-Absorption is decreased with meals or tube feeding: Should be taken one-half hour to one hour before. - Used to treat hypothyroidism.	- Absorption is decreased with meals or tube feeding - Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent.	-Absorption is decreased with meals or tube feeding - Avoid potassium sources. - ACE inhibitors used as Antihypertensive agent
	Minocycline	Pyrazinamide	Lansoprazole	Esomeprazole	Ciprofloxacin
	-Avoid dairy products. - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr. before or after - Used as tetracycline Antibacterial.	-Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). -Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr before or after. - Used as Antimycobacterial.	- Absorption is decreased with meals or tube feeding: Should be taken before - Proton pump inhibitor used as gastric acid inhibitor.	- Absorption is decreased with meals tube feeding : Should be taken 1 hour before -Proton pump inhibitor used as gastric acid inhibitor.	- Avoid with caffeine - Absorption is decreased with tube feeding: - Used as Anti-infective agent.
	Doxycycline	Omeprazole	Loratadine	Ampicillin	Ofloxacin
	- Avoid with folic acid containing Folate -Avoid dairy products. -Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr. before or after - Used as Antibiotic.	-Absorption is decreased with meals or tube feeding: Should be taken before - Proton pump inhibitor used as gastric acid inhibitor.	-Absorption is decreased with tube feeding. -Used as Antibiotic.	-Absorption is decreased tube feeding -Used as Antibiotic.	- Avoid dairy products - Absorption is decreased with tube feeding -Used as Anti-infective.

Diuretic. **Antihypertensive.** **Sedative.** **Tube feeding.** **Antibiotic,** **Anti-infective & Antibacterial.** **Antihyperlipidemic.** **Grapefruits.** **Sodium.** **Others**

Nutrients – Drug Interaction Table (Tube feeding)

	Nutrients – Drug Interaction Table (Tube feeding)				
	Eszopiclone	Cyclosporine	Sildenafil	Rifampin	Nisoldipne
Uses and Interactions	- Avoid high fat meal or with tube feeding. - Used as sedative & hypnotic (sleep medicine).	- Causes hyperglycemia - Avoid grapefruit - Avoid high fat meal or tube feeding - Immunosuppressant.	- Avoid grapefruits & high fat meal or tube feeding - Used as protease inhibitor.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). -Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr before or after. - Used as Antimycobacterial.	- Avoid grapefruit - Avoid high fat meal or tube feeding - Calcium Channel Blockers Used as Antihypertensive agent.
	Ibandronate sodium	Captopril	Tetracycline		
	- Absorption is decreased with meals or tube feeding: Should be taken 1 hr before - Avoid milk & mineral water - Used treat osteoporosis.	- Absorption is decreased with meals or tube feeding - Avoid potassium sources. - Used as Antihypertensive agent.	- Avoid dairy products. - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr. before or after- - Used as tetracycline Antibacterial.		

Nutrients – Drug Interaction Table (Antibiotics, Anti-infectives & Antibacterials)

Uses and Interactions	Levofloxacin	Penicillin	Cholestyramin	Ethambutol	Isoniazid
	- Avoid milk & yoghurt. - Used as Anti-infective agent.	- Limit Dairy products - Used as Antibiotic.	- May cause diarrhea (increase fluid intake) - Used as Antibiotic.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Used as Antimycobacterial.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Used as Antitubercular.
	Linezolid	Pyrimethamine	Doxycycline	Pentamidine	Ampicillin
	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Used as Antibacterial.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Used as Antibacterial.	- Avoid with folic acid containing Folate - Avoid dairy products. - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr. before or after - Used as Antibiotic.	- Avoid dairy products. - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr. before or after - Used as Antibiotic.	- Absorption is decreased with tube feeding - Used as Antibiotic.
	Ciprofloxacin	Minocycline	Metronidazole	Erythromycin	Loratadine
	- Avoid with caffeine - Absorption is decreased with tube feeding. - Used as Anti-infective agent.	- Avoid dairy products. - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr. before or after - tetracycline Antibacterial.	- Avoid with Fatty meal or tube feeding - Anti-infective, Antiprotozoal.	- Avoid grapefruits - Used as Antibiotic.	- Absorption is decreased with tube feeding. - Used as Antibiotic.

Nutrients – Drug Interaction Table (Antibiotic, Anti-infectives & Antibacterials)					
Uses and Interactions	Rifampin	Pyrazinamide	Cycloserine	Sulfasalazine	Azufidine
	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr before or after. - Used as Antimycobacterial.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Absorption is decreased with meals or tube feeding: Should be taken 1-2 hr before or after. - Used as Antimycobacterial.	- Avoid of high-fat meals - Monitor Diarrhea - Used as Antibiotic.	- Avoid with folic acid or with meals containing Folate - Used as Anti-infective agent, Anti GI tract inflammatory, Anti-rheumatic.	- Avoid with folic acid - Used as Anti-infective agent - Anti GI tract inflammatory, Anti-rheumatic.
	Ofloxacin	Itraconazole			
	- Avoid dairy products - Absorption is decreased with tube feeding - Used as Anti-infective.	- Avoid grapefruits - Used an Antifungal.			

Nutrients – Drug Interaction Table (Anit-hyperlipidemic drugs)					
Uses and Interactions	Atorvastatin	Fluvastatin	Lovastatin	Pravastatin	Rosuvastatin
	-Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	-Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	-Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	-Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).
	Simvastatin	Cerivastatin	Alendronate sodium	Propranolol	
	-Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	- Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	- Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	- Avoid grapefruits . -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	

Nutrients – Drug Interaction Table (Grapefruits)

	Nutrients – Drug Interaction Table (Grapefruits)				
	Fentanyl	Atorvastatin	Midazolam	Budesonide	Methylprednisolone
Uses	- Avoid grapefruits. - Used as sedative agents.	- Avoid grapefruits - Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	- Avoid grapefruits. - Used as sedative agents.	- Avoid grapefruits & - Avoid high fat food. - Used as diuretic.	- Low Na diet is needed - Avoid grapefruits. - Used as Anti-inflammatory, Immunosuppressant
Interactions	Amlodipine	Carbamazepine	Nifedipine	Nicardipine	Diltiazem
	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.	- Avoid grapefruits - Monitor Iron & Sodium intake & Levels - Used as Anticonvulsant.	- Avoid grapefruits - Used as Antihypertensive agent.	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.
	Felodipine	Cilostazol	Enalapril	Diazepam	Cyclosporine
	- Avoid grapefruits juice - Calcium Channel Blockers used as Antihypertensive agent.	- Monitor Vit K, Omega 3 rich food and garlic and ginger. - Avoid grapefruits - Avoid high fat meals - Used as Anticoagulant	- Avoid grapefruits - ACE inhibitors used as Antihypertensive agent.	- Avoid grapefruits - Used as Antianxiety.	- Avoid grapefruits - Used as immunosuppressant
	Amiodarone	Saquinavir	Indinavir	Tacrolimus	Sirolimus
	- Avoid grapefruits. - Used as Antiarrhythmic.	- Avoid grapefruits - Used as protease inhibitor.	- Avoid grapefruits - Used as protease inhibitor.	- Avoid grapefruits - Used as immunosuppressant.	- Avoid grapefruits - Used as immunosuppressant

Diuretic. **Antihypertensive.** **Sedative.** Tube feeding. Antibiotic, Anti-infective & Antibacterial. **Antihyperlipidemic.** **Grapefruits.** **Sodium.** Others

Nutrients – Drug Interaction Table (Grapefruits)					
Uses and Interactions	Sildenafil	Fluvastatin	Lovastatin	Pravastatin	Simvastatin
	- Avoid grapefruits & high fat meal or tube feeding - Used as protease inhibitor.	-Avoid grapefruits. -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	-Avoid grapefruits -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	-Avoid grapefruits -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	- Avoid grapefruits -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).
	Rosuvastatin	Alprazolam	Buspirone	Clomipramine	Erythromycin
	-Avoid grapefruits. -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	-Avoid grapefruits - Used as Antianxiety.	- Avoid grapefruits - Used as Antianxiety.	- Avoid grapefruits - Used as Antianxiety.	-Avoid grapefruits - Used as Antibiotic.
	Triazolam	Itraconazole	Ebastine	Terfenadine	Nimodipine
	- Avoid grapefruits - Used as sedative agent & Antidepressant.	- Avoid grapefruits - Used as Antifungal.	- Avoid grapefruit - Antihistamines.	- Avoid grapefruit - Antihistamines.	- Avoid grapefruit - Calcium Channel Blockers Used as cerebral vasodilator.
	Nisoldipine	Nitrendipine	Cyclosporine	Cerivastatin	Propranolol
	- Avoid grapefruit - Avoid high fat meal or tube feeding - Calcium Channel Blockers Used as Antihypertensive agent.	- Avoid grapefruit - Calcium Channel Blockers Used as Antihypertensive agent.	- Causes hyperglycemia -Avoid grapefruit - Avoid high fat meal or tube feeding - Immunosuppressant.	- Avoid grapefruits. -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).	- Avoid grapefruits. -Used to treat Antihyperlipidemic agent (inhibit cholesterol synthesis).

Nutrients – Drug Interaction Table (Sodium)					
Uses and Interactions	Hydrocortisone	Furosemide	Dexamethasone	Duloxetine	Methylprednisolone
	- Low Na diet is needed. - Anti-inflammatory & Immunosuppressant	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Diuretic and Antihypertensive agent.	- Low Na is recommended. - Anti-inflammatory agent.	- Causes hyponatremia - Monitor Na intake & levels - Used as antidepressant.	- Low Na diet is needed, avoid grapefruits - Used as Anti-inflammatory, Immunosuppressant.
	Carbamazepine	Bumetanide	Escitalopram	Chlorothiazide	Triamterene
	- Avoid grapefruits - Monitor Iron & Sodium intake & Levels - Used as Anticonvulsant.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Monitor Sodium intake & levels - Used as Antidepressant.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic and Antihypertensive agent.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium & the potassium excretion. - Used as diuretic and Antihypertensive agent.
	Metolazone	Lithium	Hydrochlorothiazide		
	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium & the potassium excretion. - Used as diuretic and Antihypertensive agent.	- Monitor Na and fluid intakes & levels - Used to treat Bipolar disorder.	- Monitor Potassium levels and intake - Inhibits reabsorption of sodium. - Used as diuretic Antihypertensive agent.		

Nutrients – Drug Interaction Table (Others)					
Uses and Interactions	Acetylsalicylic Acid	Aspirin	Warfarin	Cephalosporins	Enoxaparin
	- Avoid citrus fruits - Used as anti-inflammatory agent.	- Monitor Vit K, Omega 3 rich food and garlic and ginger. - Used as Anticoagulant.	- Monitor Vit K, Omega 3 rich food and garlic and ginger. - Used as Anticoagulant.	- Monitor Vitamin K levels and Intake - Uses as Blood Coagulant.	- Monitor Vit K, Omega 3 rich food and garlic and ginger. - Used as Anticoagulant.
	Cimetidine	Methotrexate	Digoxin	Quinidine	Isotretinoin
	- Avoid with Caffeine containing food & beverages - Antiulcer drug.	- Avoid with dairy Products - Monitor B12, Folic acid, Fat & Calcium intake & levels - Used as Antineoplastic.	- Consumption of high fiber foods or fruit juices after 2 hours from taking the drug is recommended - Monitor Potassium levels and intake - Used as Antiarrhythmic.	- Avoid with Citrus juice. - Avoid with Vitamin C supplements. - vitamin K levels and intake - Used as Antiarrhythmic.	- Increase fluid intake to avoid dehydration. - Used to treat cystic acne.
	Theophylline	Albuterol	Tranlycypromine	Phenelzine	Clozapine
	- Avoid caffeine sources. - Avoid high protein low Carb & high Carb & low Protein diets. - Used as bronchodilator.	- Avoid caffeine sources. - Used as bronchodilator.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Used as MAOIs Antidepressant.	- Avoid tyramine containing foods such as (Banana, Salted Fish, Chocolate, caffeine, Avocado, Soy Sauce, Spinach, Tofu, Tomatoes & Yeast). - Used as MAOIs Antidepressant.	- Avoid caffeine - Used as Antipsychotic.

References:

- <https://www.ncbi.nlm.nih.gov> (National Center for Biotechnology Information)
- <https://www.drugs.com>
- Avoid Food- Drug Interactions: (A Guide from the National Consumers League and U.S. Food and Drug Administration)
- Handbook of Food- Drug Interactions
edited by Beverly J. McCabe Eric H. Frankel Jonathan J. Wolfe

Diuretic. **Antihypertensive.** **Sedative.** **Tube feeding.** **Antibiotic, Anti-infective & Antibacterial.** **Antihyperlipidemic.** **Grapefruits.** **Sodium.** **Others**