

Training course on the inpatient management of severe acute malnutrition: Module 4. Feeding

Web Annex A.

F-75 reference card

Volume of F-75 to give for children without oedema or with mild or moderate oedema (O + ++)

See reverse for adjusted amounts for children with severe oedema (+++)

Child weight (kg)	Volume of F-75 per feed (ml) ^a			Daily total (130 ml/kg)	80% of daily total ^a minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
2.0	20	30	45	260	210
2.2	25	35	50	286	230
2.4	25	40	55	312	250
2.6	30	45	55	338	265
2.8	30	45	60	364	290
3.0	35	50	65	390	310
3.2	35	55	70	416	335
3.4	35	55	75	442	355
3.6	40	60	80	468	375
3.8	40	60	85	494	395
4.0	45	65	90	520	415
4.2	45	70	90	546	435
4.4	50	70	95	572	460
4.6	50	75	100	598	480
4.8	55	80	105	624	500
5.0	55	80	110	650	520
5.2	55	85	115	676	540
5.4	60	90	120	702	560
5.6	60	90	125	728	580
5.8	65	95	130	754	605
6.0	65	100	130	780	625
6.2	70	100	135	806	645
6.4	70	105	140	832	665
6.6	75	110	145	858	685
6.8	75	110	150	884	705
7.0	75	115	155	910	730
7.2	80	120	160	936	750
7.4	80	120	160	962	770

Child weight (kg)	Volume of F-75 per feed (ml) ^a			Daily total (130 ml/kg)	80% of daily total ^a minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
7.6	85	125	165	988	790
7.8	85	130	170	1014	810
8.0	90	130	175	1040	830
8.2	90	135	180	1066	855
8.4	90	140	185	1092	875
8.6	95	140	190	1118	895
8.8	95	145	195	1144	915
9.0	100	145	200	1170	935
9.2	100	150	200	1196	960
9.4	105	155	205	1222	980
9.6	105	155	210	1248	1000
9.8	110	160	215	1274	1020
10.0	110	160	220	1300	1040

^a Volumes in these columns are rounded to the nearest 5 ml.

^b Feed 2-hourly for at least the first day. Then, when little or no vomiting, modest diarrhoea (< 5 watery stools per day), and finishing most feeds, change to 3-hourly feeds.

^c After a day on 3-hourly feeds: if no vomiting, less diarrhoea, and finishing most feeds, change to 4-hourly feeds.

Volume of F-75 for children with severe (+++) oedema

Weight with +++ oedema (kg)	Volume of F-75 per feed (ml) ^a			Daily total (100 ml/kg)	80% of daily total ^a (minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
3.0	25	40	50	300	240
3.2	25	40	55	320	255
3.4	30	45	60	340	270
3.6	30	45	60	360	290
3.8	30	50	65	380	305
4.0	35	50	65	400	320
4.2	35	55	70	420	335
4.4	35	55	75	440	350
4.6	40	60	75	460	370
4.8	40	60	80	480	385
5.0	40	65	85	500	400
5.2	45	65	85	520	415
5.4	45	70	90	540	430
5.6	45	70	95	560	450
5.8	50	75	95	580	465
6.0	50	75	100	600	480
6.2	50	80	105	620	495
6.4	55	80	105	640	510

Weight with +++ oedema (kg)	Volume of F-75 per feed (ml) ^a			Daily total (100 ml/kg)	80% of daily total ^a (minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
6.6	55	85	110	660	530
6.8	55	85	115	680	545
7.0	60	90	115	700	560
7.2	60	90	120	720	575
7.4	60	95	125	740	590
7.6	65	95	125	760	610
7.8	65	100	130	780	625
8.0	65	100	135	800	640
8.2	70	105	135	820	655
8.4	70	105	140	840	670
8.6	70	110	145	860	690
8.8	75	110	145	880	705
9.0	75	115	150	900	720
9.2	75	115	155	920	735
9.4	80	120	155	940	750
9.6	80	120	160	960	770
9.8	80	125	165	980	785
10.0	85	125	165	1000	800
10.2	85	130	170	1020	815
10.4	85	130	175	1040	830
10.6	90	135	175	1060	850
10.8	90	135	180	1080	865
11.0	90	140	185	1100	880
11.2	95	140	185	1120	895
11.4	95	145	190	1140	910
11.6	95	145	195	1160	930
11.8	100	150	195	1180	945
12.0	100	150	200	1200	960

^a Volumes in these columns are rounded to the nearest 5 ml.

^b Feed 2-hourly for at least the first day. Then, when little or no vomiting, modest diarrhoea (< 5 watery stools per day), and finishing most feeds, change to 3-hourly feeds.

^c After a day on 3-hourly feeds: if no vomiting, less diarrhoea, and finishing most feeds, change to 4-hourly feeds.