

Drug food interactions



Prepared By
Pharmacist **Zaki Al Nahab**
Drug Information
Pharmaceutical Care Dept
January 2019

Drug Food Interactions

Drug	Food	Interaction
AcetaZOLAMIDE		No interaction
Aciclovir (Acyclovir)		No interaction
Acitretin		No interaction
Adefovir		No interaction
Adenosine	Caffeine	May result in increased risk of seizures; decreased therapeutic effect of adenosine.
Albendazole	Grapefruit juice	May result in increased risk of albendazole adverse effects
Alendronate	Dairy foods	May result in reduced alendronate exposure.
	Orange juice	May result in decreased alendronate bioavailability.
	Food	May result in decreased alendronate bioavailability.
alfuzosin	Grapefruit juice	May result in increased alfuzosin plasma concentrations.
Allopurinol		No interaction
ALPRAZolam	Grapefruit juice	May result in increased alprazolam exposure
	Caffeine	May result in reduced sedative and anxiolytic effects of alprazolam
Aluminium Hydroxide and Magnesium Hydroxide		No interaction
Alteplase	Celery	May result in increased risk of bleeding.
Aminophylline	Caffeine	May result in increased theophylline plasma concentrations.
	Food	May result in altered theophylline concentrations
Amantadine		No interaction
Amitriptyline		No interaction
AmLODIPine		No interaction

Drug	Food	Interaction
Amoxicillin		No interaction
Amoxicillin and Clavulanate		No interaction
Apixaban	Grapefruit juice	May result in increased bleeding risk
ARIPiprazole	Grapefruit juice	May result in increased aripiprazole exposure.
Aspirin	Celery	May result in increased risk of bleeding
Atazanavir	Food	May result in enhanced bioavailability of atazanavir and reduced pharmacokinetic variability
Atenolol		No interaction
AtorvaSTATin	Grapefruit juice	May result in increased bioavailability of atorvastatin resulting in an increased risk of myopathy or rhabdomyolysis.
AzaTHIOprine		No interaction
Azithromycin		No interaction
Carbidopa/Levodopa	High protein food	May result in decreased levodopa concentrations.
Benztropine		No interaction
Benzylpenicillin [Penicillin G]		No interaction
Benzathine		No interaction
Betahistine		No interaction
Bisacodyl	Milk	May result in gastric or duodenal irritation.
Bisoprolol		No interaction
Bosentan		No interaction
Bromocriptine	Grapefruit juice	May result in increased bromocriptine concentrations.
Cabergoline		No interaction
Calcium Carbonate	Oxalic acid foods	May result in decreased calcium exposure.
	Phytic acid foods	May result in decreased calcium effectiveness.
Calcium Folate		No interaction

Drug	Food	Interaction
Calcium Gluconate	Oxalic acid foods	May result in decreased calcium exposure.
	Phytic acid foods	May result in decreased calcium effectiveness.
Candesartan		No interaction
Captopril	Food	May result in decreased captopril concentration
Carbachol		No interaction
CarBAMazepine	Grapefruit juice	May result in increased carbamazepine bioavailability.
Carbidopa/LEVODOPA	High protein food	May result in decreased levodopa concentrations.
Carvedilol		No interaction
Castor Oil		No interaction
Cefixime		No interaction
Cefuroxime		No interaction
Celecoxib		No interaction
Cetirizine		No interaction
Charcoal, Activated		No interaction
Chloral Hydrate		No interaction
Chloroquine		No interaction
Chlorpheniramine		No interaction
ChlorproMAZINE		No interaction
CycloSPORINE	Pomelo juice	May result in an increased risk of cycloSPORINE toxicity.
	Grapefruit juice	May result in an increased risk of cycloSPORINE toxicity (renal dysfunction, cholestasis, paresthesias).
Cinacalcet	Food	May result in increased bioavailability of cinacalcet.
Cinnarizine		No interaction
Ciprofloxacin	Caffeine	May result in increased caffeine concentrations and enhanced CNS stimulation.
	Dairy foods	May result in decreased ciprofloxacin concentrations.

Drug	Food	Interaction
Citalopram		No interaction
Clarithromycin		No interaction
Clindamycin		No interaction
CloBAZam	Caffeine	May result in reduced sedative and anxiolytic effects of clobazam.
Clofazimine	Orange juice	May result in reduced clofazimine plasma concentrations.
Clomifene		No interaction
ClomiPRAMINE	Grapefruit juice	May result in an increased risk of clomiPRAMINE toxicity.
ClonazePAM	Caffeine	may result in reduced sedative and anxiolytic effects of clonazepam.
CloNIDine		No interaction
Clopidogrel	Grapefruit juice	May result in reduced exposure of the active clopidogrel metabolite.
	Celery	May result in increased risk of bleeding.
Cloxacillin		No interaction
CloZAPine	Grapefruit juice	May result in increased clozapine plasma levels.
	Caffeine	may result in an increased risk of clozapine toxicity (sedation, seizures, hypotension).
Colchicine	Grapefruit juice	May result in increased colchicine concentrations and increased risk for colchicine toxicity.
Cholecalciferol	Olestra	May result in decreased systemic cholecalciferol (vitamin D) concentrations.
Cholestyramine		No interaction
Colistimethate		No interaction
Cotrimoxazole		No interaction
Cyanocobalamin		No interaction

Drug	Food	Interaction
Cyclopentolate		No interaction
CycloSERINE		No interaction
Cyproterone		No interaction
Dabigatran		No interaction
Daclatasvir	Grapefruit juice	May result in increased daclatasvir exposure.
Dalteparin	Celery	May result in increased risk of bleeding.
Danazol		No interaction
Dantrolene		No interaction
Dapsone		No interaction
Darunavir and Cobicistat		No interaction
Dasabuvir		No interaction
Dasatinib	Grapefruit juice	May result in increased dasatinib plasma concentrations.
Deferasirox	Food	May result in variably increased deferasirox bioavailability.
Deferiprone		No interaction
Denosumab		No interaction
Desmopressin		No interaction
Dexamethasone		No interaction
Dextromethorphan	Grapefruit juice	May result in increased absorption, bioavailability, and excretion of dextromethorphan by grapefruit juice.
Diazepam	Grapefruit juice	May result in increased plasma concentrations of diazepam.
	Caffeine	May result in reduced sedative and anxiolytic effects of diazepam.
	High fat food	May result in increased diazepam concentrations.
Diazoxide		No interaction
Diclofenac		No interaction
Digoxin	Food high in bran	May result in decreased digoxin concentrations.

Drug	Food	Interaction
Diltiazem	Food	May result in increased plasma concentrations.
	Grapefruit juice	May result in increased serum diltiazem concentrations.
Dimethyl Fumarate		No interaction
Diphenhydramine		No interaction
Dolutegravir		No interaction
Domperidone	Grapefruit juice	May result in increased domperidone plasma concentrations.
Doxazosin		No interaction
Doxycycline		No interaction
Duloxetine		No interaction
Edetate CALCIUM Disodium		No interaction
Efavirenz	Food	May result in increased efavirenz absorption.
Elbasvir and Grazoprevir		No interaction
Elvitegravir, Cobicistat, Emtricitabine, and Tenofovir Alafenamide		No interaction
Emtricitabine and Tenofovir Alafenamide		No interaction
Enalapril		No interaction
Enoxaparin	Celery	May result in increased risk of bleeding.
Entecavir	Food	May result in decreased entecavir exposure.
EPHEDrine	Bitter orange	May result in hypertensive crisis (headache, hyperpyrexia, hypertension).
Erythromycin	Food	May result in altered erythromycin concentrations.
	Grapefruit juice	May result in increased erythromycin bioavailability.
Escitalopram		No interaction
Esomeprazole	Cranberry	May result in reduced effectiveness of proton pump inhibitors.

Drug	Food	Interaction
Ethambutol		No interaction
Ethionamide		No interaction
Ethosuximide		No interaction
Etravirine		No interaction
Ezetimibe		No interaction
FentaNYL	Grapefruit juice	May result in an increased risk of fentanyl toxicity (CNS depression, respiratory depression).
Ferrous Fumarate Ferrous Sulfate	Phytic acid foods	May result in reduced iron absorption.
	Dairy foods	May result in decreased iron bioavailability.
	Soy	May result in reduced iron absorption.
Finasteride		No interaction
Fingolimod		No interaction
FlavoxATE		No interaction
Fluconazole		No interaction
Fluconazole		No interaction
Fludrocortisone		No interaction
Flumazenil		No interaction
FLUoxetine		No interaction
FluPHENAZine		No interaction
Fluticasone	Grapefruit juice	May result in increased fluticasone exposure.
Fluvastatin		No interaction
Fluvoxamine	Grapefruit juice	May result in increased fluvoxamine exposure.
Folic Acid		No interaction
Follitropin Alfa		No interaction
Follitropin Beta		No interaction
Fondaparinux	Avocado	May result in reduced anticoagulant effectiveness.
	Celery	May result in increased risk of bleeding.

Drug	Food	Interaction
Furosemide	Food	May result in decreased furosemide exposure and efficacy.
Gabapentin		No interaction
Gemfibrozil		No interaction
Gentamicin		No interaction
Glibenclamide		No interaction
Gliclazide		No interaction
Glimepiride		No interaction
Griseofulvin	Food	May result in increased griseofulvin exposure.
GuaiFENesin		No interaction
Haloperidol		No interaction
Heparin	Avocado	May result in reduced anticoagulant effectiveness.
	Celery	May result in increased risk of bleeding.
HydrALAZINE	Enteral nutrition	May result in decreased hydrALAZINE concentrations.
	Food	May result in decreased hydrALAZINE exposure and efficacy.
HydroCHLOROthiazide		No interaction
Hydrocortisone		No interaction
HYDROmorphone		No interaction
Hydroquinone		No interaction
Hydroxyurea		No interaction
Hydroxychloroquine		No interaction
HydrOXYzine		No interaction
Hyoscine Butylbromide [Scopolamine]		No interaction
Ibuprofen		No interaction
Iloprost	Celery	May result in increased risk of bleeding.
Imatinib	Grapefruit juice	May result in increased imatinib plasma concentrations.
Imipramine		No interaction

Drug	Food	Interaction
Indapamide		No interaction
Indomethacin		No interaction
Irbesartan		No interaction
Iron Sucrose	Phytic acid foods	May result in reduced iron absorption.
Isoniazid	Food	May result in decreased isoniazid exposure.
	Tyramine foods	May result in increased blood pressure.
	Histamine containing food	May result in exaggerated response to foods containing histamines (eg, skipjack, tuna, other tropical fish).
Isosorbide Dinitrate		No interaction
ISOTretinoin		No interaction
Ispaghula [Psyllium]		No interaction
Itraconazole	Grapefruit juice	May result in decreased oral bioavailability of itraconazole resulting in an increased risk of clinical antifungal failure.
	Food	May result in increased or decreased bioavailability of itraconazole, depending on the dosage form of itraconazole.
Ivabradine	Grapefruit juice	May result in increased ivabradine exposure.
Labetalol	Food	May result in increased labetalol exposure.
Lactulose		No interaction
Lamivudine\Zidovudine		No interaction
Lamotrigine		No interaction
Ledipasvir\Sofosbuvir		No interaction
Lercanidipine	Grapefruit juice	May result in increased lercanidipine exposure and increased hypotension.
Letrozole		No interaction
Levetiracetam		No interaction
Levofloxacin		No interaction

Drug	Food	Interaction
Levothyroxine	Enteral nutrition	May result in hypothyroidism.
	Soy	May result in decreased effectiveness of levothyroxine
Lidocaine		No interaction
Linagliptin		No interaction
Linezolid	Avocado	May result in hypertensive crisis (headache, palpitation, neck stiffness).
	Bitter orange	May result in hypertensive crisis (headache, hyperpyrexia, hypertension).
	Tyramine foods	May result in a significant pressor response.
Liraglutide		No interaction
Lisinopril		No interaction
Loperamide		No interaction
Lopinavir\Ritonavir		No interaction
Loratadine		No interaction
LORazepam	Caffeine	May result in reduced sedative and anxiolytic effects of lorazepam.
Losartan	Grapefruit juice	May result in increased half-life ($t_{1/2}$) and decreased area under the concentration time curve (AUC) of losartan's active metabolite (E3174).
Magnesium Sulfate		No interaction
Mebendazole		No interaction
Mebeverine		No interaction
MedroxyPROGESTERone	Caffeine	May result in enhanced CNS stimulation.
Mefloquine		No interaction
Meloxicam		No interaction
Memantine		No interaction
Meropenem		No interaction
Mesalamine		No interaction
Mesna		No interaction

Drug	Food	Interaction
MetFORMIN		No interaction
Methadone	Grapefruit juice	May result in increased methadone exposure, decreased clearance of all enantiomers (R-, S-, R,S-methadone).
Methotrexate	Cola	May result in increased methotrexate serum levels and increased risk of toxicity.
Methoxy Polyethylene Glycol-Epoetin Beta		No interaction
Methyldopa		No interaction
Methylphenidate		No interaction
MethylPREDNISolone	Grapefruit juice	May result in increased methylPREDNISolone exposure.
Metoclopramide		No interaction
MetOLazone		No interaction
Metoprolol	Food	May result in increased metoprolol concentrations.
MetroNIDAZOLE		No interaction
Mexiletine	Caffeine	May result in an increased risk of caffeine adverse effects
Miconazole		No interaction
Midazolam	Grapefruit juice	May result in increased bioavailability and pharmacodynamic effects of midazolam.
	Bitter orange	May result in increased CNS depression.
	Caffeine	May result in reduced sedative and anxiolytic effects of midazolam.
Mirtazapine	Grapefruit juice	May result in increased mirtazapine exposure.
Modafinil		No interaction
Mometasone		No interaction
Montelukast	Grapefruit juice	May result in increased montelukast exposure.
Morphine		No interaction

Drug	Food	Interaction
Moxifloxacin		No interaction
Mycophenolate		No interaction
Naproxen		No interaction
Neostigmine		No interaction
NIFEdipine	Grapefruit juice	May result in severe hypotension, myocardial ischemia, increased vasodilator side effects.
Nilotinib	Grapefruit juice	May result in increased nilotinib plasma concentrations.
	Food	May result in increased nilotinib exposure.
NiMODipine	Grapefruit juice	May result in increased nimodipine bioavailability.
	Food	May result in reduced nimodipine efficacy.
Nitrofurantoin		No interaction
Norethindrone	Caffeine	May result in enhanced CNS stimulation.
Nystatin		No interaction
Ofloxacin		No interaction
OLANZapine		No interaction
Olmesartan		No interaction
Ombitasvir, Paritaprevir, and Ritonavir (Viekirax®)	Grapefruit juice	May result in increased paritaprevir exposure.
Omeprazole	Cranberry	May result in reduced effectiveness of proton pump inhibitors.
Ondansetron		No interaction
Oral Rehydration Therapy		No interaction
Oseltamivir		No interaction
Paliperidone		No interaction
Pantoprazole	Cranberry	May result in reduced effectiveness of proton pump inhibitors.

Drug	Food	Interaction
Paracetamol	Cabbage	May result in decreased acetaminophen effectiveness.
	Food	May result in decreased peak acetaminophen concentrations.
Paricalcitol		No interaction
PARoxetine		No interaction
PenicillAMINE	Food	May result in decreased penicillamine concentrations.
Pentamidine		No interaction
Pentoxifylline		No interaction
Perindopril		No interaction
Pethidine	Grapefruit juice	May result in increased Pethidine plasma concentrations and increased risk of respiratory depression.
PHENobarbital		No interaction
Phenoxymethylpenicillin		No interaction
Phentolamine		No interaction
Phenylephrine	Bitter orange	May result in hypertensive crisis (headache, hyperpyrexia, hypertension).
Phenytoin	Enteral nutrition	May result in decreased serum phenytoin levels and subsequently reduced therapeutic response to phenytoin.
	Food	May result in altered phenytoin concentrations.
Physostigmine		No interaction
Phytonadione	Olestra	May result in decreased vitamin K levels.
Pioglitazone		No interaction
Polyethylene Glycol-Electrolyte Solution		No interaction
Pramipexole		No interaction
Praziquantel		No interaction
Prazosin		No interaction
PrednisolONE		No interaction

Drug	Food	Interaction
Pregabalin		No interaction
Primaquine	Grapefruit juice	May result in increased primaquine plasma concentration.
Procyclidine		No interaction
Promethazine		No interaction
Propafenone	Grapefruit juice	May result in increased propafenone exposure.
Propranolol	Food	May result in increased propranolol concentrations.
Propylthiouracil	Food	May result in altered drug concentrations.
Pyrazinamide		No interaction
Pyridoxine	Food	May result in decreased pyridoxine exposure
Pyrimethamine		No interaction
QUetiapine	Grapefruit juice	May result in increased quetiapine exposure
QuiNIDine	Grapefruit juice	May result in decreased metabolic conversion of quinidine to 3-hydroxyquinidine.
	Food	May result in increased quinidine concentrations.
	Alkaline foods	May result in increased quinidine concentrations.
QuiNINE		No interaction
Raltegravir		No interaction
RaNITidine	Cranberry	May result in reduced effectiveness of histamine-2 blockers.
Retepase	Celery	May result in increased risk of bleeding.
Ribavirin		No interaction

Drug	Food	Interaction
Rifampicin	Food	May result in decreased rifampin concentrations.
Rifampicin/Isoniazide	Food	May result in decreased isoniazid exposure.
		May result in decreased rifampin concentrations.
	Tyramine foods	Concurrent use of isoniazid and tyramine foods may result in increased blood pressure.
	Histamine containing food	Concurrent use of isoniazid and histamine containing food may result in exaggerated response to foods containing histamines (eg, skipjack, tuna, other tropical fish).
Riluzole		No interaction
Risperidone		No interaction
Ritonavir		No interaction
Rivaroxaban		No interaction
Rosuvastatin		No interaction
Salbutamol		No interaction
Saxagliptin		No interaction
Senna		No interaction
Sertraline	Grapefruit juice	May result in elevated sertraline serum concentrations and an increased risk of adverse side effects.
Sevelamer		No interaction
Sildenafil	Grapefruit juice	May result in increased sildenafil bioavailability and delayed sildenafil absorption.
	Pomelo juice	May result in reduced plasma concentrations and efficacy of sildenafil.

Drug	Food	Interaction
Simeprevir	Grapefruit juice	May result in increased simeprevir exposure.
Simeticone		No interaction
Simvastatin	Grapefruit juice	May result in increased bioavailability of simvastatin resulting in an increased risk of myopathy or rhabdomyolysis.
	Cranberry juice	May result in increased risk of hepatitis and myopathy/rhabdomyolysis.
Sirolimus	Grapefruit juice	May result in an increased risk of sirolimus toxicity (anemia, leukopenia, thrombocytopenia, hypokalemia, diarrhea).
SITagliptin		No interaction
Sofosbuvir		No interaction
Sofosbuvir and Velpatasvir		No interaction
Somatropin		No interaction
SORafenib		No interaction
Sotalol		No interaction
Spirolactone		No interaction
Streptokinase	Celery	May result in increased risk of bleeding.
Streptomycin		No interaction
Sucralfate		No interaction
SulfaSALazine		No interaction
Sulpiride		No interaction
SUMatriptan		No interaction
Tacrolimus	Grapefruit	May result in increased tacrolimus concentration.

Drug	Food	Interaction
	Pomegranate	May result in increased tacrolimus exposure.
Tamsulosin	Grapefruit juice	May result in an increase in tamsulosin plasma exposure.
	Food	May result in decreased tamsulosin bioavailability.
Telaprevir		No interaction
Telmisartan		No interaction
Telmisartan and Hydrochlorothiazide		No interaction
Tenofovir Alafenamide		No interaction
Terbinafine		No interaction
Teriflunomide		No interaction
Tetracycline	Dairy foods	May result in decreased tetracycline concentrations.
Theophylline	Caffeine	May result in increased theophylline plasma concentrations.
	Food	May result in altered theophylline concentrations.
Thiamine		No interaction
Ticagrelor	Grapefruit juice	May result in increased ticagrelor exposure and antiplatelet activity.
Tinzaparin	Celery	May result in increased risk of bleeding.
Tirofiban	Celery	May result in increased risk of bleeding.
Tolterodine		No interaction
Topiramate		No interaction
TraMADol	Grapefruit juice	May result in increased tramadol plasma concentrations and increased risk of respiratory depression.

Drug	Food	Interaction
Tranexamic Acid		No interaction
Trihexyphenidyl		No interaction
Trimetazidine		No interaction
Tripolidine and Pseudoephedrine	Bitter orange	May result in hypertensive crisis (headache, hyperpyrexia, hypertension).
Ursodeoxycholic Acid		No interaction
ValACIclovir		No interaction
ValGANciclovir	High fat food	May result in increased ganciclovir exposure.
Valproic Acid		No interaction
Valsartan		No interaction
Valsartan and Hydrochlorothiazide		No interaction
Venlafaxine		No interaction
Verapamil	Grapefruit juice	May result in an increased risk of verapamil adverse effects (flushing, edema, hypotension, myocardial ischemia).
	Caffeine	May result in increased caffeine serum concentrations and enhanced CNS stimulation.
Vildagliptin		No interaction
Vitamin A		No interaction
Vitamin B 1 (Thiamin)		No interaction
Vitamin B 12 (Cyanocobalamin)		No interaction
Vitamin B2		No interaction
Vitamin B 6	Food	may result in decreased pyridoxine exposure.
Vitamin B Complex		No interaction
Vitamin D3	Olestra	May result in decreased systemic cholecalciferol (vitamin D) concentrations.
Vitamin E		No interaction

Drug	Food	Interaction
Vitamin K	Olestra	May result in decreased vitamin K levels.
Voriconazole	Food	May result in decreased voriconazole exposure.
Warfarin	Cranberry juice	May result in an increased risk of bleeding
	Pomegranate	May result in increased warfarin plasma concentrations and increased risk of bleeding.
	Vitamin k foods	May result in altered anticoagulant effectiveness.
	Green tea	May result in reduced anticoagulant effectiveness.
	Black tea	May result in decreased warfarin effectiveness.
	High-protein diet	May result in reduced warfarin anticoagulant effectiveness.
	Enteral nutrition	May result in decreased PT/INR response to warfarin and development of warfarin resistance.
	Avocado	May result in reduced anticoagulant effectiveness.
	Noni juice	May result in risk of acquiring warfarin resistance.
	Soy	May result in reduced warfarin effectiveness.
	Celery	May result in increased risk of bleeding.
	Parsley	May result in increased anticoagulant effectiveness.

Pharmaceutical Care Dept
Drug Information
Pharmacist Zaki Ali Al Nahab
Prepared on January 2019

Reference

IBM Micromedex Solution Online
Lexicomp Online