

Homotopy Theory and Characteristic Classes

CUI Jiaqi
East China Normal University

February 19, 2025

Abstract

This is the notes of a course given by Prof. Ma Langte in 25spring at Shanghai Jiaotong University.

Contents

I	Homotopy Theory	1
1	Cofibrations and Fibrations	2
1.1	Cofibrations	2
II	Generalized Homology	7
III	Characteristic Classes	7

Part I

Homotopy Theory

Let \mathbf{TOP} be the category of topological spaces. Then we can take a quotient of \mathbf{TOP} and get the homotopy category $h\text{-}\mathbf{TOP}$. The quotient may bring more algebraic structures. For example, $\text{Mor}(S^1, X)$, the homotopy classes of maps from S^1 to X , is the fundamental group of X . Our goal is to study functors from homotopy category to some algebraic categories.

Let \mathbf{TOP}^o be the pointed topological category, where the sum is wedge sum $(X, x_0) \wedge (Y, y_0) = X \sqcup Y / x_0 \sim y_0$ and the product is the smash product $(X, x_0) \vee (Y, y_0) = X \times Y / \{x_0\} \times Y \cup X \times \{y_0\}$. Similarly, we can take a quotient to get $h\text{-}\mathbf{TOP}^o$.

Let $\mathbf{TOP}(2)$ be the category of pairs and $h\text{-}\mathbf{TOP}(2)$ be its quotient.

Fix $K \in \text{Ob}(\mathbf{TOP})$. Let's consider \mathbf{TOP}^K , the category of spaces under K . Its objects are maps $f: K \rightarrow X$ and morphisms are maps $\alpha: X \rightarrow Y$ such that $\alpha \circ f = g$.

$$\begin{array}{ccc} & K & \\ f \swarrow & & \searrow g \\ X & \xrightarrow{\alpha} & Y \end{array}$$

If $K = \{*\}$ is a single point set, then $\mathbf{TOP}^{\{*\}} = \mathbf{TOP}^o$ is the pointed topological category. Take $X = K$. A morphism from $f: K \rightarrow X$ to $\text{id}: K \rightarrow K$ is $r: X \rightarrow K$ such that $r \circ f = \text{id}$.

$$\begin{array}{ccc} & K & \\ f \swarrow & & \searrow \text{id} \\ X & \xrightarrow{r} & K \end{array}$$

When $K \subset X$, $f = i: K \hookrightarrow X$, we say that r is a retraction.

We have $r: X \rightarrow K$ is a deformation retraction, if and only if $i \circ r \simeq \text{id}_X \text{ rel } K$, if and only if $r: X \rightarrow K$ is a homotopy equivalence in \mathbf{TOP}^K .

Fix $B \in \text{Ob}(\mathbf{TOP})$. Let's consider \mathbf{TOP}_B , the category of spaces over B , where the objects are $p: X \rightarrow B$ and morphisms are $f: X \rightarrow Y$ such that $p = q \circ f$.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ p \searrow & & \swarrow q \\ & B & \end{array}$$

Take $X = B$. A morphism from $\text{id}: B \rightarrow B$ to $q: Y \rightarrow B$ is $s: B \rightarrow Y$ such that $q \circ s = \text{id}_B$.

$$\begin{array}{ccc} B & \xrightarrow{s} & Y \\ \text{id} \searrow & & \swarrow q \\ & B & \end{array}$$

Then s is called a section of q .

Similarly, we can define $h - \mathbf{TOP}^K$ and $h - \mathbf{TOP}_B$.

1 Cofibrations and Fibrations

1.1 Cofibrations

Definition 1.1. A map $i: A \rightarrow X$ has the homotopy extension property (HEP) for a space Y if for all homotopy $h: A \times I \rightarrow Y$ and $f: X \rightarrow Y$ with $f \circ i(a) = h(a, 1)$, there exists $H: X \times I \rightarrow Y$ satisfies

$$\begin{array}{ccccc} & & X & & \\ & i \nearrow & & \searrow i_1 & \\ A & & & & \\ & i_1 \searrow & & \nearrow i \times \text{id} & \\ & & A \times I & & \\ & & \nearrow h & & \\ & & X \times I & \xrightarrow{H} & Y \end{array}$$

We say $i: A \rightarrow X$ is a cofibration if it has HEP for each $Y \in \text{Ob}(\mathbf{TOP})$.

Recall the mapping cylinder: if $i: A \rightarrow X$ is a map, then $Z(i) := (A \times I) \sqcup X / (a, 1) \sim i(a)$.

Proposition 1.2. Given a map $i: A \rightarrow X$. The followings are equivalent:

1. $i: A \rightarrow X$ is a cofibration.
2. i has HEP for $Z(i)$.

3. The map

$$\begin{aligned} s: Z(i) &\rightarrow X \times I \\ (a, t) &\mapsto (f(a), t), \\ x &\mapsto (x, 1) \end{aligned}$$

has a retraction.

Proof. (1) \implies (2) is only by definition.

(2) \implies (1): By definition, there exists $K: X \times I \rightarrow Z(i)$ such that the following diagram is commutative.

$$\begin{array}{ccccc} & & X & & \\ & \nearrow i & & \searrow i_1 & \\ A & & & & X \times I \xrightarrow{K} Z(i) \\ & \searrow i_1 & & \nearrow i \times \text{id} & \\ & & A \times I & & \end{array}$$

id

For any Y and homotopy $h: A \times I \rightarrow Y$ and $f: X \rightarrow Y$ with $f \circ i(a) = h(a, 1)$, we define

$$\begin{aligned} F: Z(i) &\rightarrow Y \\ (a, t) &\mapsto h(a, t) \\ x &\mapsto f(x). \end{aligned}$$

Then $F \circ K$ is as desired.

$$\begin{array}{ccccc} & & X & & \\ & \nearrow i & & \searrow i_1 & \\ A & & & & X \times I \xrightarrow{K} Z(i) \xrightarrow{F} Y \\ & \searrow i_1 & & \nearrow i \times \text{id} & \\ & & A \times I & & \end{array}$$

$\text{id} \quad h$

(2) \implies (3): We can easily check that the extension $K: X \times I \rightarrow Z(i)$ in the proof of (2) \implies (1) is a retraction of s .

(3) \implies (2): Let r be a retraction of s . For any homotopy $h: A \times I \rightarrow Z(i)$ and $f: X \rightarrow Z(i)$ with $f \circ i(a) = h(a, 1)$, we define

$$\begin{aligned} \sigma: Z(i) &\rightarrow Z(i) \\ (a, t) &\mapsto h(a, t) \\ x &\mapsto f(x). \end{aligned}$$

Then we can verify that $H = \sigma \circ r: X \times I \rightarrow Z(i)$ extends h .

$$\begin{array}{ccccc} & & X & & \\ & \nearrow i & & \searrow i_1 & \\ A & & & & X \times I \xrightarrow{s} Z(i) \\ & \searrow i_1 & & \nearrow i \times \text{id} & \\ & & A \times I & & \end{array}$$

$\text{id} \quad h$

□

Corollary 1.3. When $A \subset X$ is a close subset, $i: A \hookrightarrow X$ is the inclusion map. Then $i: A \rightarrow X$ is a cofibration $\iff Z(i) = A \times I \cup X \times \{1\}$ is a retraction of $X \times I$.

Therefore, we can construct many cofibrations. For example, let (X, A) be a manifold with boundary, then $i: A \hookrightarrow X$ is a cofibration.

Push-Out of Cofibration

Given a commutative diagram,

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ j \downarrow & & \downarrow J \\ X & \xrightarrow{F} & Y \end{array}$$

the push-out of j along f is the initial object of this diagram, i.e. $j: B \rightarrow Y$, $F: X \rightarrow Y$, s.t. $\forall Z$ with $J': B \rightarrow Z$, $F': X \rightarrow Z$ satisfying $J' \circ f = F' \circ j$, $\exists!$ map $p: Y \rightarrow Z$ such that the diagram is commutative.

$$\begin{array}{ccccc} & & B & & \\ & f \nearrow & & \searrow J & \\ A & & & & Y \xrightarrow{\exists! p} Z \\ & j \searrow & & \nearrow F & \\ & & X & & \end{array}$$

In our setting, we can construct $Y = X \sqcup B/f(a) \sim j(a)$ directly.

Proposition 1.4. If $j: A \rightarrow X$ is a cofibration, then the push-out of j along $f: B \rightarrow X$ is also a cofibration.

Proof. For any Z , $g: Y \rightarrow Z$, $h: B \times I \rightarrow Z$ such that $g \circ J = h \circ (i_1 \times \text{id})$, we need to find $H: Y \times I \rightarrow Z$ such that the following diagram is commutative.

$$\begin{array}{ccccc} & & X & & \\ & j \nearrow & & \searrow F & \\ A & & & & Y \xrightarrow{g} Z \\ & f \searrow & & \nearrow J & \\ & & B & & Y \times I \xrightarrow{H} Z \\ & & \searrow & \nearrow & \\ & & B \times I & & \end{array}$$

Because $j: A \rightarrow X$ is a cofibration, we have $G: X \times I \rightarrow Z$ such that the following diagram is commutative.

$$\begin{array}{ccccc} & & X & & \\ & j \nearrow & & \searrow G \circ F & \\ A & & & & X \times I \xrightarrow{G} Z \\ & \searrow & & \nearrow h \circ (f \times \text{id}) & \\ & & A \times I & & \end{array}$$

Using the fact that $J \times \text{id}: B \times I \rightarrow Y \times I$ is also the push-out of $j \times \text{id}: A \times I \rightarrow X \times I$ along $f \times \text{id}: A \times I \rightarrow B \times I$, we have unique $H: Y \times I \rightarrow Z$ such that the following diagram is commutative.

$$\begin{array}{ccccc}
 & & B \times I & & \\
 & \nearrow f \times \text{id} & & \searrow h & \\
 A \times I & & & & Y \times I \xrightarrow{H} Z \\
 & \searrow & & \nearrow G & \\
 & & X \times I & &
 \end{array}$$

The $H: Y \times I \rightarrow Z$ is the extension of $h: B \times I \rightarrow Z$, as desired. \square

In terms of categorical language, let $\Pi(A, B)$ be a category, whose objects are continue maps from A to B and morphisms are homotopy of maps from A to B . Consider $\mathbf{COF}^B \subset \mathbf{TOP}^B$ the subcategory of cofibrations under B (i.e. $J: B \rightarrow Y$). Then we have homotopy category $h - \mathbf{COF}^B$. Given a cofibration $i: A \rightarrow X$, we get a contravariant functor

$$\beta: \Pi(A, B) \rightarrow h - \mathbf{COF}^B.$$

In fact, we only need to check that if $f_0 \simeq f_1: A \rightarrow B$, then we get a morphism from $J_0: B \rightarrow Y_0$ to $J_1: B \rightarrow Y_1$. Firstly, consider the homotopy $J_0 \circ f_t: A \times I \rightarrow Y_0$, we get its extension $\Psi: X \times I \rightarrow Y_0$.

$$\begin{array}{ccccc}
 & & X & & \\
 & \nearrow i & & \searrow F_0 & \\
 A & & & & X \times I \xrightarrow{\Psi} Y_0 \\
 & \searrow & & \nearrow J_0 \circ f_t & \\
 & & A \times I & &
 \end{array}$$

Then by the universal property of the push-out $J_1: B \rightarrow Y_1$ of i along f_1 for $J_0: B \rightarrow Y_0$ and $\Psi_1: X \rightarrow Y_0$, we get a map $K: Y_1 \rightarrow Y_0$, as desired.

$$\begin{array}{ccccc}
 & & B & & \\
 & \nearrow f_1 & & \searrow J_1 & \\
 A & & & & Y_1 \xrightarrow{K} Y_0 \\
 & \searrow i & & \nearrow F_1 & \\
 & & X & &
 \end{array}$$

Replacing a Map by a Cofibration

Given a map $f: X \rightarrow Y$, consider the mapping cylinder $Z(f)$. We can notice that $Z(f)$ is the push-out.

$$\begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 i_1 \downarrow & & \downarrow s \\
 X \times I & \xrightarrow{a} & Z(f)
 \end{array}$$

We also have a map

$$\begin{aligned}
 q: Z(f) &\rightarrow Y \\
 (x, t) &\mapsto f(x).
 \end{aligned}$$

Note that by Proposition 1.2, $i_1: X \hookrightarrow X \times I$ is a cofibration $\iff X \times \{1\} \times I \cup X \times I \times \{1\}$ is a retraction of $X \times I \times I$, we have $s: Y \rightarrow Z(f)$ is a cofibration.

Proposition 1.5. Let

$$\begin{aligned} j: X &\rightarrow Z(f) \\ x &\mapsto (x, 0), \end{aligned}$$

we have

1. $j: X \rightarrow Z(f)$ is a cofibration.
2. $s \circ q \simeq \text{id}_{Z(f)} \text{ rel } Y$.
3. If f is a cofibration, then $q: Z(f) \rightarrow Y$ is a homotopy equivalence in \mathbf{TOP}^X .

Proof. (1). We construct a retraction $R: Z(f) \times I \rightarrow X \times I \cup Z(f) \times \{1\}$ as follow. Let $R': I \times I \rightarrow I \times \{1\} \cup \{0\} \times I$ be a retraction. Then we define

$$\begin{aligned} R: Z(f) \times I &\rightarrow X \times I \cup Z(f) \times \{1\} \\ ((x, s), t) &\mapsto (x, R'(s, t)) \\ (y, t) &\mapsto (y, 1) \end{aligned}$$

is as desired. By Proposition 1.2, $j: X \rightarrow Z(f)$ is a cofibration.

(2). The homotopy

$$\begin{aligned} h_t: Z(f) &\rightarrow Z(f) \\ (x, \sigma) &\mapsto (x, (1-t)\sigma + t) \end{aligned}$$

is as desired.

(3). By Proposition 1.2, there is a retraction $r: Y \times I \rightarrow Z(f)$. Define

$$\begin{aligned} g: Y &\rightarrow Z(f) \\ y &\mapsto r(y, 1). \end{aligned}$$

One can verify that g is the homotopy inverse of q . □

Summery 1. Any map $f: X \rightarrow Y$ factors into

$$X \xrightarrow{j} Z \xrightarrow{q} Y$$

where $j: X \rightarrow Z$ is a cofibration and $q: Z \rightarrow Y$ is a homotopy equivalence. Moreover, such a factorization is unique up to homotopy equivalence. In particular, we can choose $Z = Z(f)$. We define $C_f = Z(f)/\text{im } j$ as the homotopy cofibre of f , i.e. $C_f = X \times I \sqcup Y/(x, 0) \sim *, (x, 1) \sim f(x)$, is called the mapping cone of f .

$$X \xrightarrow{f} Y \xrightarrow{s} C_f$$

The Cofibre Sequence (Puppe's Sequence)

To get finer structure, we work in \mathbf{TOP}^o . Given a map $f: (X, x_0) \rightarrow (Y, y_0)$, we get an induced map

$$\begin{aligned} f^*: [Y, B]^o &\rightarrow [X, B]^o \\ [\alpha] &\mapsto [f \circ \alpha], \end{aligned}$$

where $[X, B]^o$ is the homotopy class of basepoint preserving maps. In particular, we have the constant map

$$\begin{aligned} [*]: X &\rightarrow B \\ x &\mapsto b_0. \end{aligned}$$

Definition 1.6. We say a sequence

$$(X, x_0) \xrightarrow{f} (Y, y_0) \xrightarrow{g} (Z, z_0)$$

in \mathbf{TOP}^o is h-coexact if $\forall (B, b_0) \in \text{Ob}(\mathbf{TOP}^o)$,

$$[Z, B]^o \xrightarrow{g^*} [Y, B]^o \xrightarrow{f^*} [X, B]^o$$

is exact, i.e. $(f^*)^{-1}([*]) = \text{im } g^*$.

In \mathbf{TOP}^o , we consider the reduced mapping cone $CX := X \times I / X \times \{0\} \cup \{x_0\} \times I$. The basepoint of CX is $X \times \{0\} \cup \{x_0\} \times I$. And we consider the reduced mapping cone: For $f: (X, x_0) \rightarrow (Y, y_0)$, $C(f) := CX \vee Y / (x, 1) \sim f(x)$. It is equivalent to the following push-out diagram.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ i_1 \downarrow & & \downarrow f_1 \\ CX & \longrightarrow & C(f) \end{array}$$

We will also use symbol X instead of (X, x_0) in \mathbf{TOP}^o for short.

Proposition 1.7. The sequence

$$X \xrightarrow{f} Y \xrightarrow{f_1} C(f)$$

is h-coexact.

Proof. $f_1 \circ f: X \rightarrow C(f)$ is null-homotopic. □

Part II

Generalized Homology

Part III

Characteristic Classes