

Homotopy Theory and Characteristic Classes

CUI Jiaqi
East China Normal University

April 2, 2025

Abstract

This is the notes of a course given by Prof. Ma Langte in 25spring at Shanghai Jiaotong University. The textbook is *Algebraic Topology* by Tammo tom Dieck.

Contents

I	Homotopy Theory	3
1	Cofibrations and Fibrations	4
1.1	Cofibrations	4
1.1.1	Push-Out of Cofibration	5
1.1.2	Replacing a Map by a Cofibration	7
1.1.3	The Cofibre Sequence (Puppe's Sequence)	8
1.2	Fibrations	11
1.2.1	Pull-back of Fibration	13
1.2.2	Replacing Maps by Fibration	13
1.2.3	Fibre Exact Sequence (Puppe's Sequence)	15
1.3	Duality of Cofibration and Fibration	17
1.3.1	Duality of Reduced Suspension and Loop Space	17
1.3.2	Duality of HLP and HEP	18
1.3.3	Duality of Two Puppe's Sequences	18
2	Homotopy Groups	19
2.1	Definitions and Properties	19
2.2	Change of Basepoint	20
2.3	Serre Fibration	21
2.4	Higher Connectivity	22
2.5	Excision and Suspension	23
2.6	Computation of Homotopy Groups	24
II	Generalized Homology	27
3	Homology Theory and CW-Complexes	27
3.1	Homology Theory	27
3.1.1	Suspension Isomorphism	28
3.2	CW-Complex	29
3.3	CW-Approximation	30
3.4	Eilenberg-MacLane Space	31

3.4.1	Remarks about Compactly Generated Spaces	31
3.5	Spectral Homology	34
4	Cohomology	36
4.1	Axiomatic Cohomology	36
4.1.1	Mayer-Vietoris Sequence	36
III	Characteristic Classes	37

Part I

Homotopy Theory

Let \mathbf{TOP} be the category of topological spaces. Then we can take a quotient of \mathbf{TOP} and get the homotopy category $h\text{-}\mathbf{TOP}$. The quotient may bring more algebraic structures. For example, $\text{Mor}(S^1, X)$, the homotopy classes of maps from S^1 to X , is the fundamental group of X . Our goal is to study functors from homotopy category to some algebraic categories.

Let \mathbf{TOP}^o be the pointed topological category, where the sum is wedge sum $(X, x_0) \wedge (Y, y_0) = X \sqcup Y / x_0 \sim y_0$ and the product is the smash product $(X, x_0) \vee (Y, y_0) = X \times Y / \{x_0\} \times Y \cup X \times \{y_0\}$. Similarly, we can take a quotient to get $h\text{-}\mathbf{TOP}^o$.

Let $\mathbf{TOP}(2)$ be the category of pairs and $h\text{-}\mathbf{TOP}(2)$ be its quotient.

Fix $K \in \text{Ob}(\mathbf{TOP})$. Let's consider \mathbf{TOP}^K , the category of spaces under K . Its objects are maps $f: K \rightarrow X$ and morphisms are maps $\alpha: X \rightarrow Y$ such that $\alpha \circ f = g$.

$$\begin{array}{ccc} & K & \\ f \swarrow & & \searrow g \\ X & \xrightarrow{\alpha} & Y \end{array}$$

If $K = \{*\}$ is a single point set, then $\mathbf{TOP}^{\{*\}} = \mathbf{TOP}^o$ is the pointed topological category. Take $X = K$. A morphism from $f: K \rightarrow X$ to $\text{id}: K \rightarrow K$ is $r: X \rightarrow K$ such that $r \circ f = \text{id}$.

$$\begin{array}{ccc} & K & \\ f \swarrow & & \searrow \text{id} \\ X & \xrightarrow{r} & K \end{array}$$

When $K \subset X$, $f = i: K \hookrightarrow X$, we say that r is a retraction.

We have $r: X \rightarrow K$ is a deformation retraction, if and only if $i \circ r \simeq \text{id}_X \text{ rel } K$, if and only if $r: X \rightarrow K$ is a homotopy equivalence in \mathbf{TOP}^K .

Fix $B \in \text{Ob}(\mathbf{TOP})$. Let's consider \mathbf{TOP}_B , the category of spaces over B , where the objects are $p: X \rightarrow B$ and morphisms are $f: X \rightarrow Y$ such that $p = q \circ f$.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow p & \swarrow q \\ & B & \end{array}$$

Take $X = B$. A morphism from $\text{id}: B \rightarrow B$ to $q: Y \rightarrow B$ is $s: B \rightarrow Y$ such that $q \circ s = \text{id}_B$.

$$\begin{array}{ccc} B & \xrightarrow{s} & Y \\ & \searrow \text{id} & \swarrow q \\ & B & \end{array}$$


Then s is called a section of q .

Similarly, we can define $h\text{-}\mathbf{TOP}^K$ and $h\text{-}\mathbf{TOP}_B$.

1 Cofibrations and Fibrations

1.1 Cofibrations

Definition 1.1. A map $i: A \rightarrow X$ has the homotopy extension property (HEP) for a space Y if for all homotopy $h: A \times I \rightarrow Y$ and $f: X \rightarrow Y$ with $f \circ i(a) = h(a, 1)$, there exists $H: X \times I \rightarrow Y$ satisfies



We say $i: A \rightarrow X$ is a cofibration if it has HEP for each $Y \in \text{Ob}(\mathbf{TOP})$.

Recall the mapping cylinder: if $i: A \rightarrow X$ is a map, then $Z(i) := (A \times I) \sqcup X / (a, 1) \sim i(a)$.

Proposition 1.2. Given a map $i: A \rightarrow X$. The followings are equivalent:


1. $i: A \rightarrow X$ is a cofibration.
2. i has HEP for $Z(i)$.
3. The map

$$\begin{aligned} s: Z(i) &\rightarrow X \times I \\ (a, t) &\mapsto (i(a), t), \\ x &\mapsto (x, 1) \end{aligned}$$

has a retraction.

Proof. (1) \implies (2) is only by definition.


(2) \implies (1): By definition, there exists $K: X \times I \rightarrow Z(i)$ such that the following diagram is commutative.



For any Y and homotopy $h: A \times I \rightarrow Y$ and $f: X \rightarrow Y$ with $f \circ i(a) = h(a, 1)$, we define

$$\begin{aligned} F: Z(i) &\rightarrow Y \\ (a, t) &\mapsto h(a, t) \\ x &\mapsto f(x). \end{aligned}$$

Then $F \circ K$ is as desired.




(2) \implies (3): We can easily check that the extension $K: X \times I \rightarrow Z(i)$ in the proof of (2) \implies (1) is a retraction of s .

(3) \implies (2): Let r be a retraction of s . For any homotopy $h: A \times I \rightarrow Z(i)$ and $f: X \rightarrow Z(i)$ with $f \circ i(a) = h(a, 1)$, we define

$$\begin{aligned}\sigma: Z(i) &\rightarrow Z(i) \\ (a, t) &\mapsto h(a, t) \\ x &\mapsto f(x).\end{aligned}$$

Then we can verify that $H = \sigma \circ r: X \times I \rightarrow Z(i)$ extends h .



□

Corollary 1.3. When $A \subset X$ is a close subset, $i: A \hookrightarrow X$ is the inclusion map. Then $i: A \rightarrow X$ is a cofibration $\iff Z(i) = A \times I \cup X \times \{1\}$ is a retraction of $X \times I$.


Therefore, we can construct many cofibrations. For example, let (X, A) be a manifold with boundary, then $i: A \hookrightarrow X$ is a cofibration.

1.1.1 Push-Out of Cofibration

Given a commutative diagram,

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ j \downarrow & & \downarrow J \\ X & \xrightarrow{F} & Y \end{array}$$


the push-out of j along f is the initial object of this diagram, i.e. $j: B \rightarrow Y$, $F: X \rightarrow Y$, s.t. $\forall Z$ with $J': B \rightarrow Z$, $F': X \rightarrow Z$ satisfying $J' \circ f = F' \circ j$, $\exists!$ map $p: Y \rightarrow Z$ such that the diagram is commutative.




In our setting, we can construct $Y = X \sqcup B/f(a) \sim j(a)$ directly.

Proposition 1.4. If $j: A \rightarrow X$ is a cofibration, then the push-out of j along $f: A \rightarrow B$ $J: B \rightarrow Y$ is also a cofibration.


Proof. For any $Z, g: Y \rightarrow Z, h: B \times I \rightarrow Z$ such that $g \circ J = h \circ (i_1 \times \text{id})$, we need to find $H: Y \times I \rightarrow Z$ such that the following diagram is commutative.



Because $j: A \rightarrow X$ is a cofibration, we have $G: X \times I \rightarrow Z$ such that the following diagram is commutative.



Using the fact that $J \times \text{id}: B \times I \rightarrow Y \times I$ is also the push-out of $j \times \text{id}: A \times I \rightarrow X \times I$ along $f \times \text{id}: A \times I \rightarrow B \times I$, we have unique $H: Y \times I \rightarrow Z$ such that the following diagram is commutative.




The $H: Y \times I \rightarrow Z$ is the extension of $h: B \times I \rightarrow Z$, as desired. \square

In terms of categorical language, let $\Pi(A, B)$ be a category, whose objects are continue maps from A to B and morphisms are homotopy of maps from A to B . Consider $\mathbf{COF}^B \subset \mathbf{TOP}^B$ the subcategory of cofibrations under B (i.e. $J: B \rightarrow Y$). Then we have homotopy category $h - \mathbf{COF}^B$. Given a cofibration $i: A \rightarrow X$, we get a contravariant functor

$$\beta: \Pi(A, B) \rightarrow h - \mathbf{COF}^B.$$

In fact, we only need to check that if $f_0 \simeq f_1: A \rightarrow B$, then we get a morphism from $J_0: B \rightarrow Y_0$ to $J_1: B \rightarrow Y_1$. Firstly, consider the homotopy $J_0 \circ f_t: A \times I \rightarrow Y_0$, we get its extension $\Psi: X \times I \rightarrow Y_0$.



Then by the universal property of the push-out $J_1: B \rightarrow Y_1$ of i along f_1 for $J_0: B \rightarrow Y_0$ and $\Psi_1: X \rightarrow Y_0$, we get a map $K: Y_1 \rightarrow Y_0$, as desired.

$$\begin{array}{ccccc}
 & & B & & \\
 & f_1 \nearrow & & \searrow J_1 & \\
 A & & & & Y_1 \xrightarrow{K} Y_0 \\
 & i \searrow & & \nearrow F_1 & \\
 & & X & &
 \end{array}
 \quad
 \begin{array}{c}
 \text{curved arrow } J_0 \text{ from } B \text{ to } Y_0 \\
 \text{curved arrow } \Psi_1 \text{ from } X \text{ to } Y_0
 \end{array}$$

1.1.2 Replacing a Map by a Cofibration

Given a map $f: X \rightarrow Y$, consider the mapping cylinder $Z(f)$. We can notice that $Z(f)$ is the push-out.

$$\begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 i_1 \downarrow & & \downarrow s \\
 X \times I & \xrightarrow{a} & Z(f)
 \end{array}$$

We also have a map

$$\begin{aligned}
 q: Z(f) &\rightarrow Y \\
 (x, t) &\mapsto f(x).
 \end{aligned}$$

Note that by Proposition 1.2, $i_1: X \hookrightarrow X \times I$ is a cofibration $\iff X \times \{1\} \times I \cup X \times I \times \{1\}$ is a retraction of $X \times I \times I$, we have $s: Y \rightarrow Z(f)$ is a cofibration.

Proposition 1.5. Let

$$\begin{aligned}
 j: X &\rightarrow Z(f) \\
 x &\mapsto (x, 0),
 \end{aligned}$$

we have

1. $j: X \rightarrow Z(f)$ is a cofibration.
2. $s \circ q \simeq \text{id}_{Z(f)} \text{ rel } Y$.
3. If f is a cofibration, then $q: Z(f) \rightarrow Y$ is a homotopy equivalence in \mathbf{TOP}^X .

Proof. (1). We construct a retraction $R: Z(f) \times I \rightarrow X \times I \cup Z(f) \times \{1\}$ as follow. Let $R': I \times I \rightarrow I \times \{1\} \cup \{0\} \times I$ be a retraction. Then we define

$$\begin{aligned}
 R: Z(f) \times I &\rightarrow X \times I \cup Z(f) \times \{1\} \\
 ((x, s), t) &\mapsto (x, R'(s, t)) \\
 (y, t) &\mapsto (y, 1)
 \end{aligned}$$

is as desired. By Proposition 1.2, $j: X \rightarrow Z(f)$ is a cofibration.

(2). The homotopy

$$\begin{aligned}
 h_t: Z(f) &\rightarrow Z(f) \\
 (x, \sigma) &\mapsto (x, (1-t)\sigma + t)
 \end{aligned}$$

is as desired.

(3). By Proposition 1.2, there is a retraction $r: Y \times I \rightarrow Z(f)$. Define

$$\begin{aligned} g: Y &\rightarrow Z(f) \\ y &\mapsto r(y, 1). \end{aligned}$$

One can verify that g is the homotopy inverse of q . □

Summery 1. Any map $f: X \rightarrow Y$ factors into

$$X \xrightarrow{j} Z \xrightarrow{q} Y$$

where $j: X \rightarrow Z$ is a cofibration and $q: Z \rightarrow Y$ is a homotopy equivalence. Moreover, such a factorization is unique up to homotopy equivalence. In particular, we can choose $Z = Z(f)$. We define $C_f = Z(f)/\text{im } j$ as the homotopy cofibre of f , i.e. $C_f = X \times I \sqcup Y/(x, 0) \sim *, (x, 1) \sim f(x)$, is called the mapping cone of f .

$$X \xrightarrow{f} Y \xrightarrow{s} C_f$$

1.1.3 The Cofibre Sequence (Puppe's Sequence)

To get finer structure, we work in \mathbf{TOP}^o . Given a map $f: (X, x_0) \rightarrow (Y, y_0)$, we get an induced map

$$\begin{aligned} f^*: [Y, B]^o &\rightarrow [X, B]^o \\ [\alpha] &\mapsto [f \circ \alpha], \end{aligned}$$

where $[X, B]^o$ is the homotopy class of basepoint preserving maps. In particular, we have the constant map

$$\begin{aligned} [*]: X &\rightarrow B \\ x &\mapsto b_0. \end{aligned}$$

Definition 1.6. We say a sequence

$$(X, x_0) \xrightarrow{f} (Y, y_0) \xrightarrow{g} (Z, z_0)$$

in \mathbf{TOP}^o is h-coexact if $\forall (B, b_0) \in \text{Ob}(\mathbf{TOP}^o)$,

$$[Z, B]^o \xrightarrow{g^*} [Y, B]^o \xrightarrow{f^*} [X, B]^o$$

is exact, i.e. $(f^*)^{-1}([*]) = \text{im } g^*$.

In \mathbf{TOP}^o , we consider the reduced mapping cone $CX := X \times I / X \times \{0\} \cup \{x_0\} \times I$. The basepoint of CX is $X \times \{0\} \cup \{x_0\} \times I$. And we consider the reduced mapping cone: For $f: (X, x_0) \rightarrow (Y, y_0)$, $C(f) := CX \vee Y/(x, 1) \sim f(x)$. It is equivalent to the following push-out diagram.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ i_1 \downarrow & & \downarrow f_1 \\ CX & \longrightarrow & C(f) \end{array}$$

In fact, f_1 maps y to $(y, 1)$.

We will also use symbol X instead of (X, x_0) in \mathbf{TOP}^o for short.

Proposition 1.7. The sequence

$$X \xrightarrow{f} Y \xrightarrow{f_1} C(f)$$

is h-coexact.

Proof. Consider the following sequence

$$[C(f), B]^o \xrightarrow{f_1^*} [Y, B]^o \xrightarrow{f^*} [X, B]^o$$

for any (B, b_0) .

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{f_1} & C(f) \\ & \searrow & \downarrow \alpha & \swarrow & \\ & & B & & \end{array}$$

Assume that $[\alpha] \in [Y, B]^o$ s.t. $[\alpha \circ f] = [*] \in [X, B]^o$, i.e. $\alpha \circ f$ is null-homotopic. This is equivalent that there exists a map $h: CX \rightarrow B$. The mapping cone $C(f)$ is the push-out of

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ i_1 \downarrow & & \downarrow f_1 \\ CX & \longrightarrow & C(f) \end{array}$$

Using the universal property of push-out, we have the following commutative diagram,

$$\begin{array}{ccccc} & & Y & & \\ & \nearrow f & & \searrow f_1 & \\ X & & & & C(f) \xrightarrow{\exists \beta} B \\ & \searrow i_1 & & \nearrow h & \\ & & CX & & \end{array}$$

i.e. $\alpha = \beta \circ f_1$. Therefore $[\alpha] = f_1^*[\beta]$ and this proposition follows. \square

Iterate the procedure, we get a long h-coexact sequence:

$$X \xrightarrow{f} Y \xrightarrow{f_1} C(f) \xrightarrow{f_2} C(f_1) \xrightarrow{f_3} C(f_2) \longrightarrow \dots$$

Consider the injection $j_1: CY \rightarrow C(f_1)$, we have that


$$C(f_1)/j_1(CY) = X \times I/X \times \partial I \cup \{x_0\} \times I = \Sigma X$$

is the reduced suspension of X . Then we get a quotient map

$$q(f): C(f_1) \rightarrow \Sigma X.$$

$$\begin{array}{ccccccc} \begin{array}{c} | \\ X \end{array} & \xrightarrow{f} & \begin{array}{c} | \\ Y \end{array} & \rightsquigarrow & \begin{array}{c} \triangle \\ C(f) \end{array} & \rightsquigarrow & \begin{array}{c} \triangle \\ C(f_1) \end{array} & \xrightarrow{q(f)} & \begin{array}{c} \triangle \\ \Sigma X \end{array} \end{array}$$

Claim 1. $q(f)$ is a homotopy equivalence.



Denote by $s(f): \Sigma X \rightarrow C(f_1)$ the homotopy inverse of $q(f)$. Then our original sequence becomes


$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{f_1} & C(f) & \xrightarrow{f_2} & C(f_1) \xrightarrow{f_3} C(f_2) \\ & & & & \searrow q(f) \circ f_2 & & \downarrow q(f) \\ & & & & & & \Sigma X \end{array}$$

Consider the following diagram.


$$\begin{array}{ccc} C(f_1) & \xrightarrow{f_3} & C(f_2) \\ q(f) \downarrow & \uparrow s(f) & \downarrow q(f_1) \\ \Sigma X & \xrightarrow{q(f_1) \circ f_3 \circ s(f)} & \Sigma Y \end{array}$$

Claim 2. Consider $\tau: \Sigma X \rightarrow \Sigma X$ which maps (x, t) to $(x, 1 - t)$, we have $q(f_1) \circ f_3 \circ s(f) \simeq \Sigma f \circ \tau$

To prove it, denote $p(f_1) = q(f_1) \circ f_3$. In fact, $p(f_1)$ retracts the left triangle, i.e. CX to a point.



In the following diagram, $s(f)$ is the union of id and $f \times \text{id}$, i.e. id maps the left triangle of ΣX to the left triangle of $C(f_1)$, $f \times \text{id}$ maps the right triangle of ΣX to the right triangle of $C(f_1)$. Then $\Sigma f = p(f_1) \circ s(f)$ naturally. Notice that τ flips ΣX left and right. Therefore, by symmetry, we have $p(f_1) \circ s(f) \simeq \Sigma f \circ \tau$, as desired.



Now we get

$$X \xrightarrow{f} Y \xrightarrow{f_1} C(f) \xrightarrow{p(f)} \Sigma X \xrightarrow{\Sigma f} \Sigma Y \xrightarrow{(\Sigma f)_1} C(\Sigma f)$$

Claim 3. There is a homeomorphism $\tau_1: C(\Sigma f) \rightarrow \Sigma C(f)$ such that the following diagram is commutative.

$$\begin{array}{ccc} \Sigma Y & \xrightarrow{(\Sigma f)_1} & C(\Sigma f) \\ & \searrow \Sigma f_1 & \downarrow \tau_1 \\ & & \Sigma C(f) \end{array}$$

In fact, regard both $C(\Sigma f)$ and $\Sigma C(f)$ as the quotient spaces of $X \times I \times I$ unioned with Y , τ_1 is induced from interchanging the two I -factors.

As conclusion, we have

Theorem 1.8 (Puppe's Sequence). The sequence

$$X \xrightarrow{f} Y \xrightarrow{f_1} C(f) \xrightarrow{p(f)} \Sigma X \xrightarrow{\Sigma f} \Sigma Y \xrightarrow{\Sigma f_1} \Sigma C(f) \xrightarrow{p(\Sigma f)} \Sigma^2 X \longrightarrow \Sigma^2 Y \longrightarrow \dots$$

is h-coexact.

1.2 Fibrations

Definition 1.9. A map $p: E \rightarrow B$ has the homotopy lifting property (HLP) for the space X if \forall homotopy $h: X \times I \rightarrow B$ and $a: X \rightarrow E$ s.t. $p \circ a(x) = h(x, 0)$, there exists a homotopy $H: X \times I \rightarrow E$ s.t. $p \circ H = h$. H is called a lifting of h .

$$\begin{array}{ccc} X & \xrightarrow{a} & E \\ i_0 \downarrow & \nearrow H & \downarrow p \\ X \times I & \xrightarrow{h} & B \end{array}$$

A map $p: E \rightarrow B$ is called a fibration if it has HLP for all spaces X .

Definition 1.10. Given maps $f: A \rightarrow B$ and $p: E \rightarrow B$. The pull-back of p along f is the terminal object of the following diagram,

$$\begin{array}{ccc} f^*E & \longrightarrow & E \\ \downarrow & & \downarrow p \\ A & \xrightarrow{f} & B \end{array}$$

i.e. for any C , $g: C \rightarrow E$, $h: C \rightarrow A$, there exists unique r such that the following diagram is commutative.

$$\begin{array}{ccccc} & & E & & \\ & \nearrow g & & \searrow p & \\ C & \xrightarrow{r} f^*E & & & B \\ & \searrow & & \nearrow f & \\ & & A & & \end{array}$$

Explicitly,

$$f^*E = \{(a, e) \in A \times E : f(a) = p(e)\}$$

and $\pi: f^*E \rightarrow A$ is the projection.

Denote $B^I = \text{Map}(I, B)$. Consider the pull-back

$$W(p) := \{(x, w) \in E \times B^I : p(x) = w(0)\}$$

which is given by the pull-back

$$\begin{array}{ccc} W(p) & \xrightarrow{k} & B^I \\ b \downarrow & & \downarrow e^0 \\ E & \xrightarrow{p} & B \end{array}$$

where e^0 maps w to $w(0)$.

Proposition 1.11. Given a map $p: E \rightarrow B$, the followings are equivalence:

1. $p: E \rightarrow B$ is a fibration.
2. p has HLP for $W(p)$.
- 3.

$$\begin{aligned} r: E^I &\rightarrow W(p) \\ \alpha &\mapsto (\alpha(0), p \circ \alpha) \end{aligned}$$

admits a section.

Proof. (1) \implies (2) is by definition.

(2) \implies (3): Because $W(p)$ is a pull-back, by its universal property, we have the following diagram and we want to find s such that $r \circ s = \text{id}$.

$$\begin{array}{ccccc} & & & B^I & \\ & & p^I \nearrow & & \searrow e^0 \\ E^I & \xrightleftharpoons[r]{s} & W(p) & \xrightarrow{k} & B \\ & \searrow e^0 & \downarrow b & & \nearrow p \\ & & E & & \end{array}$$

Notice that $\text{Map}(W(p), E^I) = \text{Map}(W(p) \times I, E)$, because p has HLP for $W(p)$, we have the following commutative diagram.

$$\begin{array}{ccc} W(p) & \xrightarrow{b} & E \\ \downarrow & \nearrow s & \downarrow p \\ W(p) \times I & \xrightarrow{k} & B \end{array}$$

We have $b \circ r \circ s = e^0 \circ s = b$ and $k \circ r \circ s = p^I s = k$. Using the universal property (uniqueness) of pull-back $W(p)$ for $W(p)$, we must have $r \circ s = \text{id}$, i.e. s is a section of r .

(3) \implies (1): Let s be the section of r . For any X, a, h as in the definition of fibration, we want to find H such that the following diagram is commutative.

$$\begin{array}{ccc} X & \xrightarrow{a} & E \\ i_0 \downarrow & \nearrow H & \downarrow p \\ X \times I & \xrightarrow{h} & B \end{array}$$

Using the universal property of pull-back $W(p)$, we have unique f such that the following diagram is commutative, where $h: X \rightarrow B^I$ is the same as $h: X \times I \rightarrow B$.

$$\begin{array}{ccccc}
 & & B^I & & \\
 & \nearrow h & & \searrow e^0 & \\
 X & \xrightarrow{\exists! f} & W(p) & \xrightarrow{k} & B \\
 & \searrow a & \downarrow b & \nearrow p & \\
 & & E & &
 \end{array}$$

Then because $\text{Map}(W(p), E^I) = \text{Map}(W(p) \times I, E)$, one can check that $H = s \circ f$ is as desired. In fact,

$$p \circ H(x, t) = (p \circ H(x))(t) = (k \circ r \circ s \circ f(x))(t) = (k \circ \text{id} \circ f(x))(t) = h(x, t)$$

and $H \circ i_0 = a$ is similar. \square

1.2.1 Pull-back of Fibration

Proposition 1.12. If $p: E \rightarrow B$ is a fibration, then $f^*E \rightarrow A$ is also a fibration.

Proof. In the following diagram, F is induced by HLP for fibration $p: E \rightarrow B$ and then H is induced by universal property of pull-back f^*E .

$$\begin{array}{ccccc}
 X & \xrightarrow{a} & f^*E & \xrightarrow{\pi} & E \\
 i_0 \downarrow & \nearrow H & \downarrow F & \nearrow \pi & \downarrow p \\
 X \times I & \xrightarrow{h} & A & \xrightarrow{f} & B
 \end{array}$$

\square

1.2.2 Replacing Maps by Fibration

Proposition 1.13. The evaluation $e^1: Y^I \rightarrow Y$, $w \mapsto w(1)$ is a fibration.

Proof. We can define H directly:

$$\begin{aligned}
 H: X \times I &\rightarrow Y^I \\
 (x, s) &\mapsto \begin{cases} [t \mapsto a|_X((1+s)t)], & \text{when } 0 \leq (1+s)t \leq 1 \\ [t \mapsto h(x, (1+s)t - 1)], & \text{when } (1+s)t \geq 1. \end{cases}
 \end{aligned}$$

$$\begin{array}{ccc}
 X & \xrightarrow{a} & Y^I \\
 i_0 \downarrow & \nearrow H & \downarrow e^1 \\
 X \times I & \xrightarrow{h} & Y
 \end{array}$$

\square

Given $f: X \rightarrow Y$, consider the following pull-back.

$$\begin{array}{ccc}
 W(f) = f^*Y^I & \xrightarrow{\quad} & Y^I \\
 i_0 \downarrow & & \downarrow e^1 \\
 X & \xrightarrow{f} & Y
 \end{array}$$

In fact,

$$W(f) = \{(x, w) \in X \times Y^I : f(x) = w(1)\}.$$

Denote $p: W(f) \rightarrow Y$, $(x, w) \mapsto w(1)$ and $s: X \rightarrow W(f)$, $x \mapsto (x, k_{f(x)})$ where $k_{f(x)}$ is a constant path at $f(x)$, and $q: W(f) \rightarrow X$, $(x, w) \mapsto x$. We can check that the following diagram is commutative.

$$\begin{array}{ccc} W(f) = f^*Y^I & \xrightarrow{\quad} & Y^I \\ i_0 \downarrow & \uparrow s & \downarrow e^1 \\ X & \xrightarrow{\quad f \quad} & Y \end{array}$$

Theorem 1.14. In the following commutative diagram,

$$\begin{array}{ccc} X & \xrightarrow{\quad s \quad} & W(f) \\ & \searrow f & \swarrow p \\ & & Y \end{array}$$

s is a homotopy equivalence and p is a fibration.

Proof. Consider the following fibration

$$\begin{array}{ccc} (f \times \text{id})^*Y^I & \xrightarrow{\quad} & Y^I \\ (q, p) \downarrow & & \downarrow (e^1, e^0) \\ X \times Y & \xrightarrow{\quad f \times \text{id} \quad} & Y \times Y \end{array}$$

Claim 4. $(f \times \text{id})^*Y^I = W(f)$.

To see that, notice that

$$(f \times \text{id})^*Y^I = \{(x, y, w) \in X \times Y \times Y^I : f(x) = w(1), y = w(0)\},$$

we can construct a map from $W(f)$ to $(f \times \text{id})^*Y^I$ that maps (x, w) to $(x, w(0), w)$. It's one to one.

Then $p: W(f) \rightarrow Y$ is a fibration if and only if $(f \times \text{id})^*Y^I \xrightarrow{(q, p)} X \times Y \xrightarrow{p_2} Y$ is a fibration. It's a composition of two fibration and then a fibration, as desired.

Claim 5. q is a homotopy inverse of s .

□

By this theorem, given any $f: X \rightarrow Y$, we can replace it by a fibration $p: W(f) \rightarrow Y$ homotopically. Then we can define the homotopy fibre at y_0 of $f: X \rightarrow Y$ to be

$$F(f) := p^{-1}(y_0) = \{(x, w) \in X \times Y^I : f(x) = w(1), y_0 = w(0)\}.$$

Remark 1.15. Apply HLP again, we can prove the factorization $f = s \circ p: X \rightarrow Y$ such that $s: X \rightarrow W$ is a homotopy equivalence and $p: W \rightarrow Y$ is a fibration. And this factorization is unique up to homotopy equivalence.

Theorem 1.16. Let $p: E \rightarrow B$ be a fibration and B is path-connected. Then all fibres $p^{-1}(b)$ are homotopy equivalent.

Proof. Given a path $\alpha: I \rightarrow B$, $\alpha(0) = b_0$ and $\alpha(1) = b_1$. Consider HLP property:

$$\begin{array}{ccc} p^{-1}(b_0) & \xrightarrow{\quad} & E \\ \downarrow & \nearrow H & \downarrow p \\ p^{-1}(b_0) \times I & \xrightarrow{h} & B \end{array}$$

where $h(x, t) = \alpha(t)$. Consider $H_1: p^{-1}(b_0) \rightarrow p^{-1}(b_1)$ the restriction of H at $t = 1$. Similarly, consider the reversed path $\bar{\alpha}$ of α , we get $\bar{H}_1: p^{-1}(b_1) \rightarrow p^{-1}(b_0)$.

Claim 6. $\bar{H}_1 \circ H_1 \simeq \text{id}$.

It's by applying homotopy lifting to the homotopy from $\bar{\alpha}\alpha$ to k_{b_0} . Therefore, all fibres $p^{-1}(b)$ are homotopy equivalent. \square

1.2.3 Fibre Exact Sequence (Puppe's Sequence)

Definition 1.17. We say a sequence of pointed maps

$$(X, x_0) \xrightarrow{f} (Y, y_0) \xrightarrow{g} (Z, z_0)$$

is h-coexact if $\forall (B, b_0)$, the induced sequence

$$[B, X]^o \xrightarrow{f_*} [B, Y]^o \xrightarrow{g_*} [B, Z]^o$$

is exact, i.e. $g_*^{-1}([c_{z_0}]) = \text{im } f_*$.

Recall the homotopy fibre of $f: X \rightarrow Y$ is

$$F(f) := p^{-1}(y_0) = \{(x, w) \in X \times Y^I : f(x) = w(1), y_0 = w(0)\}.$$

Denote $f^1: F(f) \rightarrow X$, $(x, w) \mapsto x$.

Proposition 1.18. For any $f: (X, x_0) \rightarrow (Y, y_0)$, the sequence

$$F(f) \xrightarrow{f^1} X \xrightarrow{f} Y$$

is h-coexact.

Proof. Assume $\alpha: B \rightarrow X$ satisfies $f \circ \alpha: B \rightarrow Y$ is null-homotopic and $f_*[\alpha] = [c_{y_0}]$. Apply HLP property:

$$\begin{array}{ccc} B & \xrightarrow{\quad} & FY = \{w \in Y^I : w(0) = y_0\} \\ \downarrow & \nearrow H & \downarrow e^1 \\ B \times I & \xrightarrow{h} & Y \end{array}$$

where h is a null-homotopy from $f \circ \alpha$ to c_{y_0} . Notice that $H_0: B \times \{1\} \rightarrow FY$ satisfies

$$\begin{array}{ccccc} & & FY & & \\ & \nearrow H_0 & & \searrow & \\ B & \xrightarrow{\beta} & F(f) & \xrightarrow{f^1} & X \\ & \searrow \alpha & & \nearrow & \\ & & X & & Y \end{array}$$

where β is induced by the universal property of the pull-back $F(f)$, such that $f^1 \circ \beta = \alpha$. Therefore, $f_*^1([\beta]) = [\alpha]$. \square

Iterate the procedure, we get a long h-exact sequence

$$\cdots \longrightarrow F(f^2) \xrightarrow{f^3} F(f^1) \xrightarrow{f^2} F(f) \xrightarrow{f^1} X \longrightarrow Y.$$

Question 1.19. How to understand $F(f^n) \xrightarrow{f^{n+1}} F(f^{n-1})$?

We consider the loop space

$$\Omega Y := \{w \in Y^I : w(0) = w(1) = y_0\}.$$

Notice that

$$(f^1)^{-1}(x_0) = \{(x, w) \in X \times Y^I : w(0) = y_0, w(1) = f(x_0) = y_0\},$$

we have $\Omega Y = (f^1)^{-1}(x_0)$. We write $i(f) : \Omega Y \rightarrow F(f)$ for the inclusion.

Theorem 1.20 (The puppe's fibre sequence). The sequence

$$\Omega^k F(f) \xrightarrow{\Omega^k f^1} \Omega^k X \xrightarrow{\Omega^k f} \Omega^k Y \xrightarrow{i(\Omega^{k-1} f)} \cdots \longrightarrow \Omega X \longrightarrow \Omega Y \longrightarrow F(f) \xrightarrow{f^1} X \longrightarrow Y$$

is h-exact.

Proof. Step 1:

$$\begin{aligned} F(f^1) &= \{(x, w, v) \in X \times Y^I \times X^I : w(0) = y_0, v(0) = x_0, w(1) = f(x), v(1) = x\} \\ &= \{(w, v) \in Y^I \times X^I : w(0) = y_0, v(0) = x_0, w(1) = f(v(1))\}. \end{aligned}$$

Define $j(f) : \Omega Y \rightarrow F(f^1)$, $w \mapsto (w, k_{x_0})$.

Claim 7. $j(f)$ is a homotopy equivalence.

In fact, define $r(f) : F(f^1) \rightarrow \Omega Y$, $(w, v) \mapsto w * \overline{(f \circ v)}$, then $r(f) \circ j(f) = \text{id}$. The homotopy from $\text{id}_{F(f^1)}$ to $j(f) \circ r(f)$ is $h_t(w, v) = (h_t^1, h_t^2)$, where $h_t^1(s) = \begin{cases} w(s(1+t)), & s(1+t) \leq 1, \\ f(v(2 - (1+t)s)), & s(1+t) \geq 1 \end{cases}$ and $h_t^2(s) = v(s(1-t))$.

Step 2: From $F(f^1) \xrightarrow{f^2} F(f) \xrightarrow{f^1} X$, we get

$$\begin{array}{ccc} F(f^2) & \xrightarrow{f^3} & F(f^1) \\ j(f^1) \uparrow & \nearrow i(f^1) & \uparrow j(f) \\ \Omega X & \xrightarrow{\Omega f} & \Omega Y \end{array}$$

Because $j(f^1)$ is a homotopy equivalence, we have $i(f^1) \simeq j(f) \circ \Omega f$.

Step 3: Now we have $\Omega X \xrightarrow{\Omega f} \Omega Y \xrightarrow{i(f)} F(f)$. Then we get $F \Omega f \longrightarrow \Omega X \xrightarrow{\Omega f} \Omega Y$.

Claim 8. $F(\Omega f)$ is homotopy equivalent to $\Omega F(f)$.

To see that, notice that $F(\Omega f)$ and $\Omega F(f)$ are all quotient of $\text{Map}(I \times I, Y)$.

Finally, we get the h-exact sequence

$$\Omega F(f) \longrightarrow \Omega X \longrightarrow \Omega Y \longrightarrow F(f) \longrightarrow X \longrightarrow Y.$$

□

1.3 Duality of Cofibration and Fibration

1.3.1 Duality of Reduced Suspension and Loop Space

Write $Y^X = \text{Map}(X, Y)$ equipped with compact-open topology. We define the adjunction

$$\begin{aligned} \alpha: Z^{X \times Y} &\rightarrow (Z^Y)^X \\ f &\mapsto [x \mapsto f(x, \cdot)]. \end{aligned}$$

Theorem 1.21. Suppose that X and Y are locally compact. Then α is a homeomorphism.

In the pointed version, we replace $X \times Y$ by $X \wedge Y = X \times Y / \{x_0\} \times Y \cup X \times \{y_0\}$ and $\text{Map}^o(X, Y)$ is the space of basepoint preserving maps. Then $\alpha^o: \text{Map}^o(X \wedge Y, Z) \rightarrow \text{Map}^o(X, \text{Map}^o(Y, Z))$ is a homeomorphism. Therefore, α^o induces a bijection $\alpha_*^o: [X \wedge Y, Z]^o \rightarrow [X, \text{Map}^o(Y, Z)]^o$.

Choose $Y = S^1 = I/\partial I$, then $X \wedge Y = X \times I / X \times \partial I \cup \{x_0\} \times I = \Sigma X$ is the reduced suspension of X and $\text{Map}^o(Y, Z) = \Omega Z$ is the loop space of Z . Therefore, we get a bijection $\alpha_*^o: [\Sigma X, Z]^o \rightarrow [X, \Omega Z]^o$.

On $[\Sigma X, Z]^o$, we have a group structure:

$$[f] +_M [g]: (x, t) \mapsto \begin{cases} f(x, 2t), & t \leq \frac{1}{2}, \\ g(x, 2t - 1), & t \geq \frac{1}{2}. \end{cases}$$

Let τ be the inversion of ΣX . For any $[f]$, $-[f] = [f \circ \tau]$.

On $[X, \Omega Z]^o$, we have

$$\begin{aligned} m: \Omega Z \times \Omega Z &\rightarrow \Omega Z \\ (u, v) &\mapsto u * v. \end{aligned}$$

Define

$$[f] +_m [g] := [m \circ (f \times g) \circ d],$$

where

$$\begin{aligned} d: X &\rightarrow X \times X \\ x &\mapsto (x, x) \end{aligned}$$

is the diagonal embedding.

One can verify that

$$\alpha_*^o([f] +_M [g]) = \alpha_*^o([f]) +_m \alpha_*^o([g]).$$

Then the adjunction map $\alpha_*^o: [\Sigma X, Z]^o \rightarrow [X, \Omega Z]^o$ is an isomorphism. In categorical language, this means $\text{Mor}(\Sigma X, Z) = \text{Mor}(X, \Omega Z)$ in \mathbf{TOP}^o . As conclusion, $\Sigma: \mathbf{TOP}^o \rightarrow \mathbf{TOP}^o$ and $\Omega: \mathbf{TOP}^o \rightarrow \mathbf{TOP}^o$ are dual functors.

1.3.2 Duality of HLP and HEP

Given a homotopy lifting diagram,

$$\begin{array}{ccc} X \times \{0\} & \longrightarrow & E \\ \downarrow & \nearrow & \downarrow p \\ X \times I & \longrightarrow & B \end{array}$$

notice that $\text{Map}(X \times I, Z) = \text{Map}(X, Z^I)$, it is equivalent to

$$\begin{array}{ccc} E & \xleftarrow{e^0} & E^I \\ \uparrow & \nearrow & \downarrow \\ X & \longrightarrow & B^I \end{array}$$

Dualize it, also by, $\text{Map}(X \times I, Z) = \text{Map}(X, Z^I)$, we have

$$\begin{array}{ccc} E' & \xrightarrow{i_0} & E' \times I \\ \downarrow & \nearrow & \uparrow \\ X' & \longleftarrow & B' \times I \end{array}$$

It is equivalent to

$$\begin{array}{ccccc} & & E' & & \\ & \nearrow & & \searrow & \\ B' & & & & X' \\ & \searrow & & \nearrow & \\ & & B' \times I & & \end{array}$$

$E' \times I \dashrightarrow X'$

which is the homotopy extension diagram.

1.3.3 Duality of Two Puppe's Sequences

Notice that $[\text{id}] \in [\Sigma X, \Sigma X]^o$, it induces $\alpha_*^o[\text{id}] = \eta: X \rightarrow \Omega \Sigma X$. For each map $f: X \rightarrow Y$, it induces

$$\begin{aligned} \eta: F(f) &\rightarrow \Omega C(f) \\ (x, w) &\mapsto \begin{cases} (x, 2t), & t \leq \frac{1}{2}, \\ w(2 - 2t), & t \geq \frac{1}{2}, \end{cases} \end{aligned}$$

where $C(f) = X \times I \sqcup Y / \{x_0\} \times I$, $f(x) \sim (x, 1)$ is the reduced cone of f . Then we get a diagram commutative up to homotopy.

$$\begin{array}{ccccc} \Omega Y & \longrightarrow & F(f) & \longrightarrow & X \\ \text{id} \downarrow & & \downarrow & & \downarrow \\ \Omega Y & \longrightarrow & \Omega C(f) & \longrightarrow & \Omega \Sigma X \end{array}$$

2 Homotopy Groups

2.1 Definitions and Properties


Given (X, x_0) , define n -th homotopy group

$$\pi_n(X, x_0) := [(I^n, \partial I^n), (X, x_0)],$$


where the identity element is the constant map and $[f] + [g]$ can be represented by

$$f +_i g: (t_1, \dots, t_n) \mapsto \begin{cases} f(t_1, \dots, 2t_i, \dots, t_n), & t_i \leq \frac{1}{2} \\ g(t_1, \dots, 2t_i - 1, \dots, t_n), & t_i \geq \frac{1}{2} \end{cases}$$

for any i . The following picture shows that $f +_i g$ and $f +_j g$ are homotopy equivalent for any $i \neq j$, where the red parts are mapped into the base point so the homotopies work. Sometimes, we write $\pi_n(X)$ for short.



Given a pair (X, A, x_0) , $J^n = \partial I^n \times I \cup I^n \times \{0\} = I^n - I^n \times \{1\} \subset I^{n+1}$,



define the $n + 1$ -th relative homotopy group to be

$$\pi_{n+1}(X, A, x_0) := [(I^{n+1}, \partial I^{n+1}, J^n), (X, A, x_0)].$$

Similarly, we sometimes use $\pi_{n+1}(X, A)$ for short.

Proposition 2.1. When $n \geq 2$, $\pi_n(X, x_0)$ and $\pi_{n+1}(X, A, x_0)$ are both abelian.

Proof. Exchanging f and g in the picture after the definition of $\pi_n(X, x_0)$, we can know that $\pi_n(X, x_0)$ is abelian for $n \geq 2$. For the relative case, we can not process homotopy in the top red region. But for $n \geq 3$, the squares of f and g should be cubes, then we can place the cubes in front and behind to get new homotopy. Therefore, $\pi_n(X, A, x_0)$ is abelian for $n \geq 3$. \square

Theorem 2.2 (Exact Homotopy Sequence). Given a pair (X, A) , we have a long exact sequence

$$\longrightarrow \pi_n(A, x_0) \xrightarrow{i_*} \pi_n(X, x_0) \xrightarrow{j_*} \pi_n(X, A, x_0) \xrightarrow{\partial} \pi_{n-1}(A, x_0) \longrightarrow \cdots \longrightarrow \pi_0(A, x_0) \xrightarrow{i_*} \pi_0(X, x_0),$$

where $j: (X, x_0, x_0) \rightarrow (X, A, x_0)$ is the inclusion and ∂ is induced from the restriction of I^n on $I^{n-1} \times \{1\}$.

Proof. Notice that each map $f: (I^n, \partial I^n) \rightarrow (X, x_0)$ induces a map

$$\begin{aligned} \overline{f_k}: I^{n-k} &\rightarrow \Omega^k(X, x_0) \\ (u_1, \dots, u_{n-k}) &\mapsto [(t_1, \dots, t_k) \mapsto f(t_1, \dots, t_k, u_1, \dots, u_{n-k})]. \end{aligned}$$

Then we get an isomorphism $\pi_n(X, x_0) \rightarrow \pi_{n-k}(\Omega^k X, c_{x_0})$. This is because $\pi_n(X, x_0) = [S^n, X]^o$ and $\Sigma S^{n-1} = S^n$, then $[S^n, X]^o = [\Sigma S^{n-1}, X]^o \cong [S^{n-1}, \Omega X]^o \cong [S^{n-k}, \Omega^k X]^o$ by duality (Section 1.3.1).

Given a pair (X, A) , the homotopy fibre of $\iota: A \hookrightarrow X$ is

$$F(\iota) = \{(a, w) \in A \times X^I : w(0) = x_0, w(1) = a\} = \{w \in X^I : w(0) = x_0, w(1) \in A\} := F(X, A).$$

Each map $f: (I^{n+1}, \partial I^{n+1}, J^n) \rightarrow (X, A, x_0)$ induces a map

$$\begin{aligned} \hat{f}: I^n &\rightarrow F(X, A) \\ (t_1, \dots, t_n) &\mapsto [t \mapsto f(t_1, \dots, t_n, t)], \end{aligned}$$

induces an isomorphism $\pi_{n+1}(X, A, x_0) \rightarrow \pi_n(F(X, A), x_0)$.

The fibre sequence of $\iota: A \hookrightarrow X$ is

$$\Omega^n F(\iota) \longrightarrow \Omega^n A \longrightarrow \Omega^n X \longrightarrow \dots \longrightarrow F(\iota) \longrightarrow A \xrightarrow{\iota} X.$$

Applying $[S^1, \cdot]^o$, we have

$$\begin{aligned} [S^1, \Omega^n F(\iota)]^o &= \pi_1(\Omega^n F(\iota)) = \pi_{n+1}(F(\iota)) = \pi_{n+2}(X, A), \\ [S^1, \Omega^n A]^o &= \pi_1(\Omega^n A) = \pi_{n+1}(A), \\ [S^1, \Omega^n X]^o &= \pi_1(\Omega^n X) = \pi_{n+1}(X). \end{aligned}$$

Then we get exact sequence

$$\pi_{n+2}(X, A) \longrightarrow \pi_{n+1}(A) \longrightarrow \pi_{n+1}(X) \longrightarrow \dots \longrightarrow \pi_1(X) \longrightarrow \pi_1(X, A) \longrightarrow \pi_0(A) \longrightarrow \pi_0(X),$$

where the exactness of the last a few places is straightforward to verify. \square

2.2 Change of Basepoint

Assume $v: I \rightarrow X$ is a continuous path with $v(0) = x_0$ and $v(1) = x_1$. We regard v as a homotopy

$$\begin{aligned} \hat{v}_t: I^n &\rightarrow X \\ u &\mapsto v(t). \end{aligned}$$

Note that $\partial I^n \hookrightarrow I^n$ is a cofibration (by Corollary 1.3), by HEP, we have the following commutative diagram,

$$\begin{array}{ccccc} & & \partial I^n \times I & & \\ & \nearrow & & \searrow & \\ \partial I^n & & & & I^n \times I \xrightarrow{\hat{v}_t} X \\ & \searrow & & \nearrow & \\ & & I^n & & \end{array}$$

f

where $[f] \in \pi_n(X, x_0)$.

Proposition 2.3. The map

$$\begin{aligned} v_\#: \pi_n(X, x_0) &\rightarrow \pi_n(X, x_1) \\ [v_0] &\mapsto [v_1] \end{aligned}$$

only depends on the homotopy class of v rel ∂_1 and defines an isomorphism.

Proof. Use HEP again. \square

Proposition 2.4. Suppose $f: (X, A) \rightarrow (Y, B)$ is a homotopy equivalence. Then $f_*: \pi_n(X, A, x_0) \rightarrow \pi_n(Y, B, f(x_0))$ is an isomorphism.

Proof. We only prove that homotopic maps induce isomorphic maps on π_n . Assume we have a homotopy $g_t: (X, A) \rightarrow (Y, B)$, we get a path in Y

$$\begin{aligned} w: I &\rightarrow Y \\ t &\mapsto g_t(x_0). \end{aligned}$$

Then we have the following commutative diagram by HEP.

$$\begin{array}{ccc} & & \pi_n(Y, B, g_0(x_0)) \\ & \nearrow^{g_{0,*}} & \downarrow w_* \\ \pi_n(X, A, x_0) & & \\ & \searrow_{g_{1,*}} & \downarrow \\ & & \pi_n(Y, B, g_1(x_0)) \end{array}$$

\square

Remark 2.5. By the proposition, we get a right action of $\pi_1(X, x_0)$ on $\pi_n(X, x_0)$.

2.3 Serre Fibration

Definition 2.6. We say $p: E \rightarrow B$ is a Serre fibration, if it has HLP for all cube $I^n, \forall n \geq 0$.

Theorem 2.7. Let $p: E \rightarrow B$ be a Serre fibration. Fix $b_0 \in B$ and $e_0 \in E$ such that $p(e_0) = b_0$. Given $B_0 \subset B$, write $E_0 = p^{-1}(B_0)$. Then $p_*: \pi_n(E, E_0, e_0) \rightarrow \pi_n(B, B_0, b_0)$ is an isomorphism for all $n \geq 1$.

Proof. **Surjectivity:** Given $h: (I^n, \partial I^n, J^{n-1}) \rightarrow (B, B_0, b_0)$. Consider the lifting problem.

$$\begin{array}{ccc} I^{n-1} \times \{0\} \cup \partial I^{n-1} \times I & \xrightarrow{c_{e_0}} & E \\ \downarrow & \nearrow H & \downarrow p \\ I^{n-1} \times I & \xrightarrow{h} & B \end{array}$$

Notice that $I^{n-1} \times \{0\} \cup \partial I^{n-1} \times I \cong I^{n-1} \times \{0\}$, the map of the first line is c_{e_0} . Then we have the lifting $H: I^n \rightarrow E$ such that $H(\partial I^n) \subset E_0 = p^{-1}(B_0)$ and $H(J^{n-1}) = e_0$.

Injectivity: Assume $p_*[f_0] = p_*[f_1]$. We get a homotopy $\phi_t: (I^n, \partial I^n, J^{n-1}) \rightarrow (B, B_0, b_0)$. Consider the lifting problem.

$$\begin{array}{ccc} I^n \times \partial I \cup J^{n-1} \times I & \xrightarrow{\quad} & E \\ \downarrow & \nearrow \phi & \downarrow \\ I^n \times I & \xrightarrow{\phi_t} & B \end{array}$$

Notice that $I^n \times \partial I \cup J^{n-1} \times I \cong I^n$, we have the lifting ϕ . \square

Corollary 2.8. Given a Serre fibration $F \hookrightarrow E \xrightarrow{p} B$ where F is a regular fibre, we have a long exact sequence

$$\pi_n(F) \xrightarrow{i_*} \pi_n(E) \xrightarrow{p_*} \pi_n(B) \longrightarrow \pi_{n-1}(F) \longrightarrow \cdots \longrightarrow \pi_0(E) \longrightarrow \pi_0(B).$$

Proof. Consider the pair (E, F) . By Theorem 2.2, we have exact sequence

$$\pi_n(F) \xrightarrow{i_*} \pi_n(E) \xrightarrow{p_*} \pi_n(B) \longrightarrow \pi_{n-1}(F) \longrightarrow \cdots$$

Choose $B_0 = b_0$ and $F = E_{b_0}$, we have $\pi_n(E, F, b_0) \cong \pi_n(E, b_0, b_0) \cong \pi_n(B, b_0)$ and this corollary follows. \square

Proposition 2.9. Every fibre bundle is a Serre fibration.

Proof. Given the lifting problem.

$$\begin{array}{ccc} I^n \times \{0\} & \xrightarrow{a} & E \\ \downarrow & \nearrow H & \downarrow \\ I^n \times I & \xrightarrow{h} & B \end{array}$$

We choose an open cover $\{U_\alpha\}_{\alpha \in \Lambda}$ of B such that finitely many U_α 's cover $\text{im } h$ and over each U_α , $E|_{U_\alpha}$ is trivialized. Choose a subdivision $\{I_\beta^n\}$ of I^n and partition $\{I_\lambda\}$ of I , such that $\forall \beta, \lambda, h(I_\beta^n \times I_\lambda) \subset U_\alpha$ for some α . Over each $I_\beta^n \times I_\lambda$, we consider

$$\begin{array}{ccc} I_\beta^n \times \partial I_\lambda \cup \partial I_\beta^n \times I_\lambda & \longrightarrow & U_\alpha \times F \\ \downarrow & \nearrow H_{\beta, \lambda} & \downarrow \\ I_\beta^n \times I_\lambda & \xrightarrow{h} & U_\alpha \end{array}$$

where $I_\beta^n \times \partial I_\lambda \cup \partial I_\beta^n \times I_\lambda \cong I_\beta^n \times \{0\}$ and $U_\alpha \times F \cong E|_{U_\alpha}$. We construct the lifting of h inductively on β and λ . \square

2.4 Higher Connectivity

Proposition 2.10. Let (X, A) be a pair, and $f: (I^n, \partial I^n) \rightarrow (X, A)$ a pointed map. The followings are equivalent.

1. f is null-homotopic.
2. f is homotopic rel ∂I^n to a map in A .

Proof. (1) \implies (2): Consider a surjective continuous map $\lambda: I^n \times I \rightarrow I^n \times I$ such that $\lambda|_{\partial I^n \times I}: (x, t) \mapsto (x, 0)$ and $\lambda|_{I \times \{0\}} = \text{id}_{I^n}$. Consider a null-homotopy $F: I^n \times I \rightarrow X$ of f , we let $H = F \circ \lambda: I^n \times I \rightarrow X$. Then H is a homotopy of f such that $H|_{\partial I^n \times \{t\}} = \text{id}_{\partial I^n}$ and $H_1(I^n) \subset A$.

(2) \implies (1): We may assume $f(I^n) \subset A$. J^{n-1} is a deformation retract of I^n . This is equivalent to that we get a homotopy $h_t: I^n \rightarrow I^n$ such that $\text{im } h_1 = J^{n-1}$ and $h_0 = \text{id}$. Then $f \circ h_t$ is a homotopy from f to c_{x_0} . \square

Remark 2.11. By (2), $\pi_n(A, A) \rightarrow \pi_n(X, A)$ is trivial.

Definition 2.12. We say a pair (X, A) is n -connected if $\pi_q(X, A) = 0$, $\forall 1 \leq q \leq n$ and $\pi_0(A) \rightarrow \pi_0(X)$ is surjective. Note that $\pi_q(X, A) = 0$ is computed for all basepoints.

Proposition 2.13. The followings are equivalent.

1. (X, A) is n -connected.
2. $j_*: \pi_q(A, *) \rightarrow \pi_q(X, *)$ is an isomorphism for $q < n$ and is an epimorphism for $q = n$.

Proof. The proof follows from exact sequence of the pair (X, A) (Proposition 2.2). \square

Definition 2.14. We say $f: X \rightarrow Y$ is n -connected if $f_*: \pi_k(X) \rightarrow \pi_k(Y)$ is an isomorphism for $1 \leq k \leq n-1$ and is an epimorphism for $k = n$.

Proposition 2.15. $f: X \rightarrow Y$ is n -connected if and only if $(Z(f), X)$ is n -connected.

Proof. The proof follows from exact sequence of the pair $(Z(f), X)$ (Proposition 2.2) and $Z(f) \simeq Y$. \square

2.5 Excision and Suspension

Theorem 2.16 (Blaskers-Massey). Let $Y = Y_1 \cup Y_2$ be union of two open subsets and $Y_0 = Y_1 \cap Y_2 \neq \emptyset$. Suppose $\pi_i(Y_1, Y_0) = 0$ for any $0 < i < p$, $p \geq 1$ and $\pi_j(Y_2, Y_0) = 0$ for any $0 < j < q$, $q \geq 1$. Then the map $\iota: \pi_n(Y_2, Y_0) \rightarrow \pi_n(Y, Y_1)$ is an isomorphism for $1 \leq n \leq p+q-3$ and is an epimorphism for $n = p+q-2$.

Proof. See textbook § 6.7. \square

Proposition 2.17. Let $j: A \hookrightarrow X$ be a cofibration. Consider a push-out diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ j \downarrow & & \downarrow J \\ X & \xrightarrow{F} & Y \end{array}$$

where $Y = X \sqcup B/f(a) \sim j(a)$. Suppose $\pi_i(X, A) = 0$, $\forall 0 < i < p$ and $\pi_i(Z(f), A) = 0$, $\forall 0 < i < q$. Then the induced map $(F, f)_*: \pi_n(X, A) \rightarrow \pi_n(Y, B)$ is an isomorphism for $1 \leq n \leq p+q-3$ and is an epimorphism for $n = p+q-2$.

Proof. Replace f by a cofibration

$$\begin{array}{ccccc} A & \xrightarrow{k} & Z(f) & \xrightarrow{p} & B \\ j \downarrow & & \downarrow & & \downarrow \\ X & \xrightarrow{K} & Z & \xrightarrow{P} & Y \end{array}$$

where $Z = Z(f) \sqcup X/(a, 0) \sim j(a)$, $f = p \circ k$, $F = P \circ K$. Since $p: Z(f) \rightarrow B$ is a homotopy equivalence and $P: Z \rightarrow Y$ is given by push-out, P is also a homotopy equivalence. Let $Z = Z_1 \cup Z_2$ where $Z_2 = X \sqcup A \times (\varepsilon, 1]/\sim$ and $Z_1 = B \sqcup A \times [0, \varepsilon]/\sim$. Then $Z_1 \cap Z_2 = A \times (\varepsilon, 1 - \varepsilon)$. Applying excision (Theorem 2.16),

$$\pi_n(X, A) \cong \pi_n(Z_2, Z_0) \rightarrow \pi_n(Z, Z_1) \cong \pi_n(Y, B)$$

has desired properties. \square

Theorem 2.18 (Quotient). Let $A \hookrightarrow X$ be a cofibration. Suppose $\pi_i(CA, A) = 0$ for $0 < i < p$ and $\pi_i(X, A) = 0$ for $0 < i < q$. Then $p_*: \pi_n(X, A) \rightarrow \pi_n(X/A, *)$ is an isomorphism for $1 \leq n \leq p+q-3$ and is an epimorphism for $n = p+q-2$.

Proof. Note $X \cup CA$ fits into the following push-out diagram.

$$\begin{array}{ccc} A & \longrightarrow & CA \\ \downarrow & & \downarrow \\ X & \longrightarrow & X \cup CA \end{array}$$

Then we get the result for

$$\pi_n(X, A) \rightarrow \pi_n(X \cup CA, CA).$$

Since $A \hookrightarrow X$ is a cofibration, $CA \hookrightarrow X \cup CA$ is also a cofibration. Notice that because CA is contractible, $X \cup CA \rightarrow X \cup CA/CA$ is a homotopy equivalence (This is left as an exercise). Then

$$\pi_n(X, A) \rightarrow \pi_n(X \cup CA, CA) \cong \pi_n(X \cup CA/CA, *) \cong \pi_n(X/A, *)$$

has desired properties. \square

Definition 2.19. We say (X, x_0) is well-pointed if $x_0 \hookrightarrow X$ is a cofibration.

Example 2.20. • For any CW-complex or manifold, it is well-pointed for any point.

- $X = \{\frac{1}{n} : n \in \mathbb{Z}^+\} \cup \{0\}$, $x_0 = 0$ is not well-pointed.

Theorem 2.21 (Freudenthal Suspension). Let (X, x_0) be a well-pointed n -connected space. Then $\Sigma_* : \pi_j(X) \rightarrow \pi_{j+1}(\Sigma X)$ is an isomorphism for $0 \leq j \leq 2n$ and is an epimorphism for $j = 2n + 1$.

Proof. The suspension map is given by

$$\pi_j(X) = [S^j, X]^o \xrightarrow{\Sigma_*} [S^{j+1}, \Sigma X]^o = \pi_{j+1}(X) .$$

We factor Σ_* into

$$\begin{array}{ccc} \Sigma_* : \pi_j(X) & \xleftarrow[\cong]{\partial} & \pi_{j+1}(CX, X) \\ & & \downarrow p_* \\ & & \pi_{j+1}(\Sigma X) \end{array}$$

To use Theorem 2.18, we verify $X \hookrightarrow CX$ is a cofibration. Consider the push-out diagram

$$\begin{array}{ccc} X \times \partial I \cup \{x_0\} \times I & \longrightarrow & X \\ \downarrow & & \downarrow \\ X \times I & \longrightarrow & CX \end{array}$$

where $CX = X \times I / X \times \{0\} \cup \{x_0\} \times I$. Because $\partial I \hookrightarrow I$ and $x_0 \hookrightarrow X$ are cofibrations, we have $\{x_0\} \times I \cup X \times \partial I \hookrightarrow X \times I$ is also a cofibration. By push-out diagram, $X \hookrightarrow CX$ is a cofibration. Now we have exact sequence

$$\begin{array}{ccc} \pi_j(CX, X) & \xrightarrow{\partial} & \pi_{j-1}(X) \longrightarrow 0 \\ \uparrow & & \uparrow \\ \pi_j(CX) & = & 0 \\ \uparrow & & \uparrow \\ \pi_j(X) & & \end{array}$$

Then (CX, X) is $(n+1)$ -connected. And $p_* : \pi_j(CX, X) \rightarrow \pi_j(\Sigma X)$ is isomorphism for $j \leq 2n - 1$ and is an epimorphism for $j = 2n$. Then we apply Theorem 2.18 with $p = q = n + 2$ and get the desired properties for $\Sigma_* : \pi_{j-1}(X) \rightarrow \pi_j(X)$. \square

2.6 Computation of Homotopy Groups

Example 2.22.

$$\pi_k(S^n) \cong \begin{cases} 0, & k < n \\ \mathbb{Z}, & k = n \end{cases} .$$

$$\pi_1(S^1) \cong \mathbb{Z}, \quad \pi_1(S^n) \cong 0, \quad \forall n \geq 2.$$

To compute $\pi_2(S^2)$, consider the Hopf fibration

$$S^1 \hookrightarrow S^3 \twoheadrightarrow S^2 .$$

This is given by the fibre bundle

$$S^2 = \mathbb{CP}^1 = \mathbb{C}^2 - \{0\}/\mathbb{C}^* = S^3/S^1.$$

We have the following fibre sequence

$$\begin{array}{ccccccc} \pi_2(S^1) & \longrightarrow & \pi_2(S^3) & \longrightarrow & \pi_2(S^2) & \xrightarrow{\partial} & \pi_1(S^1) \longrightarrow \pi_1(S^3) \\ \parallel & & & & & & \parallel & & \parallel \\ 0 & & & & & & \mathbb{Z} & & 0 \end{array}$$

Because S^1 is 0-connected, by Suspension Theorem, $\pi_1(S^1) \rightarrow \pi_2(S^2)$ is an epimorphism. Then $\pi_2(S^2) \cong \mathbb{Z}$ and $\pi_2(S^3) = 0$.

For $n \geq 2$, assume S^n is $(n-1)$ -connected, by Freudenthal's Suspension, $\pi_j(S^n) \rightarrow \pi_{j+1}(S^{n+1})$ is an isomorphism for $j \leq n \leq 2n$. By induction, $\pi_n(S^n) \cong \mathbb{Z}$ and $\pi_j(S^n) = 0$ for $j < n$.

Example 2.23. Notice that

$$\mathbb{CP}^n = \mathbb{C}^{n+1} - \{0\}/\mathbb{C}^* = S^{2n+1}/U(1)$$

for $n \geq 2$, we get a fibre bundle

$$U(1) \hookrightarrow S^{2n+1} \longrightarrow \mathbb{CP}^n.$$

Then we have fibre sequence

$$\pi_j(S^{2n+1}) \longrightarrow \pi_j(\mathbb{CP}^n) \longrightarrow \pi_{j-1}(U(1)) \longrightarrow \pi_{j-1}(S^{2n+1}).$$

Then when $j = 2$, $\pi_2(\mathbb{CP}^n) \cong \mathbb{Z}$. When $2 \neq j \leq 2n$, $\pi_j(\mathbb{CP}^n) = 0$.

Consider $\mathbb{CP}^\infty = \bigcup_{n \geq 1} \mathbb{CP}^n$,

$$\begin{array}{ccc} \mathbb{CP}^n & \hookrightarrow & \mathbb{CP}^{n+1} \\ \uparrow & & \uparrow \\ S^{2n+1} & \hookrightarrow & S^{2n+3} \\ \uparrow & & \uparrow \\ U(1) & & U(1) \end{array}$$

is induced from Five-Lemma. Then $i_*: \pi_2(\mathbb{CP}^n) \rightarrow \pi_2(\mathbb{CP}^{n+1})$ is an isomorphism. As conclusion,

$$\pi_n(\mathbb{CP}^\infty) \cong \begin{cases} \mathbb{Z}, & n = 2 \\ 0, & n \neq 2. \end{cases}$$

Example 2.24. We have the following fibre bundle by transitive group action.

$$O(n) \xrightarrow{j} O(n+1) \longrightarrow S^n.$$

Since S^n is $(n-1)$ -connected, the homotopy exact sequence for fibrations show $j: O(n) \hookrightarrow O(n+1)$ is $(n-1)$ -connected.

Write $O(\infty) = \bigcup_{n=1}^\infty O(n)$.

Theorem 2.25 (Bott-Periodicity).

$$\pi_k(O(\infty)) \cong \pi_{k+8}(O(\infty)).$$

Example 2.26 (Stiefel Manifolds). Denote $V_k(\mathbb{R}^n)$ be the orthogonal k -frames in \mathbb{R}^n . Then we have

$$V_k(\mathbb{R}^n) = O(n) / O(n-k).$$

Then we get a fibration

$$O(n-k) \hookrightarrow O(n) \twoheadrightarrow V_k(\mathbb{R}^n).$$

Notice that in

$$O(n-k) \xrightarrow{j} O(n-k+1) \hookrightarrow \cdots \hookrightarrow O(n),$$

j is $(n-k-1)$ -connected, then

$$\pi_i(O(n-k)) \xrightarrow{\cong} \pi_i(O(n)) \twoheadrightarrow \pi_i(V_k(\mathbb{R}^n))$$

for $i \leq n-k-2$. Therefore, $\pi_i(V_k(\mathbb{R}^n)) = 0$ when $i \leq n-k-1$.

Claim 9. $V_k(\mathbb{R}^n)$ is $(n-k-1)$ -connected.

Consider the projection

$$\begin{aligned} p: V_{k+1}(\mathbb{R}^{n+1}) &\rightarrow V_1(\mathbb{R}^{n+1}) \cong S^n \\ (v_1, \dots, v_{k+1}) &\mapsto v_{k+1}. \end{aligned}$$

The fibre is $V_k(\mathbb{R}^n)$. We know S^n is $(n-1)$ -connected, then $j: V_k(\mathbb{R}^n) \rightarrow V_{k+1}(\mathbb{R}^{n+1})$ is $(n-1)$ -connected. Therefore, we have $\pi_{n-k}(V_k(\mathbb{R}^n)) \cong \pi_{n-k}(V_2(\mathbb{R}^{n-k+2}))$. We know that $\pi_1(V_2(\mathbb{R}^{n-k+2})) = 0$. By Hurewicz Theorem, $H_i(V_2(\mathbb{R}^{n-k+2})) \cong \pi_i(V_2(\mathbb{R}^{n-k+2}))$ for $2 \leq i \leq n-k$, which is non-trivial. We will do these calculations later.

Part II

Generalized Homology

3 Homology Theory and CW-Complexes

3.1 Homology Theory

Denote $R - \mathbf{MOD}$ be the category of left R -modules and $\mathbf{TOP}(2)$ be the category of pairs (X, A) and

$$\begin{aligned} k: \mathbf{TOP}(2) &\rightarrow \mathbf{TOP}(2) \\ (X, A) &\mapsto (A, \emptyset) \end{aligned}$$

be the forgetful functor.

Definition 3.1 (Eilenberg-Steenrod Axioms). A homology theory on $\mathbf{TOP}(2)$ consists

1. a family of functors $h_n: \mathbf{TOP}(2) \rightarrow R - \mathbf{MOD}$,
2. a family of natural transformations $\partial_n: h_n \rightarrow h_{n-1} \circ k$ such that
 - (a) Homotopy invariance: $h_n(f_0) = h_n(f_1)$ for $f_0 \simeq f_1$.
 - (b) Exact sequence:

$$\cdots \longrightarrow h_{n+1}(X, A) \xrightarrow{\partial_{n+1}} h_n(A) \longrightarrow h_n(X) \longrightarrow h_n(X, A) \longrightarrow \cdots$$

for any pair (X, A) .

- (c) Excision: Given a pair (X, A) , for any $U \subset A$ such that $\bar{U} \subset \text{Int}(A)$, then inclusion induces an isomorphism $h_n(X - U, A - U) \rightarrow h_n(X, A)$.

Proposition 3.2. Given two pairs (X_i, A_i) , $i = 1, 2$, we get an isomorphism

$$\bigoplus_{i=1}^2 h_n(X_i, A_i) \rightarrow h_n(X_1 \sqcup X_2, A_1 \sqcup A_2).$$

Proof. Consider the commutative diagram for $A_i = \emptyset$.

$$\begin{array}{ccccc} h_n(X_1 \sqcup X_2, X_2) & & & & h_n(X_1 \sqcup X_2, X_1) \\ & \nwarrow j_1 & & \nearrow j_2 & \\ & & h_n(X_1 \sqcup X_2) & & \\ & \nearrow i_1 & & \nwarrow i_2 & \\ h_n(X_1) & & & & h_n(X_2) \end{array}$$

$\begin{array}{c} \uparrow a_1 \\ \downarrow a_2 \cong \end{array}$

Injectivity of $i_1 \oplus i_2$ is easy to check. For its surjectivity, take $c \in h_n(X_1 \sqcup X_2)$, we have $j_1(c) = j_1 \circ i_1 \circ a_1^{-1}(j_1(c))$. Then $c - i_1 \circ a_1^{-1}(j_1(c)) \in \ker j_1$. Therefore, there exists $x \in h_n(X_2)$ such that $i_2(x) = c - i_1(a_1^{-1} \circ j_1(c))$. Then $c = i_1(y) + i_2(x)$ where $y = a_1^{-1} \circ j_1(c) \in h_n(X_1)$.

The general case will be proved later. □

Let $A = *$ be a single point. Define $\tilde{h}(X) := h(X, *)$.

Assume there is a map $r: X \rightarrow A$ such that $r \circ i \simeq \text{id}$. Then $i_*: h_n(A) \rightarrow h_n(X)$ is injective. We get short exact sequences

$$0 \longrightarrow h_n(A) \xrightleftharpoons[r_*]{i_*} h_n(X) \longrightarrow h_n(X, A) \longrightarrow 0.$$

Then we have splitting $h_n(X) \cong h_n(A) \oplus h_n(X, A)$ and $h_n(X, A) = \ker r_*$. When $A = *$, take $r = c: X \rightarrow *$, then $\widetilde{h}_n(X) = h_n(X, *) = \ker(c_*: h_n(X) \rightarrow h_n(*))$.

Proposition 3.3. Let $A \hookrightarrow X$ be a cofibration. Then the quotient map induces an isomorphism $j_*: h_n(X, A) \rightarrow h_n(X/A, *)$.

Proof. Apply excision to $(X \cup CA, CA)$ for $U =$ the cone point of CA , we have $h_n(X, A) \cong h_n(X \cup CA, CA)$. When $A \hookrightarrow X$ is a cofibration, $CA \hookrightarrow X \cup CA$ is a cofibre. Since CA is contractible, $X \cup CA/CA \simeq X \cup CA$. Then $h_n(X \cup CA, CA) \cong h_n(X/A, *)$. \square

Proposition 3.4. Let $(X, *)$ and $(Y, *)$ be well-pointed spaces and $f: X \rightarrow Y$ is a pointed map. Then the cofibre sequence $X \xrightarrow{f} Y \xrightarrow{f^1} C(f)$ induces an exact sequence

$$\widetilde{h}_n(X) \xrightarrow{f_*} \widetilde{h}_n(Y) \xrightarrow{f_*^1} \widetilde{h}_n(C(f)) .$$

Proof. The proof follows the commutative diagrams

$$\begin{array}{ccccc} \widetilde{h}_n(X) & \longrightarrow & \widetilde{h}_n(Z(f)) & \longrightarrow & \widetilde{h}_n(Z(f), X) \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ \widetilde{h}_n(X) & \longrightarrow & \widetilde{h}_n(Y) & \longrightarrow & \widetilde{h}_n(C(f)) \end{array}$$

and

$$\begin{array}{ccc} X \times \partial I & \xrightarrow{(\text{id}, f)} & X \sqcup Y \\ \downarrow & & \downarrow \\ X \times I & \longrightarrow & Z(f) \end{array}$$

\square

Proposition 3.5. Given a triple (X, A, B) . Assume $B \hookrightarrow X$ is a cofibration, we get an exact sequence

$$\cdots \longrightarrow h_n(A, B) \longrightarrow h_n(X, B) \longrightarrow h_n(X, A) \xrightarrow{\partial} h_{n-1}(A, B) \longrightarrow \cdots .$$

Proof. Applying excision, we know that (X, A, B) and $(X \cup CB, A \cup CB, CB)$ have the same sequence. Applying homotopy equivalence, $(X \cup CB, A \cup CB, CB)$ and $(X, A, *)$ have the same sequence. The triple sequence of $(X, A, *)$ is the reduced pair sequence of (X, A) . \square

3.1.1 Suspension Isomorphism

Given a pair (X, A) , we have the suspension isomorphism

$$\sigma: h_n(X, A) \rightarrow h_n(\partial I \times X \cup I \times A, \{0\} \times X \cup I \times A)$$

by excision for $U = (0, 1] \times A \cup \{0\} \times X$. Consider the boundary map $\partial_{n+1}: h_{n+1}(I \times X, \partial I \times X \cup I \times A) \rightarrow h_n(\partial I \times X \cup I \times A, \{0\} \times X \cup I \times A)$. Notice that $X \simeq I \times X \simeq \{0\} \times X \cup I \times A$, we have the exact sequence

$$h_{n+1}(I \times X, \partial I \times X \cup I \times A) \xrightarrow{\partial_{n+1}} h_n(\partial I \times X \cup I \times A, \{0\} \times X \cup I \times A) \longrightarrow h_n(I \times X, \{0\} \times X \cup I \times A) = 0 .$$

Then ∂_{n+1} is an isomorphism and so is ∂_{n+1}^{-1} . We get isomorphisms

$$h_n(x, A) \longrightarrow h_n(\partial I \times X \cup I \times A, \{0\} \times X \cup I \times A) \xrightarrow{\partial_{n+1}^{-1}} h_{n+1}((I, \partial I) \times (X, A)) .$$

Choose $A = *$, define the suspension isomorphism by

$$\begin{array}{ccc} h_n(X, *) & \longrightarrow & h_{n+1}^\sigma(X \times I, \partial I \times X \cup I \times *) \\ \cong \downarrow & & \downarrow \text{quotient} \\ \widetilde{h}_n(X) & \xrightarrow{\tilde{\sigma}} & \widetilde{h}_{n+1}(\Sigma X) \end{array}$$

Assume $(X, *)$ is well-pointed, by Hurwicz map, we have the commutative diagram

$$\begin{array}{ccc} \pi_n(X) & \xrightarrow{\Sigma_*} & \pi_n(\Sigma X) \\ \downarrow & & \downarrow \\ \widetilde{h}_n(X) & \xrightarrow{\tilde{\sigma}} & \widetilde{h}_{n+1}(X) \end{array}$$

3.2 CW-Complex

Definition 3.6. We say X is obtained from A by attaching an n -cell if there exists a push-out diagram

$$\begin{array}{ccc} S^{n-1} & \xrightarrow{\varphi} & A \\ \downarrow & & \downarrow \\ D^n & \xrightarrow{\Phi} & X \end{array}$$

where φ is called attaching map and Φ is called characteristic map.

A CW-decomposition of (X, A) is a filtration $A = X^{-1} \subset X^0 \subset \dots \subset X$ such that

1. $X = \bigcup_{n \geq -1} X^n$,
2. X^n is obtained from X^{n-1} by attaching n -cells,
3. X carries the colimit topology (weak topology).

Proposition 3.7. Let (Y, B) be an n -connected pair, (X, A) be a relative CW-complex of dimension $\leq n$. Then each map $(F, f): (X, A) \rightarrow (Y, B)$ is homotopic rel. A to a map into B . When dimension $< n$, the homotopy class rel. A of maps $X \rightarrow B$ is unique.

Proof. Consider

$$\begin{array}{ccccc} \bigsqcup_k S_k^{q-1} & \longrightarrow & A & \longrightarrow & B \\ \downarrow & & \downarrow & & \downarrow \\ \bigsqcup_k D_k^q & \xrightarrow{\Phi^q} & X^q & \xrightarrow{F^q} & Y \end{array}$$

For any $q \leq n$, $\pi_q(Y, B) = 0$. Then $F^q \circ \Phi^q$ can be homotoped into B rel. $\bigsqcup_k S_k^{q-1}$.

$$\begin{array}{ccccc} \bigsqcup_k S_k^{q-1} & \longrightarrow & A & \longrightarrow & B \\ \downarrow & & \downarrow & \nearrow \text{dashed} & \downarrow \\ \bigsqcup_k D_k^q & \xrightarrow{\Phi^q} & X^q & \xrightarrow{F^q} & Y \end{array}$$

When dimension of $(X, A) < n$, apply the argument to $(X \times I, X \times \partial I \cup A \times I)$ which is a relative CW-complex of dimension $< n + 1$. \square

Theorem 3.8. Suppose $h: B \rightarrow Y$ is n -connected. Then for a CW-complex X , $h_*: [X, B] \rightarrow [X, Y]$ is bijective when $\dim X < n$ and surjective when $\dim X = n$.

Proof. We map replace Y by $Z(h)$: $B \longrightarrow Z(h) \xrightarrow{\cong} Y$.

Surjectivity: Let $A = \emptyset$. Apply Proposition 3.7 to $(X, \emptyset) \rightarrow (Z(h), B)$.

Injectivity: Apply Proposition 3.7 to $(X \times I, X \times \partial I)$. \square

Theorem 3.9 (Whitehead). Let $f: Y \rightarrow Z$ be a map between CW-complexes with $\dim Y, \dim Z \leq n \leq \infty$. If $f_*: \pi_q(Y) \rightarrow \pi_q(Z)$ is an isomorphism for $0 \leq q \leq n$, then f is a homotopy equivalence.

Proof. The map $f: Y \rightarrow Z$ is n -connected. By Theorem 3.8, $f_*: [Z, Y] \rightarrow [Z, Z]$ is surjective. Then there exists $g: Z \rightarrow Y$ such that $f \circ g \simeq \text{id}_Z$ and g is n -connected. Use Theorem 3.8 again, there exists $h: Y \rightarrow Z$ such that $g \circ h \simeq \text{id}_Y$. Therefore, g is a homotopy equivalence. \square

Theorem 3.10 (Suspension Theorem). Suppose Y is n -connected and X is a CW-complex. Then $\Sigma_*: [X, Y]^o \rightarrow [\Sigma X, \Sigma Y]^o$ is bijective if $\dim X \leq 2n$ and is surjective if $\dim X = 2n + 1$.

Proof. We know that $[\Sigma X, \Sigma Y]^o \cong [X, \Omega \Sigma Y]^o$. By Freudenthal's Suspension Theorem, $\Sigma_*: [S^k, Y]^o \rightarrow [S^{k-1}, \Sigma Y]^o$ is an isomorphism when $k \leq 2n$ and epimorphism if $k = 2n + 1$. Notice that $\pi_{k+1}(\Sigma Y) \cong \pi_k(\Omega \Sigma Y)$, $\sigma_*: [S^k, Y]^o \rightarrow [S^k, \Omega \Sigma Y]^o$ is adjoint to Σ_* and is reduced from

$$\begin{aligned} \sigma: Y &\rightarrow \Omega \Sigma Y \\ y &\mapsto [t \mapsto (y, t)]. \end{aligned}$$

Therefore, σ is $(2n + 1)$ -connected. Apply Theorem 3.8 to $\sigma_*: [X, Y]^o \rightarrow [X, \Omega \Sigma Y]^o$. \square

3.3 CW-Approximation

Proposition 3.11. Suppose X is obtained from A by attaching $(n+1)$ -cell. Then (X, A) is n -connected.

Proof. Consider the push-out diagram

$$\begin{array}{ccc} S^n & \longrightarrow & A \\ \downarrow & & \downarrow \\ D^{n+1} & \longrightarrow & X \end{array}$$

The Excision Theorem of push-out shows that $\pi_0(X, A) = 0$ and $\pi_q(D^{n+1}, S^n) = 0$ for any $1 \leq q \leq n$. Then $(\Phi, \varphi): (D^{n+1}, S^n) \rightarrow (X, A)$ is $(n - 1)$ -connected. When $k \leq n - 1$, $0 = \pi_k(D^{n+1}, S^n) \rightarrow \pi_k(X, A)$ is an isomorphism. \square

Theorem 3.12. Let $f: A \rightarrow Y$ be a k -connected map. Then for each $n > k$, there exists a relative CW-complex (X, A) with cells in $\dim \in \{k + 1, \dots, n\}$ and an n -connected extension $F: X \rightarrow Y$ of f .

Proof. When $n = 1$, $k = 0$, the proof is trivial. Consider $k = n - 1$, $n \geq 2$. Assume $f: A \rightarrow Y$ is $(n - 1)$ -connected. Replace Y by $Z(f)$:

$$\begin{array}{ccccc} A & \longrightarrow & Z(f) & \longrightarrow & Y \\ \downarrow & \nearrow & \nearrow & \nearrow & \nearrow \\ X & & & & \end{array}$$

Assume $f: A \rightarrow Y$ is an inclusion. Let $(\Phi_j, \varphi_j): (D^n, S^{n-1}) \rightarrow (Y, A)$ be a set of generators of $\pi_n(Y, A)$. Attach n -cells on A using φ_j . Regard Φ_j as a null-homotopy of $f \circ \varphi_j$. F is obtained by push-out property.

$$\begin{array}{ccccc} S^{n-1} & \xrightarrow{\varphi_j} & A & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \Phi_j & \nearrow & \nearrow \\ D^n & \longrightarrow & X & \xrightarrow{\quad} & \end{array}$$

And then $F_*: \pi_n(X, A) \rightarrow \pi_n(Y, A)$ is an epimorphism.

Consider the diagram

$$\begin{array}{ccccccccc}
\pi_n(A) & \longrightarrow & \pi_n(X) & \longrightarrow & \pi_n(X, A) & \longrightarrow & \pi_{n-1}(A) & \longrightarrow & \pi_{n-1}(X) & \longrightarrow & 0 \\
\downarrow \cong & & \downarrow F_* & & \downarrow F_* & & \downarrow \cong & & \downarrow F_* & & \downarrow \\
\pi_n(A) & \longrightarrow & \pi_n(Y) & \longrightarrow & \pi_n(Y, A) & \longrightarrow & \pi_{n-1}(A) & \xrightarrow{f_*} & \pi_{n-1}(Y) & \longrightarrow & 0
\end{array}$$

Notice that $F_*: \pi_n(X) \rightarrow \pi_n(Y)$ is also an epimorphism. Then by chasing diagram, we know that $F_*: \pi_{n-1}(X) \rightarrow \pi_{n-1}(Y)$ is an isomorphism. \square

Corollary 3.13. Given any space Y , there exists a CW-complex X and a map $F: X \rightarrow Y$ such that $F_*: \pi_n(X) \rightarrow \pi_n(Y)$ is an isomorphism for any $n \geq 0$. Such X is called a CW-approximation of Y .

Theorem 3.14. Let Y be a k -connected CW-complex. Then there exists a CW-complex X such that

1. X is homotopy equivalent to Y ;
2. $X^k = \{*\}$.

Proof. Apply Theorem 3.12 to $A = \{*\} \hookrightarrow Y$ which is a k -connected map. \square

3.4 Eilenberg-MacLane Space

3.4.1 Remarks about Compactly Generated Spaces

Definition 3.15. A Hausdorff space X is said to be compactly generated if for any compact subset K , a subset $A \subset X$ satisfies $A \cap K$ is closed, then A is closed in X .

Example 3.16. These spaces are compactly generated spaces:

- locally compact Hausdorff spaces,
- metric spaces,
- CW-complexes with finite cells in each dimension.

Given a Hausdorff space X , we can put a new topology \mathcal{T} on X by imposing:

$$A \subset X \text{ is } \mathcal{T}\text{-closed} \iff A \cap K \text{ is closed for any compact subset } K \subset X$$

such that X is compactly generated under \mathcal{T} .

Fact 3.17. If X, Y are both compactly generated spaces, then $X \times Y$ needs not to be compactly generated.

Definition 3.18. We denote by $X \times_k Y$ the product with compactly generated topology. We denote by $kF(X, Y)$ the space of continuous maps from X to Y , equipped the compactly generated topology.

Theorem 3.19. Let X, Y, Z be compactly generated spaces. Then

1. The evaluation map

$$\begin{aligned}
kF(Y, Z) \times_k Y &\rightarrow Z \\
(f, g) &\mapsto f(g)
\end{aligned}$$

is continuous.

2. The adjoint map

$$kF(X, kF(Y, Z)) \rightarrow kF(X \times_k Y, Z)$$

is a homeomorphism.

Proposition 3.20. Suppose $\pi_j(Y) = 0$ for $j > n$. Let X be obtained from A by attaching cells of $\dim \geq n + 2$. Then $\iota_*: [X, Y] \rightarrow [A, Y]$ is a bijection.

Proof. Surjectivity: Given $f: A \rightarrow Y$ and attaching map $\varphi: S^k \rightarrow A$, $k \geq n + 1$. Then $f \circ \varphi: S^k \rightarrow Y$ is null-homotopic which can be extended over X .

Injectivity: Apply the argument to $(X \times I, X \times \partial I \cup A \times I)$. \square

Definition 3.21. Let π be an abelian group. An Eilenberg-MacLane space of type $K(\pi, n)$ is a CW-complex such that

$$\pi_j(X) = \begin{cases} \pi, & j = n; \\ 0, & j \neq n. \end{cases}$$

Proposition 3.22. Suppose X_1, X_2 are $(n - 1)$ -connected CW-complex with $n \geq 2$. Then

$$\pi_n(X_1) \oplus \pi_n(X_2) \rightarrow \pi_n(X_1 \vee X_2)$$

is an isomorphism.

Proof. We can assume $X_i^{n-1} = \{*\}$ by CW-approximation. Therefore, cells in $X_1 \times X_2$ have dimension $0, n, \geq 2n$. Then $X_1 \times X_2$ is obtained from $X_1 \vee X_2$ by attaching cells of $\dim \geq 2n$. We have $\pi_n(X_1 \vee X_2) \rightarrow \pi_n(X_1 \times X_2) = \pi_n(X_1) \oplus \pi_n(X_2)$ is an isomorphism. \square

Theorem 3.23. Let X be a $(n - 1)$ -connected CW-complex. Suppose Y satisfies $\pi_j(Y) = 0, \forall j > n \geq 2$. Then the map $h_*: [X, Y]^o \rightarrow \text{Hom}(\pi_n(X), \pi_n(Y))$ is a bijection.

Proof. We can assume $X^{n-1} = \{*\}$ by Proposition 3.20. Then $[X, Y]^o = [X^{n+1}, Y]^o$. Notice that $\pi_n(X^{n+1}) = \pi_n(X)$, we only need to prove $h_X: [X^{n+1}, Y]^o \rightarrow \text{Hom}(\pi_n(X^{n+1}), \pi_n(Y))$ is a bijection.

We know $X^n = \bigvee_j S_j^n := B$. Applying homotopy, we may assume all attaching maps of $(n + 1)$ -cells are cased. Then X^{n+1} is the mapping cone $f: A := \bigvee_k S_k^n \rightarrow \bigvee S_j^n = B$.

We have the cofibre sequence

$$[A, Y]^o \longleftarrow [B, Y]^o \longleftarrow [X^{n+1}, Y]^o \longleftarrow [\Sigma A, Y]^o \longleftarrow \dots$$

Notice that

$$[\Sigma A, Y]^o = \left[\Sigma \bigvee_k S_k^n, Y \right]^o = \left[\bigvee_k \Sigma S_k^n, Y \right]^o = \left[\bigvee_k S_k^{n+1}, Y \right]^o = 0$$

because $[h] = \sum_k [h_k]$ and $\pi_{n+1}(Y) = 0$.

Claim 10.

$$\pi_n(A) \xrightarrow{f_*} \pi_n(B) \longrightarrow \pi_n(X^{n+1}) \longrightarrow 0$$

is exact.

Proof of Claim. Consider the push-out diagram:

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow i & & \downarrow \\ CA & \longrightarrow & X^{n+1} \end{array}$$

We know

$$\begin{array}{ccccccc} \pi_m(A) & \longrightarrow & \pi_m(CA) & \longrightarrow & \pi_m(CA, A) & \xrightarrow{\cong} & \pi_{m-1}(A) \longrightarrow 0 \\ & & \parallel & & & & \\ & & 0 & & & & \end{array}$$

Then $\pi_m(CA, A) = 0$ for any $m \leq n$. We know f is $(n-1)$ -connected. Applying excision, $\pi_m(CA, A) \rightarrow \pi_m(X^{n+1}, B)$ is an isomorphism for $m \leq 2n-1$. We have an exact sequence

$$\begin{array}{ccccccc} \pi_m(B) & \longrightarrow & \pi_m(X^{n+1}) & \longrightarrow & \pi_m(X^{n+1}, B) & \longrightarrow & \pi_{m-1}(B)0 \\ & & & & \parallel & & \\ & & & & 0 & & \end{array}$$

when $m \leq n$. Then

$$\begin{array}{ccccccc} \pi_{n+1}(CA, A) & \xrightarrow[\cong]{\text{excision}} & \pi_{n+1}(X^{n+1}, B) & \longrightarrow & \pi_n(B) & \longrightarrow & \pi_n(X) \longrightarrow 0 \\ \partial \downarrow \cong & & & & & & \\ \pi_n(A) & & & & & & \end{array}$$

□

Apply $\text{Hom}(-, \pi_n(Y))$, we get an exact sequence

$$\begin{array}{ccccccc} \text{Hom}(\pi_n(A), \pi_n(Y)) & \longleftarrow & \text{Hom}(\pi_n(B), \pi_n(Y)) & \longleftarrow & \text{Hom}(\pi_n(X^{n+1}), \pi_n(Y)) & \longleftarrow & 0 \\ h_A \uparrow & & h_B \uparrow & & h_X \uparrow & & \cong \uparrow \\ [A, Y]^o & \longleftarrow & [B, Y]^o & \longleftarrow & [X^{n+1}, Y]^o & \longleftarrow & 0 \end{array}$$

Claim 11. h_A and h_B are bijections.

Proof of Claim. We have

$$\begin{aligned} \text{Hom}(\pi_n(A), \pi_n(Y)) &= \text{Hom}\left(\pi_n\left(\bigvee_j S_j^n\right), \pi_n(Y)\right) = \text{Hom}\left(\bigoplus_j \pi_n(S_j^n), \pi_n(Y)\right) \\ &= \prod_j \text{Hom}(\pi_n(S_j^n), \pi_n(Y)) \cong \prod_j \pi_n(Y) \end{aligned}$$

and

$$[A, Y]^o = \left[\bigvee_j S_j^n, Y\right]^o = \prod_j [S_j^n, Y]^o = \prod_j \pi_n(Y).$$

□

Finally, by claim that $[X^{n+1}, Y]^o \rightarrow [B, Y]^o$ is injective, we get our conclusion by something like Five Lemma. □

Theorem 3.24. Let π be an abelian group and $n \geq 2$. Then the Eilenberg-MacLane space $K(\pi, n)$ exists and is unique up to homotopy.

Proof. Uniqueness: Assume X, Y are both $K(\pi, n)$. Then by Theorem 3.23,

$$h_X : [X, Y]^o \rightarrow \text{Hom}(\pi_n(X), \pi_n(Y)) = \text{Hom}(\pi, \pi)$$

is a bijection. Choose $f : X \rightarrow Y$ such that $h_X([f]) = \text{id}$. Then f is a weak homotopy equivalence. Whitehead Theorem gives us that f is in fact a homotopy equivalence.

Existence: Consider a free resolution

$$F_1 \longrightarrow F_0 \longrightarrow \pi \longrightarrow 0$$

with relators F_1 and generators F_0 . Construct X^{n+1} as the mapping cone of $g : F_1 \hookrightarrow \bigvee_k S_k^n \rightarrow \bigvee_j S_j^n \hookrightarrow F_0$. Therefore, X^{n+1} is $(n+1)$ -connected and $\pi_n(X^{n+1}) = \pi$. We attach cells of $\dim \geq n+2$ to eliminate $\pi_m(X)$ for $m \geq n+1$, by Zorn's Lemma, we finish our construction. \square

Definition 3.25. $K(\pi, 0) := \pi$ equipped with discrete topology. $K(\pi, 1)$ is constructed similar to Theorem 3.24, but the uniqueness will be proved later.

3.5 Spectral Homology

In this section, we assume that π is finitely generated and X is compactly generated.

Definition 3.26. A spectrum is a sequence of pairs $\{(E_n, e_n)\}_{n \geq 0}$ where $E(n)$ is a pointed space, $e_n : \Sigma E(n) \rightarrow E(n+1)$ is a pointed map. We say a spectrum is an Ω -spectrum if $\varepsilon_n : E(n) \rightarrow \Omega E(n+1)$ is a homotopy equivalence, where ε_n is the adjoint of e_n .

Example 3.27. 1. Sphere Spectrum: $E(n) = S^n$, $e_n : \Sigma S^n \rightarrow S^{n+1}$ is the identity map

$$\begin{aligned} \Sigma S^n &= S^n \wedge S^1 \cong S^{n+1} \\ \mathbb{R}^{n+1} \times I &\hookrightarrow \mathbb{R}^{n+2}. \end{aligned}$$

2. Eilenberg-MacLane Spectrum: Fix an abelian group π . Let $E(n) = K(\pi, n)$. Construct $e_n : \Sigma K(\pi, n) \rightarrow K(\pi, n+1)$ as follows:

(a) Milnor: $\Omega K(\pi, n+1)$ is a CW-complex. Then $[S^k, \Omega K(\pi, n+1)]^o = [S^{k+1}, K(\pi, n+1)]^o$ and then $\Omega K(\pi, n+1) \cong K(\pi, n)$. Define $e_n : \Sigma K(\pi, n) \rightarrow K(\pi, n+1)$ as the adjoint map; or

(b) Notice that $\pi_k(\Sigma K(\pi, n)) = \begin{cases} 0, & k \leq n \\ \pi, & k = n+1 \end{cases}$ because $\pi_k(K(\pi, n)) \rightarrow \pi_{k+1}(\Sigma K(\pi, n))$ is an isomorphism when $k \leq 2n-2$. Then $K(\pi, n+1)$ is obtained from $\Sigma K(\pi, n)$ by attaching cells of $\dim \geq n+3$. Take $e_n : \Sigma K(\pi, n) \rightarrow K(\pi, n+1)$ to be the inclusion map.

Definition 3.28. A reduced homology theory consists of a family of functors $\tilde{h}_n : \mathbf{TOP}^o \rightarrow R\text{-MOD}$ and isomorphisms $\sigma_n : \tilde{h}_n \rightarrow \tilde{h}_{n+1} \circ \Sigma$ that satisfy

1. Homotopy invariance: $\tilde{h}_n(f_0) = \tilde{h}_n(f_1)$ if $f_0 \simeq f_1$.
2. Exactness: each cofibre sequence

$$X \xrightarrow{f} Y \xrightarrow{f'} C(f)$$

induces an exact sequence

$$\tilde{h}_*(X) \longrightarrow \tilde{h}_*(Y) \longrightarrow \tilde{h}_*(C(f)) .$$

Remark 3.29. Unreduced theory \iff reduced theory. To see that, define $h_n(X) = \tilde{h}_n(X \sqcup \{*\})$ and $h_n(X, A) = \tilde{h}_n(C(X, A))$.

Let $E = \{(E(n), e_n)\}$ be a spectrum. We get suspension maps

$$[S^{n+k}, E(n) \wedge X]^o = \pi_{n+k}(E(n) \wedge X) \rightarrow \pi_{n+k+1}(E(n+1) \wedge X) = [S^{n+k+1}, E(n+1) \wedge X]^o$$

and

$$\Sigma(E(n) \wedge X) = S^1 \wedge (E(n) \wedge X) = \Sigma E(n) \wedge X.$$

Define $E_n(X) := \text{colim}_{k \rightarrow \infty} \pi_{n+k}(E(k) \wedge X)$, and $\sigma_n: E_n(X) \rightarrow E_{n+1}(\Sigma X)$ is defined via $[S^{n+k}, E(n) \wedge X] \rightarrow [S^{n+k+1}, E(n) \wedge \Sigma X]$.

Theorem 3.30. $\{(E_n(X), \sigma_n)\}$ defines a reduced homology theory.

Proof. Homotopy invariance is by definitions.

Injectivity of σ_n : Suppose $x \in \ker \sigma_n$, there exists $[f] \in [S^{n+k}, E(k) \wedge X]^o$ such that $[f]$ represents x and $f \wedge \text{id}_{S^1}: S^{n+k} \wedge S^1 \rightarrow (E(k) \wedge X) \wedge S^1$ is null-homotopic. Then

$$S^{n+k} \wedge S^1 \xrightarrow{f \wedge \text{id}} E(k) \wedge \Sigma X \xrightarrow{e_k \wedge \text{id}} E(k+1) \wedge X$$

is null-homotopic. Note that $[(e_k \wedge \text{id}) \circ (f \wedge \text{id})]$ represents x as well. We must have $x = 0$.

Surjectivity of σ_n : Given $g: S^{n+k+1} \rightarrow E(k) \wedge X \wedge S^1$. Then define

$$f: S^{n+k+1} \xrightarrow{g} E(k) \wedge X \wedge S^1 \xrightarrow{e_k} E(k+1) \wedge X$$

and we have $\sigma_n([f]) = [g]$.

Exactness of Cofibre Sequence: Consider

$$E_n(X) \xrightarrow{f_n} E_n(Y) \xrightarrow{f'_n} E_n(C(f)).$$

Suppose $z \in \ker f'_n$ and write $h: S^{n+k} \rightarrow E(k) \wedge Y$ to represent z . Then $(\text{id}_{E(k)} \wedge f') \circ h: S^{n+k} \rightarrow E(k) \wedge C(f)$ is null-homotopic. Consider cofibre sequences:

$$\begin{array}{ccccccc} S^{n+k} & \longrightarrow & C(\text{id}) & \longrightarrow & S^{n+k} \wedge S^1 & \longrightarrow & S^{n+k} \wedge S^1 \\ \downarrow h & & \downarrow H & & \downarrow \beta & & \downarrow h \wedge \text{id} \\ E(k) \wedge Y & \longrightarrow & C(\text{id} \wedge f) & \longrightarrow & E(k+1) \wedge X & \xrightarrow{\text{id} \wedge f} & E(k+1) \wedge Y \\ & & \downarrow \simeq & & & & \\ & & E(k) \wedge C(f) & & & & \end{array}$$

where H is given by null-homotopy of $(\text{id} \wedge f') \circ h$ and β is the quotient of H and the first two squares are commutative. These induce $h \wedge \text{id}$ such that the last square is commutative up to homotopy. Therefore, under $\text{colim}_{k \rightarrow \infty}$, we have

$$f_*[\beta] = [(\text{id} \wedge f) \circ \beta] = [h \wedge \text{id}] = [h].$$

□

Remark 3.31. In Example 3.27,

1. When $E = \{(S^n, \Sigma)\}_{n \geq 0}$,

$$E_n(X) = \text{colim}_{k \rightarrow \infty} \pi_{n+k}(S^k \wedge X) = \text{colim}_{k \rightarrow \infty} \pi_{n+k}(\Sigma^k X) = \pi_n^s(X),$$

which is the stable homotopy group.

2. When $E = \{(K(\mathbb{Z}, n), \sigma_n)\}_{n \geq 0}$,

$$E_n(X) = \operatorname{colim}_{k \rightarrow \infty} \pi_{n+k}(K(\mathbb{Z}, n) \wedge X) \cong \widetilde{H}_n(X, \mathbb{Z}),$$

which is the reduced singular homology.

Theorem 3.32 (Brown's Representation Theory). Let $\{(h_n, \partial_n)\}$ be a homology theory. Then there exists a spectrum $E = \{(E(n), e_n)\}$ and natural isomorphisms $h_n(X, A) \cong \operatorname{colim}_{k \rightarrow \infty} \pi_{n+k}(E(k) \wedge (X^+/A^+))$ for all finite CW-complexes (X, A) , where $X^+ = X \sqcup \{*\}$ and $A^+ = A \sqcup \{*\}$.

4 Cohomology

4.1 Axiomatic Cohomology

Definition 4.1. A cohomology theory consists of

1. a family of contravariant functors $h^n: \mathbf{TOP}(2) \rightarrow R\text{-MOD}$,
2. a family of natural transformations $\delta^n: h^{n-1} \circ K \rightarrow h^n$, where $K: (X, A) \rightarrow (A, \emptyset)$ is the restriction, that satisfy
 - (a) H-Invariance: $h^n(f_0) = h^n(f_1)$ if $f_0 \simeq f_1$.
 - (b) Exact Sequence: Given (X, A) ,

$$\cdots \longrightarrow h^{n-1}(A) \xrightarrow{\delta} h^n(X, A) \longrightarrow h^n(X) \longrightarrow h^n(A)$$

is exact.

- (c) Excision: Given a pair (X, A) with $U \subset A$ and $\overline{U} \subset \operatorname{Int}(A)$, then the restriction $h^n(X, A) \rightarrow h^n(X - U, A - U)$ is an isomorphism for any n .

Definition 4.2. A reduced cohomology theory is given by $\widetilde{h}^n(X) := \ker(h^n(X) \rightarrow h^n(\{*\}))$ which fits into a splitting exact sequence

$$0 \longrightarrow h^n(X, *) \longrightarrow h^n(X) \longrightarrow h^n(*) \longrightarrow 0.$$

And we have $\widetilde{h}^n(X) \cong h^n(X, *)$.

4.1.1 Mayer-Vietoris Sequence

Definition 4.3. Given $A, B \subset X$, we say the pair (A, B) is excisive if the restriction $h^*(A \cup B, A) \rightarrow h^*(B, A \cap B)$ is an isomorphism.

Lemma 4.4. The followings are equivalent:

1. (A, B) is excisive.
2. (B, A) is excisive.

Proof. The proof is given by chasing the following diagram.

$$\begin{array}{ccc}
h^*(A \cup B, A) & \xrightarrow{\quad} & h^*(B, A \cap B) \\
& \searrow & \nearrow \\
& h^*(A \cup B, A \cap B) & \\
& \nearrow & \searrow \\
h^*(A \cup B, B) & \xrightarrow{\quad} & h^*(A, A \cap B)
\end{array}$$

□

Part III

Characteristic Classes