

PSYC214: Statistics Lecture 5 – Interim summary – Part I

Michaelmas Term
Dr Richard Philpot
r.philpot@lancaster.ac.uk

1

1

WBA 1 – Q2.

• Young-min is examining the quality of care received by individuals suffering from depression. He compares the services from 10 different NHS clinics around the Lancaster area by asking their respective clients to rate their level of satisfaction on a scale from 1 to 7, 1 being not at all satisfied and 7 being extremely satisfied.



- A. Quality of care received
- B. Lancaster
- C. NHS clinics
- D. Individual suffering from depression
- E. Satisfaction scores

2

2

Variables

Independent Variable

• The variable the experimenter manipulates or changes, which may be assumed to have a direct effect (i.e., influences change) on the dependent variable.



Dependent Variable

• The outcome of interest. It is the variable being tested and measured in an experiment. It is 'dependent' on the effect (i.e., influence) of the independent variable.



3

3

WBA 1 – Q6.

- Under H_0 , the Null Hypothesis,
 - A. There is a significant difference between the conditions/groups
 - B. The samples come from different populations
 - C. The samples come from the same population
 - D. We accept the experimental hypothesis
 - E. We formulate a research hypothesis



4

4

Inferential statistics - Hypotheses

 H_0 the Null Hypothesis

- H_0 : there is no significant difference between the conditions/groups and the null hypothesis is accepted.
- Under H_0 , the samples come from the same population.

 H_1 the Experimental Hypothesis

- H_1 : there is a significant difference between the conditions/groups and the null hypothesis is rejected.
- Under H_1 , the samples come from the different populations.

5

5

WBA 1 – Q12.


- Matilda has three independent groups who each receive a different treatment to help them quit smoking – either hypnotherapy, psychotherapy or nicotine patches. To assess statistical differences between groups, Matilda should run:

- A. Three independent group t-tests
- B. A one-factor within-participant ANOVA
- C. A two-factor between-participant ANOVA
- D. A one-factor, between-participant ANOVA
- E. A set of descriptive statistics




6

6

Introduction to analysis of variance 


Factors and levels

- Factor: treatment
- 3 levels
 - Medication
 - Counselling
 - Exercise



7

7

WBA 1 – Q15. 

- The larger the F ratio


A. The larger the noise in a study

B. The smaller the signal in comparison to the noise

C. The larger the signal in comparison to the noise


D. Generally, the less likely your difference between groups will be significant


E. The harder your equation was to calculate



8

8

Introduction to analysis of variance 



$$F = \frac{\text{Signal}}{\text{Noise}}$$

$$F = \frac{\text{Signal}}{\text{Noise}}$$

The larger in magnitude the F value, the more treatment effects were standing out away from experimental error – i.e., the larger the signal is from the noise – the less likely that differences in scores are caused by chance


9

9

WBA 1 – Q18.

Which statement about the ANOVA is NOT true

- A. It is useful for examining mean differences between three or more groups
- B. It provides a greater risk of producing type I errors than a series of T-tests
- C. It requires at least one factor
- D. It requires continuous outcome (DV) data
- E. It is an Analysis of Variance



Lancaster University


10

10

Introduction to analysis of variance

Why conduct an analysis of variance?

- Compares means and variance
- Allows analysis of group differences for more than two groups
- Several means without inflating Type I error rate



Lancaster University


11

11

Introduction to analysis of variance

What do you need for a one factor between participants ANOVA?

- At least one categorical independent variable (i.e., one factor)
- One continuous dependent variable (outcome measure)



Lancaster University

12

12

PSYC214: Statistics Lecture 5 – Interim summary – Part II

Michaelmas Term
Dr Richard Philpot
r.philpot@lancaster.ac.uk

13

13

WBA 2 – Q8.

- What would be an example of a violation of the assumption of independence:
 - A. Participants are tested separately from one another.
 - B. We have disproportionately more females in one condition than another.
 - C. We have bimodal data.
 - D. We have unequal variance across our samples.
 - E. Our data have been transformed with mathematical functions.




14

14

Assumption of independence

What is it?

- Participants should be randomly assigned to a group
- Participants should not cluster, sharing a classification variable
 - Gender
 - Skill level
- There should be no influence across one data point to another



15

15

WBA 2 – Q12.

- How do planned comparisons and post hoc tests differ?
- A. Planned comparisons always happen before you run an ANOVA, whereas post hoc tests occur after.
- B. Post hoc tests always happen before you run an ANOVA, whereas planned comparisons occur after.
- C. Planned comparisons produce F statistics, whereas post hoc tests produce Q statistics.
- D. Planned comparisons test only the mean differences of interest, while post hoc tests compare all possible combinations of mean differences.
- E. They are interested in different studies



16

16

Planned comparisons

- Focused approach to examine specific group differences
- Perfect when certain hypotheses can be tested without comparing all combinations of means
- Should be pre-specified
- Need to keep the number of planned comparisons as low as possible to negate Type I errors – (number of levels – 1)

Group	\bar{A}_1	\bar{A}_2	\bar{A}_3	\bar{A}_4	\bar{A}_5
\bar{A}_1	-	-	-	-	-
\bar{A}_2		-	-	-	-
\bar{A}_3			-	-	-
\bar{A}_4				-	-
\bar{A}_5					-

17

17

Post hoc tests

- Post hoc comes from Latin for "after the event"
- Post hoc tests assess all possible combinations of differences between groups by comparing each mean with the other
- Make adjustments to p value, but more conservative than Bonferroni correction

Group	\bar{A}_1	\bar{A}_2	\bar{A}_3	\bar{A}_4	\bar{A}_5
\bar{A}_1	-	-	-	-	-
\bar{A}_2		-	-	-	-
\bar{A}_3			-	-	-
\bar{A}_4				-	-
\bar{A}_5					-

18

18

WBA 2 – Q15.

In the following image, what is the within-group variance (a.k.a error term):

- A. 611.3
B. 48.8
C. 237
D. 6.77e-06
E. 11571

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Group	2	1223	611.3	12.52	6.77e-06 ***
Residuals	237	11571	48.8		

19

19

The meaning of an ANOVA output

```
##          Df Sum Sq Mean Sq F value    Pr(>F)    ##
## Group      2    1223    611.3    12.52 6.77e-06 ***
## Residuals 237   11571     48.8                ##
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

$F = \frac{\text{between-group variance}}{\text{within-group variance}}$

$F = \frac{611.3}{48.8}$

$F = 12.52$

$p = 0.00000677$


20

20

WBA 2 – Q16.

If running 4 statistical tests, what is the approximate probability of receiving a type I error?

- A. 5%
B. 10%
C. 14%
D. 18.5%
E. 22.6%



21

21

The problem of multiple comparisons



- Type 1 error - 1 test at $p \leq 0.05 = 0.95$ (i.e., 5% chance we get noise)
- Type 1 error - 2 tests = $0.95 * 0.95 = 0.903$. (10% chance)
- Type 1 error - 3 tests = $0.95 * 0.95 * 0.95 = 0.857$ (14% chance)
- Type 1 error - 4 tests = $0.95 * 0.95 * 0.95 * 0.95 = 0.815$ (18.5% chance)
- Type 1 error - 5 tests = $0.95 * 0.95 * 0.95 * 0.95 * 0.95 = 0.774$ (22.6% chance)

22

22

WBA 2 – Q20.



- Which of the following is a type of data transformation?

- Tukey transformation.
- The square root transformation.
- The homogeneity transformation.
- The outlier removal.
- The post hoc transformation.




23

23

Dealing with 'rogue' data

Transforming data

- This involves taking every score from each participant and applying a uniform mathematical function to each
- Report both the original data and the transformed data



24

Dealing with 'rogue' data

How to transform data

Untransformed	Square-root transformed	Log transformed
38	6.164	1.580
1	1.000	0.000
13	3.606	1.114
2	1.414	0.301
13	3.606	1.114
20	4.472	1.301
50	7.071	1.699
9	3.000	0.954
38	6.164	1.580
6	2.449	0.778
4	2.000	0.602
43	6.557	1.638

<http://www.biostathandbook.com/transformation.html>

Type of Data Transformation	Nature of Data
Log Transformation $(\log(X_i))$	Whole numbers and cover wide range of values, small values with decimal fractions.
Square-root Transformation $(\sqrt{X_i})$	Small whole number & Percentage data where the range is between 0 and 30 % or between 70 and 100 %

Maidawad & Sananase (2014)

25

25

PSYC214: Statistics

Lecture 5 – Interim summary – Part III

Michaelmas Term
Dr Richard Philpot
r.philpot@lancaster.ac.uk


26

26

WBA 3 – Q2.

- Differences among participants in overall performance constitute a source of error for

- A participant
- A within-participant design
- A between-participant design
- Independent groups
- Random designs



27

27

Within-participants F ratio

Lancaster University

$$F = \frac{\text{between-group variance}}{\text{within-group variance}}$$

$$F = \frac{\text{between-group variance}}{\text{residual variance}}$$

We calculate the F ratio the same as for the between participants design, with the exception that we are not interested in how participants vary from one another!

We therefore include an additional step to remove the between-participant variability (we spoke of before) from the error term.

We remove the between-participant variability from the within-group variability – leaving only random errors behind – a.k.a., the residual variability

28


28

WBA 3 – Q9.

Lancaster University

- What is needed to calculate the between-group variance?

A. $SS_{BETWEEN}$ and the $df_{BETWEEN}$
 B. $SS_{RESIDUAL}$ and the $df_{BETWEEN}$
 C. $SS_{BETWEEN}$ and the $df_{RESIDUAL}$
 D. $SS_{RESIDUAL}$ and the $df_{RESIDUAL}$
 E. $SS_{BETWEEN}$ and the ΣY^2



29

29

What we'll need for the ANOVA

Lancaster University

$$F = \frac{\text{between-group variance}}{\text{residual variance}}$$

$$\text{between-group variance} = \frac{SS_{BETWEEN}}{df_{BETWEEN}} = \frac{47.18}{2} = 23.59$$

• $a - 1$ [i.e., number of levels - 1]

30

30

WBA 3 – Q13.

• **Low** between-participant variability means:

- A. our participants' means are similar
- B. our participants' means are very different
- C. the trends and direction of participant scores are consistent
- D. the trends and direction of participant scores are inconsistent
- E. the variability within a participant is volatile



31

31

Between-participant variability

Table 5. Burgers consumed before (A_1) and after (A_2) Conspiracy

	A_1	A_2	ΔA	P Mean
P1	5	3	-2	4
P2	9	7	-2	8
P3	3	1	-2	2
P4	7	5	-2	6
P5	4	6	2	5
A Mean	5.6	4.4		5

High between-participant variability

The extent to which participants, on average, differ from another regardless of their stage of the experiment

- In this example, there is wider variability between participant means.

32

32

Between-participant variability

The extent to which participants, on average, differ from another regardless of their stage of the experiment

- In this example, there is zero variability between participant means.
- Zero differences = zero variance.

Low between-participant variability


Table 6. Burgers consumed before (A_1) and after (A_2) Conspiracy

	A_1	A_2	ΔA	P Mean
P1	9	1	-8	5
P2	5	5	0	5
P3	4	6	2	5
P4	6	4	-2	5
P5	4	6	2	5
A Mean	5.6	4.4		5

33


33

WBA 3 – Q15.



In virtually all within-participants studies, we:


- A. would be predominately interested in between-participant differences
- B. would hypothesise the amount of change a participant would experience
- C. would hypothesise that a score at one time would significantly differ from a score at another
- D. would control for all extraneous variables
- E. would expect that we would violate the majority of data assumptions





34

34

Within-participants F ratio



Ways in which people can differ:


- Overall level of performance/score
- Trends in their scores (  )
- Both!



35


35

One factor within-participants ANOVA



Between-participant variability vs Residual variance

- In virtually all within-participant studies, we hypothesise that a score at one time would significantly differ from at another time.
- Less interested in the actual change in scores and not interested in between participant differences.
- As such, we are more interested in the residual variance than the between participant variability.



36

36

WBA 3 – Q19.

What is removed from our F ratio when calculating a within-participants F statistic?

- A. The p value
- B. The individual differences
- C. The group differences
- D. The error term
- E. The signal to noise ratio

Lancaster University

37

37

Between-participants F ratio

$$F = \frac{\text{between-group variance}}{\text{within-group variance}}$$

$$F = \frac{\text{treatment effects} + \text{experimental error}}{\text{experimental error} + \text{individual differences} + \text{random (residual) errors}}$$

$$F = \frac{\text{treatment effects} + \text{individual differences} + \text{random (residual) errors}}{\text{individual differences} + \text{random (residual) errors}}$$

Lancaster University

38

38

Within-participants F ratio


$$F = \frac{\text{between-group variance}}{\text{within-group variance}}$$

$$F = \frac{\text{treatment effects} + \text{random (residual) errors}}{\text{random (residual) errors}}$$

Lancaster University

39

39

Lancaster
University

PSYC214: Statistics

Lecture 5 – Interim summary – Part IV

Michaelmas Term
Dr Richard Philpot
r.philpot@lancaster.ac.uk

40
