

uniform distribution: (minimum, maximum)

$$\text{probability density function (PDF)} = \begin{cases} \frac{1}{\max - \min} & \min \leq x \leq \max \\ 0 & \text{otherwise} \end{cases}$$

$$\text{cumulative distribution function (CDF)} \\ P(X \leq x) = \begin{cases} \frac{x - \min}{\max - \min} & \min \leq x \leq \max \\ 1 & x > \max \end{cases}$$

poisson distribution: (mean = μ)

$$\text{pdf: } P(X = x) = \begin{cases} \frac{e^{-\mu} \mu^x}{x!} & (x \geq 0) \\ 0 & \text{otherwise} \end{cases}$$

$$\text{cdf: } P(X \leq x) = \begin{cases} Q(\lfloor x \rfloor + 1, \mu) & x \geq 0 \\ 0 & \text{otherwise} \end{cases} \quad \begin{matrix} (\lfloor x \rfloor \text{ is a floor function,} \\ Q(a, x) \text{ is the regularized incomplete gamma function}) \end{matrix}$$

binomial distribution: (number of trials " n " (positive integer), probability of success " p " (0 or 1))

$$\text{pdf: } P(X = x) = \begin{cases} \binom{n}{x} p^x (1-p)^{n-x} & 0 \leq x \leq n \\ 0 & \text{otherwise} \end{cases} \quad \binom{n}{m} \text{ is the binomial coefficient}$$

$$\text{cdf: } P(X \leq x) = \begin{cases} I_p(n - \lfloor x \rfloor, \lfloor x \rfloor + 1) & 0 \leq x < n \\ 1 & x \geq n \end{cases} \quad I_2(a, b) \text{ is the regularized incomplete beta function}$$

exponential distribution: (rate " λ " (positive))

$$\text{pdf: } \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

$$\text{cdf: } \begin{cases} 1 - e^{-\lambda x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Z distribution: (mean: $\mu=0$) (standard deviation: $\sigma=1$)

$$\text{pdf: } \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}}$$

$$\text{Z score: } Z = \frac{(x - \mu)}{\sigma} = \frac{(6 - 5)}{4} = 0.25$$

然后去 normal table 查相应数字

$$\text{cdf: } P(X \leq x) = \int_{-\infty}^x \frac{e^{-\frac{t^2}{2}}}{\sqrt{2\pi}} dt = \frac{1}{2} \operatorname{erfc}\left(-\frac{x}{\sqrt{2}}\right)$$

t distribution: (ν (positive))

$$\text{pdf: } \frac{\left(\frac{\nu}{\nu+1}\right)^{\frac{\nu+1}{2}}}{\sqrt{\nu} B(\nu/2, 1/2)}$$

$B(a, b)$ is the beta function

$$\text{cdf: } P(X \leq x) = \begin{cases} \frac{1}{2} I_{\frac{x}{x+V}} \left(\frac{V}{2}, \frac{1}{2} \right) & x \leq 0 \\ \frac{1}{2} \left(I_{\frac{x^2}{x^2+V}} \left(\frac{1}{2}, \frac{V}{2} \right) + 1 \right) & (\text{otherwise}) \end{cases}$$

confidence intervals: (with t distribution?)

$$\bar{x} \pm \frac{t_{(1-c/2)} s}{\sqrt{n}}$$

n : sample size s : sample standard deviation
 \bar{x} : sample mean c : confidence level

confidence intervals for proportion:

$$\hat{p} \pm \sqrt{\frac{(1-\hat{p})\hat{p}}{n}} Z_{(1-c/2)}$$

\hat{p} : sample proportion n : sample size
 c : confidence level

$$\hat{p} \pm (CV) \cdot SE(\hat{p})$$

↓ confidence value ↓ standard error

Confidence Intervals for z distribution:

$$\text{example: (95\% CI)} \quad \mu \pm 1.96 \frac{\sigma}{\sqrt{n}}$$

μ : mean Size: n
 σ : standard deviation

Confidence Intervals for t distribution:

$$\begin{cases} \alpha\text{-level} = 1 - \text{desired confidence level} \\ \text{df (degree of freedom)} = \text{sample size} - 1 \end{cases}$$

z-test:
 normal proportion
 population standard deviation
 t-test:
 normal proportion
 larger sample size

$$\text{Test statistic: } \frac{\bar{x} - \mu_0}{\frac{sd}{\sqrt{n}}}$$

μ_0 = H_0 value for population mean
 \bar{x} = sample mean
 n = sample size
 sd = population sd