

MedTourEasy Data Analytics Traineeship – April 2026

Name : PRAVEENKUMAR MOPURU
Email : mpk339@gmail.com
Mobile : 9985772150
Mentor : Mr. Ankit Hasija, Training Head
Course : MTE – Data Analytics Traineeship
Duration : 4 Weeks (April 2026)
Project Domain : Health Care Analytics
Study tools used : Pandas + Bayesian Statistics + Colab Environment

Project Github Link :

[https://github.com/M339KUMAR/ColabDemoFiles/blob/78482ca811fdcf319f162cef76ab319eecda5aea/MTE%20DA/MTE_DA_Apr2026\(Apr20th-1\).ipynb](https://github.com/M339KUMAR/ColabDemoFiles/blob/78482ca811fdcf319f162cef76ab319eecda5aea/MTE%20DA/MTE_DA_Apr2026(Apr20th-1).ipynb)

MTE Sponsored Coursera Google Data Analytics Professional Certificate course work Completed



9 Courses

- Foundations: Data, Data, Everywhere
- Ask Questions to Make Data-Driven Decisions
- Prepare Data for Exploration
- Process Data from Dirty to Clean
- Analyze Data to Answer Questions
- Share Data Through the Art of Visualization
- Introduction to Data Analysis Using Python
- Google Data Analytics Capstone: Complete a Case Study
- Accelerate Your Job Search with AI



Apr 13, 2026

PRAVEEN MOPURU

has successfully completed the online Professional Certificate

Google Data Analytics

Those who earn the Google Data Analytics Professional Certificate have completed nine courses, developed by Google, that include hands-on, practice-based assessments and are designed to prepare them for introductory-level roles in Data Analytics. They are competent in tools and platforms including spreadsheets, SQL, Tableau, and Python. They know how to prepare, process, analyze, and share data for thoughtful action.

Amanda Brophy
Amanda Brophy
Global Director of
Google Career
Certificates

Verify this certificate at:
<https://coursera.org/verify/professional-cert/WAPDME25N8R6>

This certificate attests to the learner's completion of an online program delivered via Coursera. It does not constitute formal enrollment at any university or entity and does not itself grant academic credit, grades, or a degree. Institutions or organizations may, at their discretion, recognize this learning toward their own programs or credentials.

Project & Dataset Description :

The Present Study of Healthcare Analytics domain is mainly aimed at the Claim (or) Hypothesis that survey conducted by the **Nation Geographic** at **1986** and study performed **in1990** on the data collected about the **Cause** of death of people who are left hander's at a particular age is **9** years early compared to that of the right hander's.



Project & Dataset Description :

Present Study uses 2 Datasets ::

1. ***lh_data.csv*** (3 Columns & 77 Rows) - Left Handedness Dataset
which describes the rate of left handedness across different ages, separated by Males & Females-
Age : Column Ranges from 10 to 86 in the order of youngest to oldest age,
Male : Proportion of Left hander's in Males at a given particular age,
Female : Proportion of Left hander's in Females at a given particular age.
2. ***cdc_vs00199_table310.tsv*** (4 Columns & 125 Rows) - Death Distribution Dataset
Age : Column represents Age group of people under study,
Male : Column indicates number of people who died at a particular age given they are Male ,
Female : Column indicates number of people who died at a particular age, given they are Female,
Both Sexes : Column indicates number of people died both Male & Female Combined

Tools used for Performing the current Study:

1. **Python Programming** for performing the operations/Task on the Given data,
2. **Pandas** Library which provides Series & Data-Frame data structures for loading the data,
3. **Numpy** Library for performing Numerical and Mathematical Operations on the given data,
4. **Bayesian Statistics & Probability** which uses the concept of Bayes Theorem

What is Bayes Theorem?

- Bayes Theorem is a method of calculating conditional probability.
- The traditional method of calculating conditional probability (the probability that one event occurs given the occurrence of a different event) is to use the conditional probability formula, calculating the joint probability of event one and event two occurring at the same time, and then dividing it by the probability of event two occurring.
- However, conditional probability can also be calculated in a slightly different fashion by using Bayes Theorem.

Bayes' theorem

- Bayes' theorem can be derived using product rule and conditional probability of event A with known event B:
- As from product rule we can write:
- $P(A \cap B) = P(A|B) P(B)$ or
- imilarly, the probability of event B with known event A:
- $P(A \cap B) = P(B|A) P(A)$
- Equating right hand side of both the equations, we will get:

$$P(A|B) = \frac{P(B|A) P(A)}{P(B)} \quad \dots(a)$$

Data Analysis Steps :

1. Ask
2. Prepare
3. Process
4. Analyze
5. Share
6. Act


Data Analysis Phase - 1 : Ask



“Left-handed people have a **9-year shorter** life expectancy than right-handed people.

The Population's Story

Who Dies, and When



Data Analysis Phase - 1 : Ask

Null Hypothesis H_0 :

There is no Significant difference in the average age at death between left handed individuals & right handed individuals,

Alternate Hypothesis H_1 :

The Average age at death for left handed individuals is 9 years earlier than the average age at death for right handed individuals.

w.r.t above hypothesis to perform the current study, asking for correct data sources and collecting the data necessary is a important step.

The primary source of the data is available and was provided by the MedTourEasy Team.

Dataset -1 : ***lh_data.csv*** (left handedness rates in Male vs Female)

Dataset-2 : ***cdc_vs00199_table310.tsv*** (*Death Dist Data in Male vs Female*)

Refer to slide-4 for details

Data Analysis Phase - 2 : Prepare

This phase of the analysis involves the loading the dataset details in to the Pandas DataFrame using the Pandas library

Both Datasets, Left handedness data & Death Distribution Data are loaded in to the DataFrame respectively for further steps in the analysis process,

!.Loading the Data

```
1  #Left Handers Data
2  data_url_1 = '/content/sample_data/lh_data.txt'
3
4  #Death Distribution Data
5  data_url_2 = '/content/sample_data/cdc_vs00199_table310.tsv'
```

```
1  df1= pd.read_csv(data_url_1)
2  df1
```

Data Analysis Phase - 2 : Prepare

Left Handedness DataFrame

```
1 df1= pd.read_csv(data_url_1)
2 df1
```

	Age	Male	Female
0	10	12.717558	12.198041
1	11	15.318830	11.144804
2	12	14.808281	11.549240
3	13	13.793744	11.276442
4	14	15.156304	11.572906
...	...	...	...
72	82	7.350204	4.874899
73	83	6.471204	5.672536
74	84	6.510858	5.774881
75	85	7.337968	7.051459
76	86	5.059387	4.680948

77 rows × 3 columns

Death Distribution DataFrame

```
1 df3 = pd.read_csv(data_url_2, sep='\t', skiprows=[1])
2 df3
```

	Age	Both Sexes	Male	Female
0	0	27937.0	15646.0	12291.0
1	1	1989.0	1103.0	886.0
2	2	1376.0	797.0	579.0
3	3	1046.0	601.0	445.0
4	4	838.0	474.0	364.0
...	...	...	...	...
120	120	1.0	NaN	1.0
121	121	NaN	NaN	NaN
122	122	NaN	NaN	NaN
123	123	NaN	NaN	NaN
124	124	NaN	NaN	NaN

125 rows × 4 columns

Data Analysis Phase - 3 : Process

This phase of the analysis primarily involves checking the data for Null Values, Missing Values, Range of the Data, Format or Data type of the columns in the dataset, and performing the necessary steps to make the data ready for analysis.

	Age	Both Sexes	Male	Female
114	114	6.0	NaN	6.0
115	115	3.0	NaN	3.0
116	116	NaN	NaN	NaN
118	118	1.0	NaN	1.0
119	119	2.0	NaN	2.0
120	120	1.0	NaN	1.0
121	121	NaN	NaN	NaN
122	122	NaN	NaN	NaN
123	123	NaN	NaN	NaN
124	124	NaN	NaN	NaN

Death Dist Dataset

```
1 #index of rows with NaN
2 rows_with_nan.index

Index([114, 115, 116, 118, 119, 120, 121, 122, 123, 124], dtype='int64')

1 #Number of rows with NaN
2 num_rows_with_nan = df3.isna().any(axis=1).sum()
3 num_rows_with_nan

np.int64(10)
```

```
1 df3.info()
```

```
<class 'pandas.core.frame.DataFrame'>
RangeIndex: 125 entries, 0 to 124
Data columns (total 4 columns):
#   Column          Non-Null Count  Dtype
---  ---
0   Age             125 non-null   int64
1   Both Sexes      120 non-null   float64
2   Male            115 non-null   float64
3   Female          120 non-null   float64
dtypes: float64(3), int64(1)
memory usage: 4.0 KB
```

```
1 df4=df3.dropna()
```

```
<class 'pandas.core.frame.DataFrame'>
Index: 115 entries, 0 to 117
Data columns (total 4 columns):
#   Column          Non-Null Count  Dtype
---  ---
0   Age             115 non-null   int64
1   Both Sexes      115 non-null   float64
2   Male            115 non-null   float64
3   Female          115 non-null   float64
dtypes: float64(3), int64(1)
memory usage: 4.5 KB
```


Data Analysis Phase - 4 : Analyze

Task 1: Instructions

Load the handedness data from the National Geographic survey and create a scatter plot.

- Import `pandas` as `pd` and `matplotlib.pyplot` as `plt`.
- Load the data into a pandas DataFrame named `lefthanded_data` using the provided `data_url_1`. Note that the file is a CSV file.
- Use the `.plot()` method to create a plot of the "Male" and "Female" columns vs. "Age"

```
1 plt.plot(lefthanded_data.Age, lefthanded_data.Male, marker='o', label='Male')
2 plt.plot(lefthanded_data.Age, lefthanded_data.Female, marker='o', label='Female')
3 plt.title("Male vs Female", size=25, color='Red')
4 plt.xlabel('Age', size=15, color='Green')
5 plt.ylabel('Gender', size=15, color='Green')
6 plt.legend()
7 plt.axis()
8 plt.show()
```




Data Analysis Phase - 4 : Analyze


Task 2: Instructions

Add two new columns, one for birth year and one for mean left-handedness, then plot the mean as a function of birth year.

- Create a column in `lefthanded_data` called `Birth_year`, which is equal to 1986 - Age (since the study was done in 1986).
- Create a column in `lefthanded_data` called `Mean_lh` which is equal to the mean of the `Male` and `Female` columns.
- Use the `.plot()` method to plot `Mean_lh` vs. `Birth_year`.



```
1 plt.plot(lefthanded_data.Age, lefthanded_data.Mean_lh, marker='o', label='Mean')
2 plt.title('Age vs Mean_lh', size=20, color = 'Red')
3 plt.xlabel("Age", color='green', size=15)
4 plt.ylabel("Mean Age", color= 'green', size=15)
5 plt.legend()
6 plt.axis()
7 plt.show()
```



Data Analysis Phase - 4 : Analyze

Task 3: Instructions

Create a function that will return $P(LH | A)$ for particular ages of death in a given study year.

- Import the `numpy` package aliased as `np`.
- Use the last ten `Mean_lh` data points to get an average rate for the early 1900s. Name the resulting DataFrame `early_1900s_rate`.
- Use the first ten `Mean_lh` data points to get an average rate for the late 1900s. Name the resulting DataFrame `late_1900s_rate`.
- For the early 1900s ages, fill in `P_return` with the appropriate left-handedness rates for `ages_of_death`. That is, input `early_1900s_rate` as a fraction, i.e., divide by 100.
- For the late 1900s ages, fill in `P_return` with the appropriate left-handedness rates for `ages_of_death`. That is, input `late_1900s_rate` as a fraction, i.e., divide by 100.

When calculating `early_1900s_rate` and `late_1900s_rate`, remember that because the original data was from youngest age to oldest age, that means that the data is organized from latest birth year to earliest birth year. You will use the first ten `Mean_lh` data points to get an average rate for the late 1900s and the last ten for the early 1900s.

Data Analysis Phase - 4 : Analyze

Task 3:

```
1 import numpy as np
```

```
1 def P_lh_given_A(ages_of_death, study_year= 1990):
2
3     """
4     inputs: death age & study year
5     output: Prob of left handed
6             given that a subject died
7             in study-year at 'ages_of_death'
8     """
9
10    early_1900s_rate = lefthanded_data['Mean_lh'][-10:].mean()
11    late_1900s_rate = lefthanded_data['Mean_lh'][:10].mean()
12
13    Y_age = 10 + (study_year - 1986)
14    O_age = 86 + (study_year - 1986)
15
16    Mid_Rate = lefthanded_data.loc[lefthanded_data['Birth_year'].isin(study_year-ages_of_death)]['Mean_lh']
17
18    P_return = np.zeros(ages_of_death.shape)
19
20    P_return[ages_of_death > O_age] = early_1900s_rate/100
21    P_return[ages_of_death < Y_age] = late_1900s_rate/100
22
23    P_return[np.logical_and((ages_of_death <= O_age), (ages_of_death >= Y_age))] = Mid_Rate/100
24
25    return P_return
```

Data Analysis Phase - 4 : Analyze

Task 4: Instructions

Load death distribution data for the United States and plot it.

- Load death distribution data in the provided `data_url_2` into `death_distribution_data`, setting `sep = '\t'` and `skiprows=[1]` to account for the dataset's format.
- Drop the NaN values from the `Both Sexes` column.
- Use the `.plot()` method to plot the number of people who died as a function of their age.

Task 4:

```
1 #Loading Death Distribution Data in the given variable
2
3 death_distribution_data = pd.read_csv(data_url_2, sep='\t', skiprows = [1])
4 death_distribution_data
```


...

	Age	Both Sexes	Male	Female
0	0	27937.0	15646.0	12291.0
1	1	1989.0	1103.0	886.0
2	2	1376.0	797.0	579.0
3	3	1046.0	601.0	445.0
4	4	838.0	474.0	364.0
...	...	...	...	...
120	120	1.0	NaN	1.0
121	121	NaN	NaN	NaN
122	122	NaN	NaN	NaN
123	123	NaN	NaN	NaN
124	124	NaN	NaN	NaN

125 rows × 4 columns

Data Analysis Phase - 4 : Analyze

```
1 plt.figure(figsize=(25,10))
2 plt.plot(death_distribution_data.Age, death_distribution_data['Both Sexes'], marker='o', markersize=5, linewidth=2, label='Age vs Total Deaths')
3 plt.title('Age vs Both Seces', size=20, color='Red')
4 plt.xlabel('Age', size=15, color='green')
5 plt.ylabel('Both Sexes', size=15, color='green')
6 plt.legend()
7 plt.show()
```



Data Analysis Phase - 4 : Analyze **Task 5: Instructions**

Create a function called `P_lh()` which calculates the overall probability of left-handedness in the population for a given study year.

- Create a series, `p_list`, by multiplying the number of dead people in the `Both Sexes` column with the probability of their being lefthanded using `P_lh_given_A()`.
- Set the variable `p` equal to the sum of that series.
- Divide `p` by the total number of dead people by summing `death_distribution_data` over the `Both Sexes` column. Return result from the function.

$P(LH | A)$ was defined in Task 3. $N(A)$ is the value of `Both Sexes` in the `death_distribution_data` DataFrame where the `Age` column is equal to `A`. The denominator is total number of dead people, which you can get by summing over the entire dataframe in the `Both Sexes` column.

Task 5:

```
1 def P_lh(death_distribution_data, study_year=1990):
2     """
3     Overall Prob of left-handedness
4     in the population for a given study year
5
6     input  : death_dist_data, study_year
7     output : prob of death as lefthanded
8     """
9
10    p_list = death_distribution_data['Both Sexes']*P_lh_given_A(death_distribution_data['Age'], study_year)
11    p = np.sum(p_list)
12
13    return p/np.sum(death_distribution_data['Both Sexes'])
```

Data Analysis Phase - 4 : Analyze

Task 6: Instructions

Write a function to calculate `P_A_given_lh()`.

- Calculate `P_A`, the overall probability of dying at age A, which is given by `death_distribution_data` at age A divided by the total number of dead people (the sum of the `Both Sexes` column of `death_distribution_data`).
- Calculate the overall probability of left-handedness $P(LH)$ using the function defined in Task 5.
- Calculate $P(LH | A)$ using the function defined in Task 3.

Task 6:

```
1 def P_A_given_lh(ages_of_death, death_distribution_data, study_year=1990):
2     """
3     P_A, The Overall Probability of
4     dying at age A, which is given
5     by death_distribution_data at
6     age A divided by total number of
7     dead people who are left handed.
8
9     input  : ages_of_death, study_year
10    output : prob of dying at age A given lefthanded
11    """
12
13    P_A = death_distribution_data['Both Sexes'][ages_of_death]/np.sum(death_distribution_data['Both Sexes'])
14
15    P_left = P_lh(death_distribution_data, study_year)
16
17    P_lh_A = P_lh_given_A(ages_of_death, study_year)
18
19    return (P_lh_A*P_A)/P_left
```

Data Analysis Phase - 4 : Analyze

Task 7: Instructions

Write a function to calculate `P_A_given_rh()`.

- Calculate `P_A`, the overall probability of dying at age A, which is given by `death_distribution_data` at age A divided by the total number of dead people. (This value is the same as in task 6.)
- Calculate the overall probability of right-handedness $P(RH)$, which is $1 - P(LH)$.
- Calculate $P(RH | A)$, which is $1 - P(LH | A)$.

Task 7:

```
1 def P_A_given_rh(ages_of_death, death_distribution_data, study_year=1990):
2     """
3     The Overall Probability of
4     dying at age A, which is given
5     by death_distribution_data at
6     age A divided by total number of
7     dead people who are right handed.
8
9     input  : ages_of_death, study_year
10    output : prob of dying at age A given lefthanded
11    """
12
13    P_A = death_distribution_data['Both Sexes'][ages_of_death]/np.sum(death_distribution_data['Both Sexes'])
14
15    P_right = 1 - P_lh(death_distribution_data, study_year)
16
17    P_rh_A = 1 - P_lh_given_A(ages_of_death, study_year)
18
19    return (P_rh_A*P_A)/P_right
```

Data Analysis Phase - 4 : Analyze

Task 8: Instructions

Plot the probability of being a certain age at death given that you're left- or right-handed for a range of ages.


- Calculate `P_A_given_lh` and `P_A_given_rh` using the functions defined in Task 6.
- Use the `.plot()` method to plot the results versus age.

Data Analysis Phase - 4 : Analyze

Task 8:

```
1  ages = np.arange(0, 115,1)
2  #print(ages)
3
4  left_handed_prob = P_A_given_lh(ages, death_distribution_data, 1990) # for age in ages ]
5  right_handed_prob = P_A_given_rh(ages, death_distribution_data, 1990) #for age in ages ]
6
7  plt.plot(ages, left_handed_prob, label = "Left Handed")
8  plt.plot(ages, right_handed_prob, label = "Right Handed")
9  plt.title("Left vs Right Handed Prob", color = 'Red', size = 15)
10 plt.xlabel("Ages", size=15, color='green')
11 plt.ylabel("Probabilities at Age of Death", size= 15, color= 'green')
12 plt.legend()
13
14 plt.show()
```

...



Data Analysis Phase - 4 : Analyze

Task 9: Instructions

Find the mean age at death for left-handers and right-handers.

- Multiply the `ages` list by the left-handed probabilities of being those ages at death, then use `np.nansum` to calculate the sum. Assign the result to `average_lh_age`.
- Do the same with the right-handed probabilities to calculate `average_rh_age`.
- Print `average_lh_age` and `average_rh_age`.
- Calculate the difference between the two average ages and print it.

To make your printed output prettier, try using the `round()` function to round your results to two decimal places.

Task 9:

```
1 Avg_lh_age = np.nansum(ages*np.array(left_handed_prob))
2 Avg_rh_age = np.nansum(ages*np.array(right_handed_prob))
3
4 print(round(Avg_lh_age,2))
5
6 print(round(Avg_rh_age,2))
7
8 print("The difference between the two average ages is : "+ str(round(Avg_rh_age-Avg_lh_age,2))+ "Years")
```

67.26
72.8
The difference between the two average ages is : 5.54Years

Data Analysis Phase - 4 : Analyze

Task 10: Instructions

Redo the calculation from Task 8, setting the `study_year` parameter to 2018.

- In the call to `P_A_given_lh`, set `age_of_death` to `ages`, `death_distribution_data` to `death_distribution_data`, and `study_year` to 2018.
- Do the same for `P_A_given_rh`.

Task 10:



```
1 #Given study Year = 2018
2
3 left_hand_prob_2018 = P_A_given_lh(ages, death_distribution_data, study_year=2018)
4
5 right_hand_prob_2018 = P_A_given_rh(ages, death_distribution_data, study_year=2018)
6
7 Avg_lh_age_2018 = np.nansum(ages*np.array(left_hand_prob_2018))
8
9 Avg_rh_age_2018 = np.nansum(ages*np.array(right_hand_prob_2018))
```

```
1 print("The difference between the two average ages is : "+ str(round(Avg_rh_age_2018-Avg_lh_age_2018,2))+"Years"+"\\n"+"if the study were to take place on 2018")
```

The difference between the two average ages is : 2.34Years
if the study were to take place on 2018

Data Analysis Phase – 5 : Share

Findings on the Claim:

Original Claim (1991): Suggested a 9-year difference in age at death between left and right handers.

Current Study : finding the mortality rate differences are not statistically significant.

Data Misled: The percentage of left-handers was higher in younger populations than older ones. Therefore, a snapshot study of dead people will make it look like left-handers die younger, simply because there are fewer old left-handers in the general population.

Age Gap :

From the Data,

Older People (80-90 Years old) Appear mostly Right Handed

Younger population appear to be recorded as left-handed correctly

So, it looked like Left Hander's Die Young, Right Handers live longer

But Actually,

Older Left handers are just being misclassified as right handers.

Data Analysis Phase – 6 : Act

Results & Conclusions



Data Analysis Phase – 6 : Act

Results & Conclusions :

People around 1900 only 3% were left handed, Today/Recent study indicates it is around 11%.

As per the study conducted in 1990

the difference between average age of Right Handers and Left Handers from the given data is found to $[72.8 - 67.26 = 5.54 \text{ Years}]$

If the study were to be conducted in the year 2018, the difference between average age of Right Handers and Left Handers from the given data is found to 2.34 Years

From the above analysis there is no statistical evidence that the age gap between the Right Handers & Left Handers is 9 Year at the time of death of a person given he/she is left handed.


From the Present Study it can be said that The death of a subject at a particular age is 9 years early given that he is left handed compared to that of right handed is a **complete Myth** rather than a Reality

Recommendations


Exact Reporting of the left-handedness data at all ages / generations without any data bias / cohort bias & Sampling Bias can provide accurate result of future studies.

Recommendations for Future Study

```
1 plt.plot(lefthanded_data.Birth_year, lefthanded_data.Mean_lh, marker='o', label='Mean')
2 plt.title('Birth Year vs Mean_lh', size=20, color = 'Red')
3 plt.xlabel("Birth Year", color='green', size = 15)
4 plt.ylabel("Mean Age", color= 'green', size = 15)
5 plt.legend()
6 plt.axis()
7 plt.show()
```



```
1 plt.plot(df4.Age, df4.Male, label='Male')
2 plt.plot(df4.Age, df4.Female, label='Female')
3 plt.legend()
4 plt.axis()
5 plt.show()
```



When using age-based left handedness data, results suggest left-hander's die younger. However, after adjusting for cohort effects using Birth-Year based Handedness, the effects disappear indicating the original finding was due to confounding rather than a real difference in lifespan – Referring to Fig -A

The data indicates that there might be slight **correlation** between the death of subject at a given age but it is definitely not the **cause** that he/she is Left handed – Referring to Fig -A

Referring to Fig -B

The graph plotted against **age vs death distribution data** indicates that data is **negatively skewed** not perfectly normal distribution, which can indicate a sampling bias

Workbook/ Notebook Generated from Google Colab Environment

Project Github Link :

[https://github.com/M339KUMAR/ColabDemoFiles/blob/78482ca811fd319f162cef76ab319eecda5aea/MTE%20DA/MTE_DA_Apr2026\(Apr20th-1\).ipynb](https://github.com/M339KUMAR/ColabDemoFiles/blob/78482ca811fd319f162cef76ab319eecda5aea/MTE%20DA/MTE_DA_Apr2026(Apr20th-1).ipynb)

[ColabDemoFiles](#) / [MTE DA](#) / 

 **M339KUMAR** Add files via upload

Name
 ..
 1
 1.png
 2.png
 3.png
 4.png
 5.png
 6.png
 MTE_DA_Apr2026(Apr19th-1).ipynb
 MTE_DA_Apr2026(Apr20th-1).ipynb

Thank You

I Express my Sincere Gratitude to **MedTourEasy** for giving the opportunity to be part of their Team and work on the Healthcare Data Analytic Project,

I Express my Thanks to **Mr. Ankit Hasija**, Training Head, MTE
For guiding me in this project, and providing the necessary support to complete this project as per the schedule,

Thanking you,
PRAVEENKUMAR MOPURU
mpk339@gmail.com
9985772150