

FREDHOLM OPERATORS OVER HILBERT C^* -MODULES

Mezzovilla

November 23, 2022

Contents

1	Introduction	5
2	K-theory of Banach Algebras	7
2.1	General portrait of homological theories	7
2.2	The K_0 group	9
2.3	The K_1 group	13
2.4	The index map	14
3	Hilbert C^*-modules	17
3.1	The interest object	17
3.2	Adjointable operators	21
3.3	Compact and Finite rank operators	24
3.4	Kasparov Stabilization Theorem	29
3.5	Rank definition of Finite Rank Hilbert Modules	31
3.6	Quasi-stably-isomorphic Hilbert modules	36
4	Fredholm Operators	41
4.1	Regular Fredholm operators	44
4.2	Regularization of Fredholm operators	48
4.3	Fredholm Picture of $K_0(A)$	54
5	Application to Morita Equivalence	57
5.1	Preliminars on Hilbert C^* -bimodules	57
5.2	K -theory and Hilbert C^* -bimodules	58

Chapter 1

Introduction

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Chapter 2

K-theory of Banach Algebras

2.1 General portrait of homological theories

A homological theory for a category \mathcal{C} consist in a sequence of covariant functors $H_n : \mathcal{C} \longrightarrow \mathbf{GrpAb}$ for each $n \in \mathbb{N}$ which satisfies some set of axioms, which depends on what theory one is interested. For example, if \mathcal{C} contains a nice homotopical concept, its rather common to ask for homotopical invariance. If exact sequences naturally pops in the domain encoding a lot of information, some other axioms are required to obtain long exact sequences. The usual notation is:

$$\begin{array}{ccc} H_n : \mathcal{C} & \longrightarrow & \mathbf{GrpAb} \\ A & \longmapsto & H_n(A) \\ \phi \downarrow & & \phi_n \downarrow \\ B & \longmapsto & H_n(B) \end{array}$$

We also need a way to translate short exact sequences of the form

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$


from the original category to higher counterparts obtained by H_n , hence, every homology theory seeks to define a connecting morphism $\delta_n : H_n(C) \longrightarrow H_{n+1}(A)$ into a long exact sequence:

$$\begin{array}{ccccccc} H_0(A) & \longrightarrow & H_0(B) & \longrightarrow & H_0(C) & \longrightarrow & \\ \downarrow & & \delta_0 & & \downarrow & & \\ H_1(A) & \longrightarrow & H_1(B) & \longrightarrow & H_1(C) & \longrightarrow & \\ \downarrow & & \delta_1 & & \downarrow & & \\ H_2(A) & \longrightarrow & H_2(B) & \longrightarrow & H_2(C) & \longrightarrow & \dots \end{array}$$

In the other hand, as everything containing the prefix “co”, cohomology theories are consisted of contra-variant functors $(H^n)_n$ with the same pay-off. The position of the index on the notation usually indicates what sort of theory one is dealing with.

Here we are concerned with a homology theory for complex Banach Algebras $\mathcal{B}\text{-Alg}$ or, more popularly, for C^* -algebras $C^*\text{-Alg}$, a.k.a., K -theory for Operator Algebras. It is the mirror image of Topological K -theory, in light of *Gelfand Duality* connecting the category of Locally Compact Hausdorff spaces and complex abelian C^* -algebras, but not restricted to commutative spaces, which is often referred to as the "Non-Commutative Topology".

In his work to reformulate Riemman-Roch theorem [2], *A. Grothendieck* introduces the group $K(A)$ associated to a subcategory of an abelian category, which nowadays, it is the so called “Grothendieck’s group”. That’s where is from the letter K , which he had chosen for “Klassen”. His reformulation famously contains his legendary drawing:



Riemann-Roch Theorem: the new black: The diagram

[...] is commutative!

I would need to misuse about 2h of my listener’s time in order to impart only an approximal understanding of this statement for $f : X \longrightarrow Y$.

In cold print (as in Springer’s lecture notes) this would take around 400-500 pages.

A thrilling example of how our urge for knowledge and discoveries decays into a lifeless and ideological delirium while life itself goes thousandfold to the devil - and is threatened with final destruction.

It’s high time to change our course!

(16.12.1971)

Alexander Grothendieck

With his work settled in what is about to be algebraic K -theory, topological K -theory would be a product of *M. Atiyah* and *F. Hirzebruch* replicating Grothendieck’s construction for topological vector bundles over compact Hausdorff spaces.

We'll construct functors $K_n : \mathcal{B}\text{-}\mathbf{Alg} \longrightarrow \mathbf{GrpAb}$ and the connecting maps will be call *index map*, denoted by ∂ . A remarkable aspect of operator K -theory is the *Bott periodicity*: $K_n \simeq K_{n+2}$, which then describes for any short exact sequence $0 \longrightarrow I \longrightarrow A \longrightarrow A/I \longrightarrow 0$, where $I \triangleleft A$, a six-term exact sequence:

$$\begin{array}{ccccc} K_0(I) & \longrightarrow & K_0(A) & \longrightarrow & K_0(A/I) \\ \partial \uparrow & & & & \downarrow \partial \\ K_1(A/I) & \longleftarrow & K_1(A) & \longleftarrow & K_1(I) \end{array}$$

The details will be spared in what is outside of our scope, which will include the definition of the groups K_0 and K_1 for complex Banach algebras, and the index map mentioned. The connecting map will be used in the classification of finite rank modules, and later on, the index of our Fredholm operators. Hence it is important to define it in a helpfull way.

2.2 The K_0 -group

Our object is to deal with Hilbert C^* -modules, witch are right A -modules with a generalized A -valued inner product for a given C^* -algebra A (plus some other details), generalizing the concept of Hilbert space. Therefore, it is reasonable to understand some K -theory for C^* -algebras, as they are our underlying space. Unfortunately, as we'll see later on, there is no *Riesz representation lemma* and, there exists bounded linear operators that aren't adjointable between Hilbert modules. Hence, dealing with self-adjoint operators is a restriction for sure. Thankfully, the K -theory for Banach algebras is good enough in order to fill our needs.

In topological K -theory, in order to define the 0th K group, one would consider a complex vector bundle E over $X \in \mathbf{CHaus}$ and take the right $C(X)$ -module $\Gamma(X, E)$ of continuous sections $s : X \longrightarrow E$ with pointwise scalar multiplication¹. Compactness of X implies that $\Gamma(X, E)$ is a projective $C(X)$ -module, and *Serre-Swan* theorem [10, Thr. 6.18] states that $E \longmapsto \Gamma(X, E)$ induces an equivalence between the category of complex vector bundles and finitely generated projective $C(X)$ -modules. Hence, $K^0(X)$ is the *Grothendieck group* of the set of equivalence classes of isomorphisms between vector bundles over X .

For a given Banach Algebra A , the following definitions and constructions mimics the above paragraph, by replacing vector bundles by finitely generated projective A -modules.

Definition 2.2.1. In any given Banach algebra A , for two idempotent elements x and y , define the following notions of equivalence:


¹That is to say, for $f \in C(X)$ and $s \in \Gamma(X, E)$, let $x \longmapsto s(x)f(x)$.

- (i) **Murray-von-Neuman equivalent:** There are elements $p, q \in A$ such that $x = pq$ and $y = qp$.
- (ii) **Similarly equivalent:** There exists an invertible² element $u \in \text{GL}(A)$ such that $x = u^{-1}yu$.
- (iii) **Homotopic:** There are a continuous path $\gamma \in C([0, 1], A)$ of idempotents between x and y , i.e.,

$$\gamma(0) = x, \gamma(1) = y \quad \text{and} \quad \forall t \in [0, 1], \gamma(t)^2 = \gamma(t).$$

If A is assured to be a C^* -algebra, those definitions are concerned with self-adjoint idempotent elements, a.k.a., projections. Two projections x, y are equivalent if there exists a *partial isometry* u such that $x = u^*u$ and $y = uu^*$.

For the canonical embedding $x \mapsto \text{diag}(x, 0)$ over matrices, consider the inductive limit $\mathbb{M}_\infty(A) := \varinjlim_{n \in \mathbb{N}} \mathbb{M}_n(A)$, which can be seen as the set of infinite matrices over A but only finitely many of the entries are non-zero.

Remark 2.2.2. Note that $\mathbb{M}_\infty(A)$ contains no unity, but that doesn't stop us to declaring two elements x, y to be similar when they are similar in some square matrix space $\mathbb{M}_n(A)$. Therefore, all equivalence relations listed in the definition 2.2.1 coincide in $\mathbb{M}_\infty(A)$. 

Simply shouting “Let A be an C^* -algebra” in the crowd is a powerful classification tool, whenever is a mathematicians crowd³.

- (i) If you hear in response “unital or not?”, you know that there is some C^* -algebraic fellow around you.
- (ii) If the crowd contains mathematicians and no-one ask whether A contains a unity or not, no C^* -algebraist is contained in the crowd. They are instantly assuming the unity is there.

This is because dealing without unital rings outside C^* -theories are usually simple. Just unitize and go on. However, the presence of unity in C^* -algebras is crucial to determine their underlying hidden topology, as explicitly is made in *Gelfand's duality* theorem.

The next definition is in charge to define the functor K_0 for both cases, but some intermediate steps are required from one to another.

Definition 2.2.3. Let A be a Banach algebra. The set of equivalence classes over $\mathbb{M}_\infty(A)$ considering any relation \sim contained in 2.2.1 is an abelian semi-group with $[x] + [y] := [\text{diag}(x, y)]$. Before defining K_0 , in order to include

²Assuming that A is unital.

³Otherwise, you are just playing creepy at dinner table again.

the non necessarily unital algebras, it is needed to be considered an auxiliar functor K_{00} much closer to the topological counterpart K^0 . This is necessary in order to obtain the Bott periodicity result for Banach algebras, and other good functorial properties.

- (i) **K_{00}** : It is the Grothendieck group construction associated with the semi-group $V(A) := \mathbb{M}_\infty(A) / \sim$ where addition is given by $[x] + [y] := [\text{diag}(x, y)]$, generalising the construction of \mathbb{Z} from \mathbb{N} considering formal differences. In lighter sheets, for pairs (a, b) and (c, d) of elements in $V(A)$, let $(a, b) \equiv (c, d)$ whenever there exists⁴ $z \in V(A)$ such that $a + d + z = c + b + z$. This is an equivalence relation over the pairs, and $[\cdot]_{00}$ will denote the related equivalence class.

We are mimicking the formal differences construction, so it's natural to define the addition operation coordinate-wise and let $x - y := [(x, y)]_{00}$. Therefore, it is well defined the following covariant functor:

$$\begin{array}{ccccc}
 K_{00}: & \mathcal{B}\text{-Alg} & \longrightarrow & \mathbf{GrpAb} & \\
 & A \longmapsto V(A) \times V(A) / \equiv & & x - y & \\
 & \phi \downarrow & & \phi_{00} \downarrow & \downarrow \\
 & B \longmapsto V(B) \times V(B) / \equiv & & \phi(x) - \phi(y) &
 \end{array}$$

Since every element in $V(A)$ is the class of some idempotent matrix p , we can state that every element in $K_{00}(A)$ is on the form $[p]_{00} - [q]_{00}$. Two formal differences $[p]_{00} - [q]_{00}$ and $[x]_{00} - [y]_{00}$ coincide in $K_{00}(A)$ precisely when the operators $\text{diag}(p, y)$ and $\text{diag}(x, q)$ are *stably* homotopic.

- (ii) **K_0** : In our next step, it's crucial to know exactly who $K_{00}(\mathbb{C})$ is. Hence, remeber that two idempotents in $\mathbb{M}_n(\mathbb{C})$ are similar if, and only if, their images has the same dimension. Therefore $V(\mathbb{C}) \simeq \mathbb{N}$, and by historical nightmares with Analysis I exercises constructing the integer numbers, it is easy to infer that $K_{00}(\mathbb{C}) = \mathbb{Z}$.

For non necessarily unital A , consider $\tilde{A} := A \oplus \mathbb{C}$ the *unifization* of A and the complex projection $\varepsilon : \tilde{A} \twoheadrightarrow \mathbb{C}$, which induces the short exact sequence:

$$0 \longrightarrow A \hookrightarrow \tilde{A} \xrightarrow{\varepsilon} \mathbb{C} \longrightarrow 0$$

The urge to obtain Bott periodicity theorem for Banach algebras, which is a relation between K_0 and K_1 in the presence of short exact

⁴Since it is only a semi-group, the cancelation property do not hold necessarily over $V(A)$. One might check that this is the case if, and only if, the inclusion of $V(A)$ at the Grothendieck's associated group is injective.

sequences, will oblige the exactness of the following:

$$0 \longrightarrow K_0(A) \hookrightarrow K_0(\tilde{A}) \xrightarrow{\varepsilon_0} K_0(\mathbb{C}) \longrightarrow 0$$

Since it is a morphism between unital Banach algebras, the induced map $\varepsilon_0 : K_0(\tilde{A}) \rightarrow \mathbb{Z}$ is a well defined morphism, hence, it is possible to define the following:

$$\begin{array}{ccccc} K_0: & \mathcal{B}\text{-Alg} & \longrightarrow & \mathbf{GrpAb} & \\ & A \longmapsto \ker(K_0(\tilde{A}) \rightarrow \mathbb{Z}) & & a + z & \\ & \phi \downarrow & & \phi_0 \downarrow & \downarrow \\ & B \longmapsto \ker(K_0(\tilde{B}) \rightarrow \mathbb{Z}) & & \phi(a) + z & \end{array}$$

Notice that $K_0(A)$ is precisely the set of elements $[p]_0 - [q]_0 \in K_0(\tilde{A})$ such that $\varepsilon(p) \sim \varepsilon(q)$. If A is already unital, it is possible to show that $K_0(A) \simeq K_{00}(A)$.

Remark 2.2.4. The argument to show that $V(\mathbb{C}) \simeq \mathbb{N}$ is equivalent for compact operators in an infinite-dimensional Hilbert space H , i.e., $V(\mathcal{K}(H)) \simeq \mathbb{N}$, hence $K_0\mathcal{K}(H) = \mathbb{Z}$. On the other hand, any two infinite rank projections in $\mathcal{B}(H)$ are equivalent, hence $V\mathcal{B}(H) \simeq \mathbb{N} \cup \{\infty\}$, which is a semi-group without the cancellation property. Since everyone is equivalent to ∞ , it is obtained that $K_{00}\mathcal{B}(H) \simeq 0$. The semi-group $V(A)$ possesses the cancellation property if, and only if, the inclusion $V(A) \hookrightarrow K_{00}(A)$ is injective. \blacksquare

Proposition 2.2.5 (Standard portrait of K_0). Every element of $K_0(A)$ can be written as $[x + p_n]_0 - [p_n]_0$.

Proof. Let $p, q \in \mathbb{M}_\infty(\tilde{A})$ be some idempotent square matrices with order no longer than n , such that $\varepsilon_0([p]_0 - [q]_0) = 0$, i.e., $[p]_0 - [q]_0 \in K_0(A)$. Matrices $p \in \mathbb{M}_n(\tilde{A})$ can be written as $(p_A, p_C) \in \mathbb{M}_n(A) \oplus \mathbb{M}_n(\mathbb{C})$, i.e., an algebraic part p_A and a scalar part p_C . Stating that $\varepsilon_0([p]_0 - [q]_0) = 0$ means that the scalar parts of p and q coincide.

The identity $I_n \in \mathbb{M}_\infty(\tilde{A})$ can be seen as the projection operator of the first n -th coordinates, by filling it with 0's, but to avoid confusions, let it be p_n . With $y \leq x$ be given by $xy = yx = y$, one may see that $p < p_n$ and $y < p_n$. Notice that $\text{diag}(0, p) \in \mathbb{M}_{2n}(\tilde{A})$ is similar to p and orthogonal to I_n , i.e.,

$$\begin{pmatrix} 0 & 0 \\ 0 & p \end{pmatrix} \begin{pmatrix} I_n & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} I_n & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & p \end{pmatrix} = 0.$$

Hence, $x := \text{diag}(-q, p)$ is such that $x + p_n$ is an idempotent operator and:

$$\begin{aligned} [x + p_n]_0 - [p_n]_0 &= [\text{diag}(0, p)]_0 + [p_n - q]_0 - ([p_n - q]_0 + [q]_0) \\ &= [p]_0 - [q]_0. \end{aligned} \quad \square$$

2.3 The K_1 -group

While K_0 is build upon equivalence classes of idempotent, K_1 uses invertible elements, but simpler. Therefore, let $\mathrm{GL}_\infty(A) := \varinjlim_{n \in \mathbb{N}} \mathrm{GL}_n(A)$ considering the embedding $x \mapsto \mathrm{diag}(x, 1)$. Calculus is back, and we shall consider exponentials inside a unital algebra A :

$$\exp(a) := \sum_{n=0}^{\infty} \frac{a^n}{n!} \quad \text{and} \quad \log(1+a) := \sum_{n=1}^{\infty} -\frac{a^n}{n} \quad (a \in A)$$

where the log is defined whenever $\|a\| < 1$. This is the case since elements of the form $z - a$ for complex z are invertible if $\|a\| \leq |z|$. If a and b doesn't commute, $\exp(a)\exp(b) \neq \exp(a+b)$, which means that the set of exponentials isn't closed by multiplications.

Lemma 2.3.1. For a unital Banach algebra A , the component of the unity is the group generated by $\{\exp(a) \mid a \in A\} \subset \mathrm{GL}(A)$, denoted by $\exp(A)$.

Proof. Let $\mathrm{GL}^{(0)}(A)$ be the referred set of connected components of 1. Notice that $t \mapsto \exp(tb)$ for $t \in [0, 1]$ is a continuous path of invertible elements between 1 and $\exp(b)$ for any b , hence $\exp(A) \subset \mathrm{GL}^{(0)}(A)$. It remains only to show the converse inclusion.

For some a with $\|1 - a\| < 1$, let $b := \log(1 + (a - 1)) = \log(a)$, i.e., $a = \exp(b)$. Therefore, if $u \in \mathrm{GL}(A)$ and $\|v - u\| < \|u^{-1}\|^{-1}$, this means that $v = \exp(b)u$ for some b . From this treatment, it follows that $\exp(A)$ is a open and closed topological subspace of $\mathrm{GL}^{(0)}(A)$ which contains the unity, i.e., $\mathrm{GL}^{(0)}(A)$ coincides with $\exp(A)$. \square

Remark 2.3.2. Let $M \in \mathrm{GL}_n(\mathbb{C})$. Since 0 cannot be an eigenvalue of M (which is a finite set), it's possible to find $\alpha \neq 0$ such that $[0, \infty) \cdot \alpha$ doesn't contains any of the eigenvalues of M or 1. Therefore, $1 - \alpha t \neq 0$ for all $t \geq 0$ and $M_t := (1 - \alpha t)^{-1}(M - t\alpha I_n)$ is a continuous path from M to the identity, i.e., $\mathrm{GL}_n(\mathbb{C})$ is connected. \blacksquare

In a not so long future, the following result will be important in the presence of an ideal $I \triangleleft A$, the considering of the projection $A \twoheadrightarrow A/I$.

Corollary 2.3.3. Any continuous surjection $A \twoheadrightarrow B$ induces a lift from every element in $\mathrm{GL}_n^{(0)}(B)$ to one in $\mathrm{GL}_n^{(0)}(A)$.

Proof. Using 2.3.1, write $\prod_i \exp(b_i) \in \mathrm{GL}_n(B)^{(0)}$ for any desired element. Since there is a surjection, there exists lifts to each b_i , i.e., $a_i \in \mathrm{GL}_n(A)$ such that $\prod_i \exp(a_i) \in \mathrm{GL}_n(A)_0$. \square

Considering the homotopy equivalence relation, two elements in $\mathrm{GL}_\infty(A)$ are homotopical whenever they are in the same connected component in some $\mathrm{GL}_n(A)$. Denote the equivalence class by $[\cdot]_1$. Whence, the quotient


$\mathrm{GL}_\infty(A)/\mathrm{GL}_\infty^{(0)}(A)$ is an abelian group with the multiplication $[x]_1[y]_1 = [xy]_1$, which is commutative once you note it is possible to find a connected path⁵ between $\mathrm{diag}(y, 1)$ and $\mathrm{diag}(1, y)$, hence

$$[x]_1[y]_1 = [xy]_1 = \left[\begin{pmatrix} xy & 0 \\ 0 & 1 \end{pmatrix} \right]_1 = \left[\begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \right]_1 = \left[\begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix} \right]_1$$

and similarly, one shows that $[xy]_1 = [\mathrm{diag}(y, x)]_1 = [y]_1[x]_1$. There we have, our $K_1(A)$ group. Since $\mathrm{GL}_n(\mathbb{C})$ is connected, it follows immediately that $K_1(\mathbb{C}) = 0$ and, therefore, we can deal with units the way it is intended: for non necessarily unital algebras A , let $K_1(A) := K_1(\tilde{A})$.

Definition 2.3.4. The functor K_1 can be seen as the following:

$$\begin{array}{ccccc} K_1: & \mathcal{B}\text{-Alg} & \longrightarrow & \mathbf{GrpAb} & \\ & A & \longmapsto & \mathrm{GL}_\infty(\tilde{A})/\mathrm{GL}_\infty^{(0)}(\tilde{A}) & [x]_1 \\ & \phi \downarrow & & \phi_1 \downarrow & \downarrow \\ & B & \longmapsto & \mathrm{GL}_\infty(\tilde{B})/\mathrm{GL}_\infty^{(0)}(\tilde{B}) & [\phi(x)]_1 \end{array}$$

Remark 2.3.5. It should be stated that if one is dealing with a C^* -algebra, then K_1 can be obtained by the set of unitary matrices $U_n(A)$, i.e., $u^* = u^{-1}$; Since $U_n(A)/U_n^{(0)}(A) \cong \mathrm{GL}_n(A)/\mathrm{GL}_n^{(0)}(A)$, one can obtain a deformation retraction from $\mathrm{GL}_n(A)$ to $U_n(A)$ by the polar decomposition, hence, $K_1(A)$ is isomorphic to $U_\infty(A)/U_\infty^{(0)}(A)$. 

2.4 The index map

We are now ready to define the so called index map. This name comes from the Fredholm operator theory since what we are about to construct is a generalization of the index of those operators. Consider $\mathcal{B}(H)$ the C^* -algebra of bounded operators between a Hilbert space H , and $\mathcal{K}(H)$ the ideal of compact operators. The *Atkinson* theorem states precisely that the *Calkin* algebra $\mathcal{Q}(H) := \mathcal{B}(H)/\mathcal{K}(H)$ is a classifying one: T is a Fredholm operator if, and only if, $(T \bmod \mathcal{K}(H)) \in \mathrm{GL} \mathcal{Q}(H)$.

Since $K_0\mathcal{K}(H) = \mathbb{Z}$ and $K_1\mathcal{Q}(H)$ can be seen the set of Fredholm operators up to homotopy⁶, the index map $\mathbf{ind} : K_1\mathcal{Q}(H) \longrightarrow K_0\mathcal{K}(H)$ is well defined. Our index map ∂ will generalize this application.


⁵Let $z(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ be the rotation matrix by some angle θ . Therefore, the continuous map $[0, \pi/2] \ni \theta \longrightarrow z(\theta) \mathrm{diag}(y, 1) z(\theta)^{-1}$ is the desired path.

⁶Remind that two Fredholm operators in the same realm have the same index if, and only if they are homotopic.

Construction 2.4.1. Let $I \triangleleft A$ and consider the following short exact sequence:

$$0 \longrightarrow I \hookrightarrow A \twoheadrightarrow A/I \longrightarrow 0$$

We are in position to construct $\partial : K_1(A/I) \longrightarrow K_0(I)$. For $[x]_1 \in K_1(A/I)$, let n be such that $x \in \mathrm{GL}_n(\widetilde{A/I})$. It's about time to the corollary 2.3.3 to shine: Since the projection $A \twoheadrightarrow A/I$ is a continuous surjection, so it is the unification induced morphism between the algebras, hence, one can lift the element $\mathrm{diag}(x, x^{-1}) \in \mathrm{GL}_{2n}^{(0)}(\widetilde{A/I})$ to some $w \in \mathrm{GL}_{2n}^{(0)}(\widetilde{A})$.

If $\pi : \mathrm{GL}_\infty(\widetilde{A}) \twoheadrightarrow \mathrm{GL}_\infty(\widetilde{A/I})$ is the quotient projection, notice that $\pi(wp_n w^{-1}) = p_n$, so that $wp_n w^{-1} \in \widetilde{I}$. Since $wp_n w^{-1}$ is also an idempotent, notice that $[wp_n w^{-1}]_0 - [p_n]_0 \in K_0(I)$. And this is the image of the index map ∂ of some element $[x]_1$. 

An anxious mind would immediately panic. We have a TO-DO list before calling it a day:

- (i) Check that $[wp_n w^{-1}]_0 - [p_n]_0$ doesn't depend on the lift w chosen;
- (ii) Check that $\partial([x]_1) = \partial([y]_1)$ for $x \equiv y$.
- (iii) Check that ∂ is a group morphism.

Proof of TO-DO list items. If v is another lift of $\mathrm{diag}(x, x^{-1})$, notice that

$$vp_n v^{-1} = (vw^{-1})wp_n w^{-1}(vw^{-1})^{-1},$$

i.e., $vp_n v^{-1}$ is similar to $wp_n w^{-1}$. This is enough to take care of (i).

In order to show that the index is well defined, suppose that $y \in \mathrm{GL}_n(\widetilde{A/I})$ is another representant of class $[x]_1$. Notice that

$$x^{-1}y \in \mathrm{GL}_n^{(0)}(\widetilde{A/I}) \quad \text{and} \quad \begin{pmatrix} x & 0 \\ 0 & I_n \end{pmatrix} \begin{pmatrix} I_n & 0 \\ 0 & y^{-1} \end{pmatrix} \in \mathrm{GL}_{2n}^{(0)}(\widetilde{A/I})$$

so by the corollary 2.3.3 again, let $a \in \mathrm{GL}_n^{(0)}(\widetilde{A})$ and $b \in \mathrm{GL}_{2n}^{(0)}(\widetilde{A})$ be the lifts respectively. But then $u := w \mathrm{diag}(a, b)$ is a lift of $\mathrm{diag}(y, y^{-1})$. From the fact that p_n commutes with $\mathrm{diag}(a, b)$, it is obtained that $up_n u^{-1} = wp_n w^{-1}$. Since we already show that the choice of lift doesn't matter, (ii) is checked.

For $x, y \in \mathrm{GL}_n(\widetilde{A/I})$, suppose that w is a lift of $\mathrm{diag}(x, x^{-1})$ and v is a lift of $\mathrm{diag}(y, y^{-1})$. Notice that $\varpi := \mathrm{diag}(w, v)$ is a lift of $\mathrm{diag}(x, y, x^{-1}, y^{-1})$,

hence

$$\begin{aligned}
\partial([x]_1[y]_1) &= [\varpi p_{2n} \varpi^{-1}]_0 - [p_{2n}]_0 \\
&= \left[\begin{pmatrix} w & 0 \\ 0 & v \end{pmatrix} \begin{pmatrix} p_n & 0 \\ 0 & p_n \end{pmatrix} \begin{pmatrix} w & 0 \\ 0 & v \end{pmatrix}^{-1} \right]_0 - \left[\begin{pmatrix} p_n & 0 \\ 0 & p_n \end{pmatrix} \right]_0 \\
&= \left[\begin{pmatrix} wp_n w^{-1} & 0 \\ 0 & vp_n v^{-1} \end{pmatrix} \right]_0 - \left[\begin{pmatrix} p_n & 0 \\ 0 & p_n \end{pmatrix} \right]_0 \\
&= [wp_n w^{-1}]_0 - [p_n]_0 + [vp_n v^{-1}]_0 - [p_n]_0 = \partial[x]_1 + \partial[y]_1
\end{aligned}$$

Therefore, it is a group morphism as our final item (iii) assures. \square

Definition 2.4.2 (Index map in *K*-theory). Using construction 2.4.1, the group morphism so called *index* map is given by

$$\begin{aligned}
\partial : K_1(A/I) &\longrightarrow K_0(I) \\
[x]_1 &\longmapsto [wp_n w^{-1}]_0 - [p_n]_0
\end{aligned}$$


whenever $x \in \text{GL}_n(\widetilde{A/I})$ and $w \in \text{GL}_{2n}^{(0)}(\widetilde{A})$ is a lift of $\text{diag}(x, x^{-1})$.

Example 2.4.3. In a unital *C**-algebra *A*, if a unitary⁷ idempotent element *u* in $\text{GL}_n(A/I)$ lifts to $v \in \mathbb{M}_n(A)$, the element

$$w := \begin{pmatrix} v & I_n - v^*v \\ I_n - vv^* & v^* \end{pmatrix}$$

is a lift for $\text{diag}(u, u^{-1})$. Therefore:

$$\begin{aligned}
\partial[u]_1 &= [wp_n w^{-1}]_0 - [p_n]_0 \\
&= \left[\begin{pmatrix} v & I_n - v^*v \\ I_n - vv^* & v^* \end{pmatrix} \begin{pmatrix} I_n & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} v^* & I_n - vv^* \\ I_n - vv^* & v \end{pmatrix} \right]_0 - \left[\begin{pmatrix} I_n & 0 \\ 0 & 0 \end{pmatrix} \right]_0 \\
&= \left[\begin{pmatrix} vv^* & 0 \\ 0 & I_n - vv^* \end{pmatrix} \right]_0 - \left[\begin{pmatrix} I_n & 0 \\ 0 & 0 \end{pmatrix} \right]_0 \\
&= [I_n - v^*v]_0 - [I_n - vv^*]_0
\end{aligned}$$

Notice that when $A = \mathcal{B}(H)$ and $I = \mathcal{K}(H)$, we dealing again with Fredholm operators living in the Calking algebra $\mathcal{Q}(H)$ and ∂ coincides with the Fredholm index. 

⁷I.e., $u^* = u^{-1}$.

Chapter 3

Hilbert C^* -modules

Hilbert modules first appear in the work of *I. Kaplaski* [9] and *W. Paschke* [15] later. There are three main areas where Hilbert C^* -modules are heavily used to formulate mathematical concepts involving:

- (i) Induced representations of Morita equivalence [3], [19], [18];
- (ii) Kasparov's KK -theory [11];
- (iii) C^* -algebraic quantum groups.

In what is tangible to this work, we address the Morita equivalence target by building a Fredholm operator approach between Hilbert modules, introduced by Ruy Exel [6]. Hence, this chapter is responsible for defining and studying those objects.

The material source contains for this chapter contains the well written textbooks like [12], [8], [13].

3.1 The interest object

Definition 3.1.1 (Inner product Module). A right module E over a C^* -algebra (non-necessarily unital) blessed with an generalized inner product $\langle \cdot, \cdot \rangle : E \times E \longrightarrow A$ will be said to be a *Inner product module* when $\langle \cdot, \cdot \rangle$ attends the following properties:

- (i) **Twisted A -sesquilinear**: The first coordinate are involuted-linear and the second one linear, i.e.,

$$\begin{cases} \langle x + ya, z \rangle = \langle x, z \rangle + a^* \langle y, z \rangle \\ \langle z, x + ya \rangle = \langle z, x \rangle + \langle z, y \rangle a \end{cases} \quad \left(\begin{array}{l} x, y, z \in E \\ a \in A \end{array} \right)$$

- (ii) **A -Hermitian symmetry**: $\langle x, y \rangle = \langle y, x \rangle^*$ whenever $x, y \in E$.

(iii) **Positive definite:** For any $x \in E$, $\langle x, x \rangle = 0 \Leftrightarrow x = 0$. By (ii), we can say that $\langle x, x \rangle \geq 0$ since it is self-adjoint.

One could argue that we only need the inner product to be linear in the second coordinate and by the Hermitian symmetry conclude as a proposition that every inner product over Inner product modules is indeed twisted sesquilinear.

Proposition 3.1.2 (Cauchy-Schwartz inequality). For any Inner product module E over A , the following inequality holds:

$$(3.1) \quad \|\langle x, y \rangle\|^2 \leq \|\langle x, x \rangle\| \cdot \|\langle y, y \rangle\|. \quad (x, y \in E)$$

Proof. Given the fact that $0 \leq \langle a, a \rangle$ for $a \in A$, notice that with the accessory elements $a := \langle x, x \rangle$, $b := \langle y, y \rangle$ and $c := \langle x, y \rangle$,

$$\begin{aligned} 0 &\leq \langle x - ytc^*, x - ytc^* \rangle \\ &= \langle x, x - ytc^* \rangle - tc\langle y, x - ytc^* \rangle \\ &= \langle x, x \rangle - \langle x, y \rangle tc^* - tc\langle y, x \rangle + tc\langle y, y \rangle tc^* \\ &= a - 2tcc^* + t^2cbc^* \end{aligned} \quad (t \in \mathbb{R})$$

Since $2tcc^*$ is self-adjoint, we can add in both sides and maintain the inequality in the C^* -realm. Using the A -norm and assumig $t \geq 0$, by ??,

$$\begin{aligned} 2t\|cc^*\| &\leq \|a\| + t^2\|cbc^*\| \\ &\leq \|a\| + t^2\|c\|\|b\|\|c^*\| \\ (3.2) \quad \Rightarrow \quad 2t\|c\|^2 &\leq \|a\| + t^2\|b\|\|c\|^2 \end{aligned}$$

With a fairly nice quadratic polynomial in $\mathbb{R}[t]$ calved by (3.2) in our hands witch is allways non negative, the discriminant must be non positive. Therefore:

$$\begin{aligned} (-2\|c\|^2)^2 - 4\|b\|\|c\|^2\|a\| &\leq 0 \\ (3.3) \quad \Rightarrow \quad \|\langle x, y \rangle\|^4 - \|\langle y, y \rangle\|\|\langle x, y \rangle\|^2\|\langle x, x \rangle\| &\leq 0 \end{aligned}$$

Assuming $\|\langle x, y \rangle\|^2 \neq 0$ means that (3.3) can be simplified into Cauchy-Schwartz inequality (3.1) by cancelling $\|\langle x, y \rangle\|^2$. Otherwise¹, $\langle x, y \rangle = 0$ is a trivial case of the desired inequality. \square

For any A -valued inner product as above, we define a norm $\|x\| := \sqrt{\|\langle x, x \rangle\|_A}$ on a Inner product C^* -module. Which means that for arbitrary $x, y \in E$ and $a \in A$, the following holds:

¹Note that $\|\langle x, y \rangle\|^2 = 0$ if and only if $\langle x, y \rangle = 0$.

- (i) $\|x\| = 0 \Leftrightarrow x = 0$.
- (ii) $\|xa\| = \|a\|_A \|x\|$.
- (iii) $\|x + y\| \leq \|x\| + \|y\|$.

Notice that the triangle inequality (iii) is a direct consequence of 3.1.2:

$$\begin{aligned}
 \|x + y\|^2 &= \|\langle x + y, x + y \rangle\|_A \\
 &= \|\langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle\|_A \\
 &\leq \|x\|^2 + \|\langle x, y \rangle\|_A + \|\langle x, y \rangle^*\|_A + \|y\|^2 \\
 &= \|x\|^2 + 2\|\langle x, y \rangle\|_A + \|y\|^2 \\
 &\stackrel{3.1.2}{\leq} \|x\|^2 + 2\|x\|\|y\| + \|y\|^2 = (\|x\| + \|y\|)^2
 \end{aligned}$$

as in the good old days. One identity that still remains is the polarization one: For every sesquilinear form $\varsigma : E \times E \rightarrow A$

$$(3.4) \quad 4\varsigma(y, x) = \sum_{n=0}^3 i^n \varsigma(x + i^n y, x + i^n y). \quad (x, y \in E)$$

Since it should be a normed space, hence a complex vector space, one may be concerned about the fact that A doesn't necessarily have a unit and therefore, zx for $z \in \mathbb{C}$ should be an worry.

Proposition 3.1.3. All Inner product modules are naturally complex vector spaces, even the ones over non necessarily unital C^* -algebras.

Proof. Any Inner product module E is a \mathbb{Z} -module naturally because it is an abelian group with respect to the addition, and so is that $-\langle x, y \rangle = \langle x, -y \rangle$. Therefore, since the proof of Cauchy-Schwartz inequality 3.1.2 doesn't depend on the unity of A , we safe unitl now. For any approximate unit $(u_\lambda)_\lambda \subset A$, $(xu_\lambda)_\lambda \subset E$ converges to x , whence, for $z \in \mathbb{C}$, let $zx := \lim_\lambda x(zu_\lambda)$. Since A is a vector space, all properties are guaranteed and we are done. \square

Definition 3.1.4. Inner product modules are called *Hilbert C^* -modules* when the induced norm is complete in the Cauchy sense.

Proposition 3.1.5. For a Hilbert C^* -module E over A , let EA denote the linear span of elements given by xa , for $x \in E$ and $a \in A$. Therefore $\overline{EA} = E$.

Proof. If $(u_\lambda)_\lambda \subset A$ is a approximate unit for A , then for all $x \in E$:

$$\begin{aligned}
 \lim_\lambda \langle x - xu_\lambda, x - xu_\lambda \rangle &= \lim_\lambda (\langle x, x \rangle - u_\lambda \langle x, x \rangle) \\
 &\quad - \lim_\lambda (\langle x, x \rangle u_\lambda - u_\lambda \langle x, x \rangle u_\lambda) = 0.
 \end{aligned}$$

Hence the elements of the form xu_λ are dense in E . \square

Remark 3.1.6. Let A and B be C^* -algebras. If E is a Hilbert B -module and the ideal I of the closure of the elements spanned by $\langle x, y \rangle$ is contained in A , then there is a way to make E into a Hilbert A -module without changing the inner product. Namely, let $(u_\lambda)_\lambda$ be an approximate unit for I . Then the identity

$$\begin{aligned} \langle xu_\eta a - xu_\lambda a, xu_\eta a - xu_\lambda a \rangle &= a^* u_\eta \langle x, x \rangle u_\eta a + a^* u_\lambda \langle x, x \rangle u_\lambda a \\ &\quad - a^* u_\eta \langle x, x \rangle u_\lambda a - a^* u_\lambda \langle x, x \rangle u_\eta a, \end{aligned}$$

holds for all $x \in E$ and $a \in A$, showing that $(xu_\lambda a)_\lambda$ converges in E . We can define $xa = \lim xu_\lambda a$, and it is straightforward to check that this makes E into a Hilbert A -module. This is particularly when dealing with non unital C^* -algebras A , and we might have a look into the same module over \tilde{A} . \blacksquare

Examples 3.1.7.

- (i) Any traditional complex Hilbert space is a Hilbert \mathbb{C} -module.
- (ii) Let $(E_i)_{i \in I}$ be a family of Hilbert C^* -modules over A . The direct sum will be:

$$\bigoplus_{i \in I} E_i := \left\{ x \in \prod_{i \in I} E_i \mid \sum_{i \in I} \langle x_i, x_i \rangle \in A \right\}$$

It should be noticed that the convergence of $\sum_i \langle x_i, x_i \rangle$ is a weaker condition than requiring that the series of norms $\sum_i \|\langle x_i, x_i \rangle\|$ should converge. With the addition inner product $\langle x, y \rangle = \sum_i \langle x_i, y_i \rangle_{E_i}$, $\bigoplus_i E_i$ is a Hilbert C^* -module itself.

- (iii) Subexamples of (ii) are: A it-self endowed with $\langle a, b \rangle := a^* b$; $A^n = \bigoplus_{i=1}^n A$ for any natural number n .
- (iv) **The standard Hilbert A -module \mathcal{H}_A :** A more especif subexample of (ii) can be given by $\mathcal{H}_A := \bigoplus_{n \in \mathbb{N}} A$, consisting of all sequences $(a_n)_n \subset A$ which $\sum_n a_n^* a_n$ converges.
- (v) Given a Hilbert space H , the algebraic tensor product of H by A can be seeing as a Inner product C^* -module, with the bond:

$$\langle x \otimes a, y \otimes b \rangle := \langle x, y \rangle_H a^* b$$

$H \otimes A$ stands for its completion.

- (vi) Let $X \in \mathbf{CHaus}$ and $E \rightarrow X$ a complex vector bundle. As we mention, $C(X)$ is a unital C^* -algebra. Whenever $d : E \times E \rightarrow [0, \infty)$ is an Hermitian metric over E , the set $\Gamma(E)$ of continuous sections over E holds the title of Hilbert module over $C(X)$ when endowed with

$$\begin{aligned} \langle \cdot, \cdot \rangle : \Gamma(E) \times \Gamma(E) &\longrightarrow C(X) \\ (a, b) &\longmapsto d(a(\cdot), b(\cdot)) \end{aligned}$$

as an inner product.



Lemma 3.1.8. Given two nets $(x_\lambda)_\lambda$ and $(y_\lambda)_\lambda$ and x, y in a Hilbert module E over a C^* -algebra A such that $x_\lambda \rightarrow x$ and $y_\lambda \rightarrow y$, $\lim_\lambda \langle x_\lambda, y_\lambda \rangle = \langle x, y \rangle$ holds.

Proof. From the Cauchy-Schartz inequality 3.1.2, is easy to obtain that

$$\|\langle x_\lambda - x, z \rangle\|_A \stackrel{(3.1)}{\leq} \|x_\lambda - x\|_E \|z\|_E \quad (z \in E, \lambda \in \mathbb{A})$$

Analogously, $\|\langle z, y_\lambda - y \rangle\|_A \leq \|y_\lambda - y\|_E \|z\|_E$. For each and every index λ , it is possible to obtain the following inequality:

$$\begin{aligned} \|\langle x_\lambda, y_\lambda \rangle - \langle x, y \rangle\| &= \|\langle x_\lambda, y_\lambda \rangle - \langle x_\lambda, y \rangle + \langle x_\lambda, y \rangle - \langle x, y \rangle\| \\ &\leq \|\langle x_\lambda, y_\lambda - y \rangle\| + \|\langle x_\lambda - x, y \rangle\| \\ &\leq \|y_\lambda - y\| \|x_\lambda\| + \|y\| \|x_\lambda - x\| \end{aligned}$$

Let $\varepsilon > 0$. Notice that $x_\lambda \rightarrow x$, means that $\|x_\lambda\| \rightarrow \|x\|$. By $\|y_\lambda - y\| \rightarrow 0$, there exists λ_0 in which $\|y_\lambda - y\| \|x_\lambda\| < \varepsilon/2$. Similarly, there allways exists λ_1 such that $\|x_\lambda - x\|_E < \varepsilon/2(\|y\| + 1)$ for $\lambda \succcurlyeq \lambda_1$. Since it exists λ_2 such that $\lambda_2 \succcurlyeq \lambda_0$ and $\lambda_2 \succcurlyeq \lambda_1$, we conclude that $\|\langle x_\lambda, y_\lambda \rangle - \langle x, y \rangle\| < \varepsilon$ for all $\lambda \succcurlyeq \lambda_2$. \square

3.2 Adjointable operators

Definition 3.2.1 (Adjoint). Let E, F be Hilbert modules over a C^* -algebra A . A function $T : E \rightarrow F$ is said to be *adjointable* if there exists a function $T^* : F \rightarrow E$ which satisfies the following relation:

$$\langle Tx, y \rangle_F = \langle x, T^*y \rangle_E \quad ((x, y) \in E \times F)$$

Besides talking about Hilbert modules, we had defined the adjoint concept for any function between Hilbert modules. That's because inner product relations naturally require these functions to be linear operators, and if they exist, the adjoint is unique. For an adjointable T , T^* is unique, adjointable and $T^{**} = T$. Moreover, $(ST)^* = T^*S^*$ for adjointable operators T and S .

Proposition 3.2.2. Every adjointable operator $T : E \rightarrow F$, between Hilbert A -modules is bounded and continuous.

Proof. Since the set $\{\langle Tx, y \rangle_F = \langle x, T^*y \rangle_E \mid \|x\| \leq 1\} \subset A$ is bounded for all $y \in F$, Banach-Steinhaus theorem 3.2.3 implies that T is bounded.

Summoning 3.2.3 (Banach-Steinhaus “Uniform Boundness Principle” - [20]). Let \mathcal{F} be a family of bounded linear operators from a Banach space X to a normed linear space Y . If \mathcal{F} is

pointwise bounded, then \mathcal{F} is norm-bounded, i.e.,


$$\forall x \in X, \sup_{T \in \mathcal{F}} \|Tx\| < \infty \Rightarrow \sup_{T \in \mathcal{F}} \|T\| < \infty.$$



By 3.1.8, for a convergent net $x_\lambda \rightarrow x$, the following holds for all $y \in E$:

$$\begin{aligned} 0 &= \lim_\lambda [\langle T^*y, x_\lambda \rangle - \langle T^*y, x \rangle] \\ &= \lim_\lambda \langle y, Tx_\lambda \rangle - \lim_\lambda \langle T^*y, x_\lambda \rangle \\ &= \langle y, \lim_\lambda Tx_\lambda \rangle - \langle T^*y, \lim_\lambda x_\lambda \rangle \\ &= \langle y, \lim_\lambda Tx_\lambda \rangle - \langle T^*y, x \rangle \\ &= \langle y, \lim_\lambda Tx_\lambda \rangle - \langle y, Tx \rangle = \langle y, \lim_\lambda Tx_\lambda - Tx \rangle \end{aligned}$$

Especially when $y := \lim_\lambda Tx_\lambda - Tx$, so that $\lim_\lambda Tx_\lambda = Tx$. \square

Example 3.2.4. Given $x, y \in E$, the maps $y\langle x, \cdot \rangle$ and $x\langle y, \cdot \rangle$ are adjoints of each other. The linear span of those operator are what we will call the *finite rank* operators. 

In traditional Hilbert spaces, every bounded operator is adjointable, thanks to the Riesz Lemma ([17]). But when talking about Hilbert modules, that can't be the case anymore:


Counterexample 3.2.5 (*Non-adjointable bounded operator* - [15]). Suppose that J is a closed right ideal of a unital C^* -algebra A such that no element of J^* acts as a left multiplicative identity on J^2 . Consider the right module $J \times A$ with inner product defined by

$$\langle (a_1, b_1), (a_2, b_2) \rangle = a_2^*a_1 + b_2^*b_1$$

for $a_1, a_2 \in J$ and $b_1, b_2 \in A$. In this new space we have $\|(a, b)\|_{J \times A}^2 = \|a^*a + b^*b\| \leq \|a\|^2 + \|b\|^2$, hence $\|\cdot\|_{J \times A}$ is complete, i.e., $J \times A$ is a Hilbert module.

The operator $T(a, b) := (0, a)$ for each $(a, b) \in J \times A$ is obviously a bounded one, but if we suppose that there exists T^* and let $(x, y) := T^*(0, 1)$, notice that

$$a = \langle T(a, b), (0, 1) \rangle = \langle (a, b), T^*(0, 1) \rangle = x^*a + y^*b$$

for all (a, b) . Necessarilly, it is the case that $y = 0$ and $x^*a = a$ for all $a \in J$, hence x^* acts as a left multiplicative identity on J . But $x^* \in J^*$, and this contradicts our assumption of J . 

²For instance, the algebra of complex valued continuous functions on the unit interval $C[0, 1]$, and the ideal $C_0[0, 1]$ of functions which vanish at 0.

Counterexample 3.2.6. Let $X \in \mathbf{CHaus}$ and $Y \subset X$ a closed non-empty subset with dense complement. Let $E := \{f \in C(X) \mid f(Y) = \{0\}\}$ and $\iota : E \hookrightarrow C(X)$ the bounded inclusion map. If ι were adjointable, $E \ni \iota^*(I_{C(X)}) = I_{C(X)} \notin E$, i.e., the inclusion is a non-adjointable bounded operator. \blacksquare

Proposition 3.2.7. With the operator norm $\|T\| := \sup_{\|x\|=1} \|Tx\|$, the adjointable operators $\mathcal{L}(E, F)$ is C^* -algebra.

Proof. It is straightfoward checking that $\mathcal{L}(E, F)$ is an involution Banach algebra. To check the C^* -norm property, for each adjointable T ,

$$(3.5) \quad \begin{aligned} \|Tx\|^2 &= \|\langle Tx, Tx \rangle\| = \|\langle T^*Tx, x \rangle\| \stackrel{3.1.2}{\leq} \|T^*Tx\| \|x\| \\ &\leq \|T^*T\| \|x\|^2 \leq \|T^*\| \|T\| \|x\|^2 \end{aligned}$$

for all $x \in E$. A direct calculation using (3.5), shows that

$$\|T\|^2 = \sup_{\|x\|=1} \|Tx\|^2 \leq \sup_{\|x\|=1} \|T^*\| \|T\| \|x\|^2 = \|T^*\| \|T\|,$$

which means: $\|T\| \leq \|T^*\|$ and by extension, $\|T\| = \|T^*\|$. This automatically guarantee the C^* -norm property. \square

Proposition 3.2.8 (Equivalence of positivity). Being positive some element T in the C^* -algebra $\mathcal{L}(E)$ of adjointable automorphisms of a Hilbert C^* -module is equivalent to being positive in the inner algebra: $\langle Tx, x \rangle \geq 0$ for all $x \in E$.

Proof. Assume that T is a positive element in the C^* -algebra of operators. In a C^* -algebra, we know that $a \geq 0 \Leftrightarrow a = b^*b$ for some b . With this in hands, let S be such that $T = S^*S$. Therefore:

$$(3.6) \quad \langle Tx, x \rangle = \langle S^*Sx, x \rangle = \langle Sx, Sx \rangle \stackrel{3.1.1(iii)}{\geq} 0. \quad (x \in E)$$

Conversely, positive elements are self-adjoint, i.e., $\langle Tx, x \rangle = \langle x, Tx \rangle$. From the polarization identity 3.4, one can see that $\langle Tx, y \rangle = \langle x, Ty \rangle$ for all $x, y \in E$, showing that T is self-adjoint. By the Hahn decomposition³, there exists two positive elements T_+ and T_- such that $T = T_+ - T_-$ and $T_+T_- = T_+T_- = 0$. Then $\langle T_-y, y \rangle \leq \langle T_+y, y \rangle$ for any $y \in E$. In particular,

$$\langle T_-^3x, x \rangle = \langle T_-^2x, T_-x \rangle \leq \langle T_+T_-x, T_-x \rangle = 0.$$

On the other hand, $T_- \geq 0$ and $T_-^3 \geq 0$, hence $\langle T_-^3x, x \rangle \geq 0$ (because the statement in this direction is already proved). So the only possibility left is $\langle T_-^3x, x \rangle = 0$ for any x . By the polarization equality 3.4, this implies $\langle T_-^3x, y \rangle = 0$ for all $x, y \in \mathcal{M}$, hence $T_-^3 = 0$, $T_- = 0$. Thus, $T = T_+ \geq 0$. \square

³Every element $a \in A$ in a C^* -algebra can be written as $a = a_+ - a_-$ with $a_+, a_- \geq 0$, $a_+a_- = a_-a_+ = 0$. This is called the Hahn decomposition.

Proposition 3.2.9. If $T \in \mathcal{L}(E, F)$ and $x \in E$, then $\langle Tx, Tx \rangle \leq \|T\|^2 \langle x, x \rangle$.

Proof. Let ρ be a state of A . By repeated application of the Cauchy-Schwartz inequality to $\rho(\langle \cdot, \cdot \rangle)$:

$$\begin{aligned}
 \rho(\langle T^*Tx, x \rangle) &\leq \rho(\langle T^*Tx, T^*Tx \rangle)^{\frac{1}{2}} \rho(\langle x, x \rangle)^{\frac{1}{2}} \\
 &= \rho(\langle (T^*T)^2x, x \rangle)^{\frac{1}{2}} \rho(\langle x, x \rangle)^{\frac{1}{2}} \\
 &\leq \rho(\langle (T^*T)^4x, x \rangle)^{\frac{1}{4}} \rho(\langle x, x \rangle)^{\frac{1}{2} + \frac{1}{4}} \\
 &\vdots \\
 &\leq \rho(\langle (T^*T)^{2^n}x, x \rangle)^{\frac{1}{2^n}} \rho(\langle x, x \rangle)^{\frac{1}{2} + \frac{1}{4} + \dots + \frac{1}{2^n}} \\
 &\leq \|x\|^{2^{1-n}} \|T^*T\| \rho(\langle x, x \rangle)^{1 - \frac{1}{2^n}}
 \end{aligned}$$

As $n \rightarrow \infty$, one deduces that $\rho(\langle Tx, Tx \rangle) \leq \|T\|^2 \rho(\langle x, x \rangle)$. Since this is true for all states ρ , the desired inequality holds. \square

3.3 Compact and Finite rank operators

We are heading towards the definition of generalized Fredholm operators between Hilbert modules, and for that, we need a replacement for the finite dimensional condition. Henceforth, we shall explore the example 3.2.4.

Let M be a Hilbert A -module. Consider the following operator:

$$\begin{aligned}
 (3.7) \quad \Omega : M^n &\longrightarrow \mathcal{L}(A^n, M) \\
 (x_i)_i &\longmapsto \left((a_i)_i \xrightarrow{\Omega_x} \sum_{i=1}^n x_i a_i \right)
 \end{aligned}$$

In order to obtain the adjoint operator, as far as algebraic manipulation goes, $\Omega_x^* : M \longrightarrow A^n$ has no other option else besides being the coordinate inner decomposition $(\langle x_i, \cdot \rangle)_i$: For $a \in A^n$ and $\xi \in M$,


$$\begin{aligned}
 \langle \Omega_x a, \xi \rangle_M &= \left\langle \sum_{i=1}^n x_i a_i, \xi \right\rangle_M \\
 &= \sum_{i=1}^n a_i^* \langle x_i, \xi \rangle_M \\
 &= \begin{bmatrix} a_1^* & \cdots & a_n^* \end{bmatrix} \begin{bmatrix} \langle x_1, \xi \rangle_M \\ \vdots \\ \langle x_n, \xi \rangle_M \end{bmatrix} \\
 &= \left\langle a, (\langle x_1, \xi \rangle_M, \dots, \langle x_n, \xi \rangle_M) \right\rangle_{A^n} = \langle a, \Omega_x^* \xi \rangle_{A^n}
 \end{aligned}$$

One should note that for $x \in M^n$ and $y \in N^n$, $\Omega_y \Omega_x^*$ rises a fair notion of *finite rank*, since it's image elements are given by

$$(3.8) \quad \Omega_y \Omega_x^* \xi = \Omega_y^n (\langle x_1, \xi \rangle_M, \dots, \langle x_n, \xi \rangle_M) = \sum_{i=1}^n y_i \langle x_i, \xi \rangle_M.$$

Definition 3.3.1 (Compact and finite-rank operators). Every operator of the form $\Omega_y \Omega_x^* : M \rightarrow N$, where $(x, y) \in M^n \times N^n$ will be said to be a *A-finite rank operator*. The set of finite-rank operators will be denoted by $\text{FR}_A(M, N)$. The set of *A-compact operators* between M and N are defined as the topological closure of $\text{FR}(M, N)$ and it's denoted as $\mathcal{K}_A(M, N)$.

Unfortunately, *A-compact* operators need not to be compact in the sense of Banach spaces:

Counterexample 3.3.2. In a unital C^* -algebra A , the identity can be viewed as $\Omega_1 \Omega_1^* = I_A$ on the Hilbert module A . Hence $I_A \in \mathcal{K}(A)$, but it is a compact operator on the Banach space A if and only if A is finite-dimensional, since it is a invertible compact⁴. 

Proposition 3.3.3. In the standard Hilbert module \mathcal{H}_A over a unital C^* -algebra A , the classical compact notion of compact operator is well rescued: If $L_n(A) \subset \mathcal{H}_A$ denotes the free submodule generated by the first n canonical elements e_i ($i \leq n$), the following are equivalent:

- (i) $K \in \mathcal{K}_A(A)$.
- (ii) The norms of restrictions of K onto the orthogonal complements $L_n(A)^\perp$ of the submodules $L_n(A)$ vanish as $n \rightarrow \infty$.

Proof.

(i) \Rightarrow (ii) Let $p_n : \mathcal{H}_A \rightarrow L_n(A)^\perp$ by the orthogonal projection. Then, for any $z \perp L_n(A)$, one has

$$\begin{aligned} \|\Omega_x \Omega_y^* z\|^2 &= \|\langle \Omega_x \Omega_y^* z, \Omega_x \Omega_y^* z \rangle\| \\ &= \|\langle x \langle y, z \rangle, x \langle y, z \rangle \rangle\| \\ &= \|\langle y, z \rangle^* \langle x, x \rangle \langle y, z \rangle\| \\ &\leq \|x\|^2 \|\langle y, z \rangle\|^2 \\ &= \|x\|^2 \|\langle p_n y, z \rangle\|^2 \\ &\leq \|x\|^2 \|p_n y\|^2 \|z\|^2. \end{aligned}$$

⁴If $T \in \text{Hom}_{\mathcal{B}\text{an}}(X, Y)$ is a invertible compact operator between Banach spaces, the boundedness of T^{-1} rises a constant C such that $\|T^{-1}y\| \leq C\|y\|$, and by invertibility, $\|x\| \leq C\|Tx\|$. Thus, the image by T of the unit ball in X contains an open ball in Y . Since T is compact, Y is finite-dimensional, and so do X .

Since $\|p_n y\|$ tends to zero, the same is true for the norm of the restriction of the operator $\Omega_x \Omega_y^*$ to the submodule $L_n(A)^\perp$, hence, for the norm of any compact operator K .

- (i) \Leftarrow (ii) For a operator $K \in \mathcal{L}(\mathcal{H}_A)$, suppose that $\|K \upharpoonright_{L_n(A)^\perp}\|$ vanishes when $n \rightarrow \infty$. If z is an orthogonal element with respect to $L_n(A)$, i.e., $\langle e_i, z \rangle = 0$ for any $i \leq n$, notice that if $e^n := (e_1, \dots, e_n)$, then $\Omega_{K e^n} \Omega_{e^n}^*(z) = 0$. Therefore:

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sup_{\substack{z \in L_n(A)^\perp \\ \|z\| \leq 1}} \left\| Kz - \sum_{i=1}^n K e_i \langle e_i, z \rangle \right\| \\ &= \lim_{n \rightarrow \infty} \sup_{\substack{z \in L_n(A)^\perp \\ \|z\| \leq 1}} \|Kz\| = \lim_{n \rightarrow \infty} \|K \upharpoonright_{L_n(A)^\perp}\| = 0. \end{aligned}$$

If $z \in L_n(A)$, one can see that $Kz = \Omega_{K e^n} \Omega_{e^n}^*(z)$, so that the supreme can be taken in the hole \mathcal{H}_A . Therefore:

$$K = \lim_{n \rightarrow \infty} \sum_{i=1}^n \Omega_{K e_i} \Omega_{e_i}^* \in \mathcal{K}_A(A). \quad \square$$

Examples 3.3.4.

- (i) Let A be your favorite C^* -algebra. We shall see that $\mathcal{K}(A) \simeq A$. Notice that the map

$$\begin{aligned} L: A &\longrightarrow \mathcal{L}(A) \\ a &\longmapsto (b \mapsto L_a b) \end{aligned}$$

is well defined since L_{a^*} is the adjoint of L_a . It is no mistery that $\|L_a\| \leq \|a\|$, but notice that $\|L_a(a^*)\| = \|a\| \|a^*\|$, hence L is an isomorphism of A onto a closed $*$ -subalgebra of $\mathcal{L}(A)$.

Since $a \langle b, \cdot \rangle = L_{ab^*}$, it follows that $\mathcal{K}(A)$ is the clousure of (the image under L of) the linear span of products in A . But every C^* -algebra contains an approximate identity, such products are dense, thus L is an isomorphism between A and $\mathcal{K}(A)$.

- (ii) If A is a unital algebra, $\mathcal{K}(A) = \mathcal{L}(A)$ since any adjointable operator T consists of left multiplication by $T(1)$.
- (iii) Reviewing 3.1.7(v), one can obtain that $\mathcal{K}(H \otimes A) \simeq \mathcal{K}(H) \otimes A$.
- (iv) $\mathcal{K}(E^m, F^n) \simeq \mathbb{M}_{m \times n}(\mathcal{K}(E, F))$ and $\mathcal{L}(E^m, F^n) \simeq \mathbb{M}_{m \times n}(\mathcal{L}(E, F))$.



Proposition 3.3.5. For each $x \in M^n$, Ω_x is a A -compact operator.

Proof. Let $(u_\lambda)_\lambda \subset A$ be an approximate unit. Considering A it-self as a Hilbert module, $\Omega_{u_\lambda}^* : A^1 \longrightarrow A$ is the standard multiplication by u_λ . Therefore, for each $a \in A$:

$$(3.9) \quad \begin{aligned} \Omega_x a &= x a = x \lim_\lambda u_\lambda a \\ &\stackrel{u_\lambda = u_\lambda^*}{=} \lim_\lambda x \langle u_\lambda, a \rangle_A \stackrel{(3.8)}{=} \lim_\lambda \Omega_x \Omega_{u_\lambda}^* a \end{aligned}$$

So every finite 1-rank operators is indeed compact. For a general n , we arrive at a sum of finite 1-rank operators. Given $x \in M^n$ and $a \in A^n$, for each λ , we wish to obtain $y \in M^n$ and $b \in (A^n)^n$ such that,

$$\sum_{i=1}^n \Omega_{x_i} \Omega_{u_\lambda}^* a_i \stackrel{(*)}{=} \Omega_y \Omega_b^* a$$

in order to write a general Ω_x as a compact operator. To address $(*)$, let $y := x$ and $b := \text{Diag}(u_\lambda) \in \mathbb{M}_{n \times n}(A) \simeq (A^n)^n$ the diagonal matrix whose non zero entries are u_λ . Notice that

$$(3.10) \quad \begin{aligned} \Omega_y \Omega_b^* a &= \Omega_x \Omega_{\text{Diag}(u_\lambda)}^* a \\ &= \sum_{i=1}^n x_i \underbrace{\langle b_i, a \rangle_{A^n}}_{\sum_{j=1}^n \langle b_{ij}, a_j \rangle_A} \\ &= \sum_{i=1}^n x_i \langle b_{ii}, a_i \rangle_A \quad (a \in A^n) \\ &\stackrel{u_\lambda = b_{ii}}{=} \sum_{i=1}^n \Omega_{x_i} \Omega_{u_\lambda}^* a_i \end{aligned}$$

Therefore, for any $a \in A^n$, the following holds and the claim is proved.

$$\Omega_x a = \sum_{i=1}^n x_i a_i \stackrel{(3.9)}{=} \lim_\lambda \sum_{i=1}^n \Omega_{x_i} \Omega_{u_\lambda}^* a_i \stackrel{(3.10)}{=} \lim_\lambda \Omega_x \Omega_{\text{Diag}(u_\lambda)}^* a. \quad \square$$

Definition 3.3.6 (Finite-rank Hilbert Module). A Hilbert Module M over an C^* -algebra A whose identity operator I_M is A -finite rank will be said to be an A -finite rank module.

Proposition 3.3.7. For an A -finite rank Hilbert module M over A , the family of finite rank automorphisms $\text{FR}(M)$ over A is a two-sided ideal of the adjointable operators $\mathcal{L}(M)$.

Proof. Extending coordinatewise, let $Ty := (Ty_1, \dots, Ty_n)$ and notice that, for $T, S \in \mathcal{L}(M)$,

$$\begin{aligned} T \Omega_y \Omega_x^* + \Omega_w \Omega_z^* S &= \sum_{i=1}^n T y_i \langle x_i, \cdot \rangle + \sum_{i=1}^n w_i \langle S^* z_i, \cdot \rangle \\ &= \Omega_{(Ty, w)} \Omega_{(x, S^* z)}^*. \end{aligned}$$

Therefore, any $\mathcal{L}(M)$ -linear combination of finite rank operators is itself a finite rank one, as showed above, i.e., $\text{FR}(M) \triangleleft \mathcal{L}(M)$. \square

Proposition 3.3.8. If the identity I_E in a Hilbert module E is a compact operator, then E has finite rank, i.e., $I_E \in \mathcal{K}(E) \Rightarrow I_E \in \text{FR}(E)$.

Proof. Suppose that I_M is a compact operator in a given Hilbert module E . By construction, $\text{FR}(E)$ is a dense subset of compact operators, so every open non empty set $U \subset \mathcal{K}(E)$ obeys

$$U \cap \text{FR}(E) \neq \emptyset \quad (U \subset \mathcal{K}(E))$$

Since $\mathcal{K}(E)$ is a unital C^* -algebra, the invertible operators $\text{GL}(\mathcal{K}(E))$ constitute a non empty open set, hence there is a finite-rank invertible operator $F \in \text{GL}(\mathcal{K}(E)) \cap \text{FR}(E)$. Since $\text{FR}(E)$ contains an invertible and is an ideal, it follows that $I_E = FF^{-1} \in \text{FR}(E)$, i.e., the identity has finite rank. \square

In order to fully characterize finite rank Hilbert Modules over a given C^* -algebra, the same K -theoretic bias is seen right here through the representation by idempotents.

Theorem 3.3.9. A Hilbert module M has finite rank if, and only if, there exists an idempotent matrix $p \in \mathbb{M}_{n \times n}(A)$ such that M is isomorphic, as Hilbert A -modules, to pA^n .

Proof. Assume that M has finite rank, i.e., $I = \Omega_y \Omega_x^*$ for some $x, y \in M^n$. As presented in (3.11), $\Omega_x^* \Omega_y \in \mathcal{L}(A^n)$ is an idempotent operator, which corresponds to left multiplication by the idempotent matrix $p := (\langle x_i, y_j \rangle)_{i,j} \in \mathbb{M}_{n \times n}(A)$.

$$(3.11) \quad I = \Omega_y \Omega_x^* \Rightarrow \Omega_x^* = (\Omega_x^* \Omega_y) \Omega_x^* \Rightarrow \Omega_x^* \Omega_y = (\Omega_x^* \Omega_y)^2$$

$\Omega_x^* : M \longrightarrow pA^n$ is invertible: The middle term of (3.11) tells us that $\Omega_x^* = p\Omega_x^*$. Therefore, consider the following:

$$\begin{aligned} T : pA^n &\longrightarrow M \\ pa &\longmapsto \Omega_y a \end{aligned}$$

That operator show us that Ω_x^* is an invertible operator: Given $a \in A^n$, $\xi \in M$, one obtains that $\Omega_x^* T(pa) = \Omega_x^* \Omega_y a = pa$ and $T\Omega_x^* \xi = T(p\Omega_x^* \xi) = \Omega_y \Omega_x^* \xi = \xi$, i.e., $T = \Omega_x^{*-1}$.

Since it is an adjointable, functional continuous calculus allow us to extract the square root $|\Omega_x^*| := (\Omega_x \Omega_x^*)^{1/2}$, which is self-adjoint. Besides being

a linear bijection, Ω_x^* doesn't preserves inner products. But notice that $U := \Omega_x^* |\Omega_x^*|^{-1}$ does: For $\xi, \zeta \in M$,

$$\begin{aligned} \langle U\xi, U\zeta \rangle_{A^n} &= \langle \Omega_x^* |\Omega_x^*|^{-1} \xi, \Omega_x^* |\Omega_x^*|^{-1} \zeta \rangle_{A^n} \\ &= \langle \xi, |\Omega_x^*|^{-1} \Omega_x \Omega_x^* |\Omega_x^*|^{-1} \zeta \rangle_M \\ &= \langle \xi, |\Omega_x^*|^{-1} |\Omega_x^*|^2 |\Omega_x^*|^{-1} \zeta \rangle_M = \langle \xi, \zeta \rangle_M. \end{aligned}$$

Hence U is a Hilbert isomorphism between M and pA^n . \square

3.4 Kasparov Stabilization Theorem

In what follows through this text, one important application is a construction of a specific isomorphism between $K_0(A)$ and $K_0(B)$, without any separability condition assumed in the inner C^* -algebras A and B . Technically, we still need the main theorem of this section, requiring some countability condition. But keep calm and relax, since this enumerability shall be dribbled in Lemma 3.6.2.

We follow [14] in the following road map: Defining strictly positive elements in a C^* -algebra and characterizing operators in $\mathcal{K}(E)$ whose range is dense. With this characterization, we prove Kasparov stabilization theorem.

Definition 3.4.1. If a is a positive element in a C^* -algebra A and $\phi(a) \neq 0$ for all states⁵ ϕ on A , then a is said to be *strictly positive*.

Proposition 3.4.2. A positive element $a \geq 0$ in a unitary C^* -algebra is strictly positive if and only if it is a invertible element.

Proof due to [7]. Suppose $a \in A$ is strictly positive. Since A is unital, the state space of A is weak*-compact, and it follows that the element $\varepsilon := \inf\{\phi(a) \mid \phi \text{ is a state on } A\}$ is strictly positive, i.e., $\varepsilon > 0$. Therefore $a - \varepsilon$ is a positive element, so that the spectrum of a is contained in $[\varepsilon, \infty)$, i.e., a is invertible since 0 is not in the spectrum of a .


Conversely, suppose that a element $a \in A$ is positive and invertible. Whence, its spectrum is a compact subset of $(0, \infty)$, and thus $a - \varepsilon$ is positive for some $\varepsilon > 0$. If ϕ is a non-zero positive linear functional on A , we have

$$\phi(a) = \varepsilon\phi(1) + \phi(a - \varepsilon) \geq \varepsilon\|\phi\| + 0 > 0. \quad \square$$

Lemma 3.4.3. Let $a \in A$ be a positive element. Therefore a is strictly positive if, and only if, aA is dense in A .

Proof. Conjure the following statement:

⁵Norm 1 positive linear functional $\phi : A \longrightarrow \mathbb{C}$.

Summoning 3.4.4 ([4] - Lemma 2.9.4). Let A be a C^* -algebra and L, L' two closed left ideals of A such that $L \subseteq L'$. Suppose every positive form on A that vanishes on L also vanishes on L' . Then $L = L'$. 

Suppose that $L := aA \subset A =: L'$ isn't dense, i.e., $L \neq L'$. By the summoning, there is a state of A vanishing on aA . Such a state must vanish on a , so a is not strictly positive. Conversely, if ϕ is a state which $\phi(a) = 0$. Then, by Cauchy-Schwarz inequality for states:

$$|\phi(ab)|^2 \leq \phi(b^*b)\phi(a^*a) = 0 \quad (b \in A)$$

i.e., ϕ vanishes on aA , hence it is not dense. \square

Proposition 3.4.5. Let E be a Hilbert A -module and T a positive element in the C^* -algebra $\mathcal{K}(E)$. Then T is strictly positive if and only if T has dense range.

Proof. If T is strictly positive, by 3.4.3 then $\overline{T\mathcal{K}(E)} = \mathcal{K}(E)$. Since $\overline{\mathcal{K}(E)E} = E$, we have that $\overline{\text{Im } T} = \overline{T\mathcal{K}(E)E} = \overline{\mathcal{K}(E)E} = E$, i.e., T has dense image. Conversely, suppose that T has dense range. Therefore, for any $x, y \in E$, choose a sequence $(z_n)_n \subset E$ with $Tz_n \rightarrow x$. Therefore,

$$\Omega_x \Omega_y^* = \lim_{n \rightarrow \infty} T \Omega_{z_n} \Omega_y^* \in \overline{T\mathcal{K}(E)}.$$

So $T\mathcal{K}(E)$ is dense and T is strictly positive. \square

A Hilbert A -module M is countably generated if there is a sequence $(x_n)_n \subset X$ such that every x is the limit A -linear combinations of $(x_n)_n$.

Theorem 3.4.6 (Kasparov Stabilization Theorem). If M is a countably generated Hilbert A -module, then $\mathcal{H}_A \simeq M \oplus \mathcal{H}_A$.

Proof. Consider the case only when M is Hilbert \tilde{A} -module, which is sufficient since $\overline{MA} = M$ and $\overline{\mathcal{H}_A A} = \mathcal{H}_A$. Therefore, assume that A is a unital C^* -algebra.

Let $(\eta_n)_n$ be a bounded sequence of generators for M , with each generator repeated infinitely often. Let $(e_n)_n$ be the canonical orthonormal basis for \mathcal{H}_A , i.e., only the n -th coordinate of e_n is 1 and 0 elsewhere. Define $T : \mathcal{H}_A \rightarrow M \oplus \mathcal{H}_A$ linearly extending by $Te_n := 2^{-n}\eta_n + 4^{-n}e_n$. Notice that, for the elements $\zeta_n := \eta_n + 2^{-n}e_n$, T can be written as

$$T = \sum_{n=1}^{\infty} 2^{-n} \Omega_{\zeta_n} \Omega_{e_n}^* = \sum_{n=1}^{\infty} 2^{-n} (\eta_n + 2^{-n}e_n) \langle e_n, \cdot \rangle.$$

Therefore, T is a compact bijection. Since each η_n is repeated infinitely often, it is true that $\eta_n + 2^{-m}e_m = T(2^m e_m) \in \text{Im } T$ for infinitely many m which $\eta_n = \eta_m$. Going through the limit when $m \rightarrow \infty$, we see that

$$\eta_n + 0 = \lim_{m \rightarrow \infty} \eta_n + 2^{-m}e_m = \lim_{m \rightarrow \infty} T(2^m e_m) \in \overline{\text{Im } T}$$

and $0 + e_n = 4^n(Te_n - 2^{-n}(\eta_n + 0))$. Therefore, both $\eta_n + 0$ and $0 + e_n$ are in the closure $\overline{\text{Im } T}$.

Since $\{\eta_n + 0, 0 + e_n\}_n$ generates a dense submodule of $M \oplus \mathcal{H}_A$, T has dense range. Define new operators S and R given by $Se_n := 0 + 4^{-n}e_n$ and $Re_n := 2^{-n}\eta_n + 0$, in order that

$$\begin{aligned} T^*T &= S^*S + R^*R \\ &= \begin{pmatrix} 4^{-4} & 0 & 0 & \dots \\ 0 & 4^{-8} & 0 & \dots \\ 0 & 0 & 4^{-12} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} + \begin{pmatrix} 4^{-2}\langle\eta_1, \eta_1\rangle & 4^{-3}\langle\eta_1, \eta_2\rangle & 4^{-4}\langle\eta_1, \eta_4\rangle & \dots \\ 4^{-3}\langle\eta_2, \eta_1\rangle & 4^{-4}\langle\eta_2, \eta_2\rangle & 4^{-5}\langle\eta_2, \eta_3\rangle & \dots \\ 4^{-4}\langle\eta_3, \eta_1\rangle & 4^{-5}\langle\eta_3, \eta_2\rangle & 4^{-6}\langle\eta_3, \eta_1\rangle & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \end{aligned}$$

Hence $T^*T \geq S^*S$. Notice that S^*S has dense range, and by 3.4.5, it is strictly positive and so do T^*T . Therefore, $T|T|^{-1}$ is the desired isomorphism between those Hilbert modules. \square

Proposition 3.4.7. A Hilbert A -module E is countably generated if, and only if $\mathcal{K}(E)$ contains a strictly positive element, i.e., is σ -unital.

Proof. If A is unital and $(e_n)_n \subset \mathcal{H}_A$ is the canonical orthonormal basis, the operator $h := \sum_n 2^{-n}e_n\langle e_n, \cdot \rangle$ is a strictly positive element in $\mathcal{K}(\mathcal{H}_A)$ since it has dense range. If $P \in \mathcal{L}(\mathcal{H}_A)$ is an idempotent self-adjoint operator, then PhP is strictly positive in $\mathcal{K}(P\mathcal{H}_A)$. Finally, notice that A is countably generated if, and only if \tilde{A} is countably generated. Therefore $\mathcal{K}_A(E) = \mathcal{K}_{\tilde{A}}(E)$ is σ -unital if, and only if E is countably generated. \square

3.5 Rank definition of Finite Rank Hilbert Modules

In order to define the actual *rank* of a finite rank module M , we wish to deal with the K_0 -group as the K -algebraic theoreticals like, so we shall visualise our A -modules over the unification \tilde{A} of the underlying C^* -algebra A , regardless if it already possesses a unity or not. This process naturally enrich our module, as showed in 3.5.2.

Definition 3.5.1 (Projective module). Let M be an A -module. M is *projective* whenever there exists a map h such that it is commutative the following diagram:

$$\begin{array}{ccc} & M & \\ \swarrow h & \downarrow & \\ E & \twoheadrightarrow & F \end{array} \quad (E, F \in \mathbf{A}\text{-Mod})$$

That is to say, for every epimorphism⁶ $g : E \twoheadrightarrow F$ and $f : M \twoheadrightarrow F$, there always exists a map h such that $g \circ h = f$.

Proposition 3.5.2. Every finite rank Hilbert A -module is a finitely generated projective Hilbert \tilde{A} -module.

Proof. Let M be a finite rank Hilbert A -module. If $(a, \lambda) \in \tilde{A}$, letting $(a, \lambda) \cdot \xi := a\xi + \lambda\xi$ turns M into an Hilbert \tilde{A} -module. If $p \in \mathcal{L}(\tilde{A}^n)$ is the idempotent such that $M \simeq p\tilde{A}^n$ given by 3.3.9, notice that:

(i) $\tilde{A}^n \simeq M \oplus \ker p$: We shall see that it is a direct sum:

$$(3.12) \quad \tilde{A}^n = p\tilde{A}^n \oplus (I_{\tilde{A}^n} - p)\tilde{A}^n$$

Indeed, if $x \in p\tilde{A}^n \cap (I_{\tilde{A}^n} - p)\tilde{A}^n$, then $x = pa = (I_{\tilde{A}^n} - p)b$ for some tuples a, b . Therefore

$$x = pa = p(pa) = p(I_{\tilde{A}^n} - p)b = (p - p^2)b = 0,$$

and the sum in (3.12) is fact direct. Since $\ker p = (I_{\tilde{A}^n} - p)\tilde{A}^n$, the desired isomorphism holds by the givenness of p .

(ii) **If there exists Q such that $M \oplus Q$ is free, then M is projective:**

Let E, F be \tilde{A} -modules, $g : E \twoheadrightarrow F$ a surjective map and $f : M \twoheadrightarrow F$. Let $(b_i)_i$ be a basis of $M \oplus Q$. By surjectivity, for all i , there is always $x_i \in M$ such that $g(x_i) = f(b_i)$.

Define $\tilde{h} : M \oplus Q \rightarrow E$ extending linearly $\tilde{h}(b_i) := x_i$, in order that $g \circ \tilde{h} = f$. Therefore, $h := \tilde{h}|_M$ is the one necessary so that M is projective.

(iii) **If M is a direct summand of a free rank module, then it is finitely generated:** Let $M \oplus Q \simeq \tilde{A}^n$ for some \tilde{A} -module Q . That way, Q can be both projected and embbded in \tilde{A}^n by morphisms. Let $\psi : \tilde{A}^n \rightarrow \tilde{A}^n$ be the compositions of those. Then

$$\text{Im } \psi = \{(0, q) \in M \oplus Q \mid q \in Q\}$$

is the kernel of the canonical projection $\Pi_M : \tilde{A}^n \rightarrow M$. Therefore, the composition $\Pi_M \circ \psi : \tilde{A}^n \rightarrow M$ is a surjection, telling the world that M must be finitely generated.

Since \tilde{A}^n is free module, setting $Q := \ker p$, one can see that M is projective by (ii) and finitely generated by (iii). \square

Lemma 3.5.3. Let p, q be idempotent square matrices with entries living in A . The following are equivalent:

⁶For our purpose and needs, surjective morphism

- (i) As Hilbert modules, $pA^m \simeq qA^n$.
- (ii) p and q are Murray von-Neumann equivalent: There are r, s such that $p = rs$ and $q = sr$.

Proof.

- (i) \Rightarrow (ii) Let $T : pA^m \longrightarrow qA^n$ be an isomorphism. There exist unique matrices $r \in \mathbb{M}_{n \times m}(A)$ and $s \in \mathbb{M}_{m \times n}(A)$ which corresponds $s pa = T(pa)$ and $r qb = T^{-1}(qb)$ for every $a \in A^m$ and $b \in A^n$. Notice that

$$rs(pa) = T^{-1}T(pa) = pa = p^2a \quad \text{and} \quad sr(qb) = TT^{-1}(qb) = qb = q^2b$$

Therefore $p = rs$ and $q = sr$.

- (i) \Leftarrow (ii) The left multiplications maps $s : pA^m \longrightarrow qA^n$ and $r : qA^n \longrightarrow pA^m$ are mutual inverses of each other:

$$\begin{cases} s(r(qb)) = (sr)^2b = q^2b = qb, & (b \in A^n). \\ r(s(pa)) = (rs)^2a = p^2a = pa, & (a \in A^m). \end{cases}$$

Therefore $pA^m \simeq qA^n$. \square

For a A -finite rank module M , M can be viewed as a finitely generated projective \tilde{A} -module, hence let p be as in 3.3.9 embedded in $\mathbb{M}_\infty(\tilde{A})$. As an element of $V(\tilde{A})$, the equivalence class of idempotents which represents M , is the set:

$$[M] := \{q \in \mathbb{M}_\infty(\tilde{A}) \mid q^2 = q \in \mathbb{M}_n(\tilde{A}), M \simeq qA^n\} = [p] \in V(\tilde{A})$$

which by 3.5.3 is well defined.

Letting $[\cdot]_0 : V(\tilde{A}) \hookrightarrow K_0(\tilde{A})$ be the natural inclusion, $[q]_0 := [q] - [s(q)]$, one may see $[M]_0 \in K_0(\tilde{A})$ as an element of $K_0(A)$. In fact, if $\varepsilon : \tilde{A} \longrightarrow \mathbb{C}$ is the projection of the complex component, $\varepsilon(p) = 0$ since M is originally a Hilbert A -module. Therefore,

$$\begin{aligned} \varepsilon_0([M]_0) &= \varepsilon_0([p] - [s(p)]) = [\varepsilon(p)] - [\varepsilon(s(p))] = 0 \\ &\Rightarrow [M]_0 \in \ker \varepsilon_0 = K_0(A). \end{aligned}$$

Proposition 3.5.4. Let $P \in \mathcal{K}(E)$ be a compact self-adjoint idempotent operator over a Hilbert A -module E . Therefore, $\text{Im } P$ is an A -finite rank one.

Proof. Notice that $I_{\text{Im } P} = P$. Since it is compact in M , there are nets $(y_\lambda)_\lambda, (x_\lambda)_\lambda \subset E^\infty$ such that $P = \lim_\lambda \Omega_{y_\lambda} \Omega_{x_\lambda}^*$. Therefore:

$$\begin{aligned} I_{\text{Im } P} &= P = P^3 = P \left(\lim_\lambda \Omega_{y_\lambda} \Omega_{x_\lambda}^* \right) P \\ &= \lim_\lambda P \Omega_{y_\lambda} \Omega_{x_\lambda}^* P = \lim_\lambda \Omega_{Py_\lambda} \Omega_{P^*x_\lambda}^* \stackrel{P^*=P}{=} \lim_\lambda \Omega_{Py_\lambda} \Omega_{P^*x_\lambda}^*. \end{aligned}$$

In light of the 3.3.8, $\text{Im } P$ is indeed a finite-rank module. \square

Remark 3.5.5. The range of an idempotent operator coincide with the range of some projection, i.e., self-adjoint idempotent operator. To see this, suppose that $a \in A$ is an idempotent in a unital C^* -algebra A . Let

$$h := 1 + (a - a^*)(a^* - a) = 1 + aa^* - a^* - a + a^*a$$


With h in hands, one can draw the following conclusions:

(i) $h^* = h$.

(ii) Notice that $(a - a^*)(a^* - a) \geq 0$, hence $\text{Spec}((a - a^*)(a^* - a)) \subset [0, \infty)$. Conjure one of the classical theorems in spectral theory:

Summoning 3.5.6 (Spectral mapping theorem). Given an unital Banach algebra A and $a \in A$, the following holds:

$$\text{Spec}(p(a)) = \{p(\lambda) \mid \lambda \in \text{Spec}(a)\} = p(\text{Spec}(a))$$

for each and every complex polynomial $p \in \mathbb{C}[x]$. 

By the theorem 3.5.6,

$$\text{Spec}(h) = 1 + \text{Spec}((a - a^*)(a^* - a)) \subset [1, \infty).$$

Since $0 \notin \text{Spec}(h)$, h is invertible.

(iii) $ah = aa^*a = ha$ and $a^*h = a^*aa^* = ha^*$.

(iv) $p := aa^*h^{-1} = h^{-1}aa^*$. Indeed:

$$hp = haa^*h^{-1} = aa^*hh^{-1} = aa^*.$$


(v) p is self-adjoint:

$$p^* = (h^{-1})^*aa^* = (h^*)^{-1}aa^* = h^{-1}aa^* = p$$

(vi) p is idempotent:

$$p^2 = h^{-1} \underbrace{(aa^*a)}_{ha} a^* h^{-1} = h^{-1}h(aa^*h^{-1}) = p.$$

(vii) $pa = a$ and $ap = p$.

In particular, if $a := Q \in \mathcal{L}(E)$ is an idempotent operator, $\text{Im } Q$ coincides with the range of p by (vii), which is a self-adjoint idempotent operator. In particular $E = \text{Im } Q \oplus \text{Im } Q^\perp$. 

Definition 3.5.7. The *rank* of a finite rank Hilbert A -module will be defined as the class $\mathbf{rank}(M) := [p]_0 \in K_0(A)$. By proposition 3.5.4 and the remark 3.5.5, we shall also define the rank of a given compact idempotent operator P as the rank of $\text{Im } P$.

Lemma 3.5.8. Let E be a Hilbert module and $P, Q \in \mathcal{L}(E)$ be two compact idempotent operators, and let p and q be the idempotent square matrices given in the proof of Theorem 3.3.9. If P and Q are similar, i.e., there exists $u \in \text{GL } \mathcal{L}(E)$ such that $P = uQu^{-1}$, the matrices p and q are Murray-von Neumann equivalent.

Proof. We have that $P = uQu^{-1}$, $\text{Im } P \simeq pA^n$ and $\text{Im } Q \simeq qA^n$. Since their ranges are finite-rank modules (3.5.5 and 3.5.4), there are $x, y \in (\text{Im } P)^n$ and $z, w \in (\text{Im } Q)^m$ such that

$$P = I_{\text{Im } P} = \Omega_y \Omega_x^* \quad \text{and} \quad Q = I_{\text{Im } Q} = \Omega_w \Omega_z^*.$$

We know that $p = (\langle x_i, y_j \rangle)_{i,j}$ and $q = (\langle z_i, w_j \rangle)_{i,j}$ by the argument used in 3.3.9. Therefore:

$$\Omega_y \Omega_x^* = P = uQu^{-1} = u\Omega_w \Omega_z^* u^{-1} = \Omega_{u(w)} \Omega_{(u^{-1})^*(z)}^*,$$

which means that

$$\sum_{i=1}^n y_i \langle x_i, \cdot \rangle = \sum_{i=1}^m u(w_i) \langle (u^{-1})^*(z_i), \cdot \rangle.$$

Notice that the matrix $\hat{q} := (\langle (u^{-1})^*(z_i), u(w_j) \rangle)_{i,j}$ obeys $\text{Im } uQu^{-1} \simeq \hat{q}A^m$ by the construction of 3.3.9. But since $\langle (u^{-1})^*(z_i), u(w_j) \rangle = \langle z_i, w_j \rangle$, one concludes that $\hat{q} = q$, hence, we obtain two isomorphisms such that the following diagram commutes.

$$\begin{array}{ccccc} \text{Im } P & \xlongequal{\quad} & \text{Im } uQu^{-1} & \dashrightarrow & \text{Im } Q \\ \downarrow & & \downarrow & & \downarrow \\ pA^n & \dashrightarrow & qA^m & \xlongequal{\quad} & qA^m \end{array}$$

Since $pA^n \simeq qA^m$ if, and only if p and q are Murray-von Neumann equivalent by 3.5.3, the result follows. \square

Corollary 3.5.9. Let P and Q be compact idempotent operators in $\mathcal{L}(E)$. If $[P]_0 = [Q]_0$, then $\mathbf{rank}(P) = \mathbf{rank}(Q)$.

3.6 Quasi-stably-isomorphic Hilbert modules

If X, Y, Z and W are Hilbert A -modules and T is in $\mathcal{L}(X \oplus Y, Z \oplus W)$, then T can be represented by a matrix

$$T = \begin{pmatrix} T_{ZX} & T_{ZY} \\ T_{WX} & T_{WY} \end{pmatrix}$$

where T_{ZX} is in $\mathcal{L}(X, Z)$ and similarly for the other matrix entries. Matrix notation is used to define our next important concept.

We are in touch with pretty algebraic properties of Hilbert modules, and in our case of finite rank ones. Since those modules can be seen as projective finitely generated, an algebraist might convince you that two generators $[M]_0, [N]_0 \in K_0(A)$ are equal if, and only if they are *stably-isomorphic*, i.e., there exists n such that $M \oplus A^n \simeq N \oplus A^n$. We will present a generalization of this concept in terms of the *rank*, darkly hidden:

Definition 3.6.1 (Quasi-stably-isomorphic finite rank). Two Hilbert modules E and F are said to be *quasi-stably-isomorphic* if there exists a Hilbert module X and an invertible operator $T \in \text{GL } \mathcal{L}(E \oplus X, F \oplus X)$ such that $I_X - T_{XX}$ is compact.

Notice that this is an equivalence relation. Before lighting that relationship, first we enrich the definition on the special case of finite-rank modules.

Lemma 3.6.2. Assume M and N are A -finite rank modules. If M and N are quasi-stably-isomorphic then the module X referred to in 3.6.1 can be taken to be countably generated.

Proof. Let X and T as in 3.6.1. Using matrix notation, we have:

$$\begin{aligned} T : M \oplus X &\longrightarrow N \oplus X \\ (\xi, \eta) &\longmapsto \begin{pmatrix} T_{NM} & T_{NX} \\ T_{XM} & T_{XX} \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix} \end{aligned}$$

Since it is an invertible operator, consider the inverse given in matrix notation as well:

$$T^{-1} = \begin{pmatrix} S_{MN} & S_{MX} \\ S_{XN} & S_{XX} \end{pmatrix}$$

We shall construct inductively a enumerable collection $\mathcal{C} \subset X$ in order to generate a specific submodule. Such a construction will be given by a collection $(\mathcal{C}_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$, in order that each $\mathcal{C}_n \subset X$ satisfies:

- (i) The images of the operators T_{XM}, S_{XN}, T_{NX}^* and S_{XM}^* are contained in the submodule of X generated by \mathcal{C}_n .
- (ii) $I_X - T_{XX}$ can be approximated by finite rank operators of the form $\Omega_y \Omega_x^*$, where the components of x and y belong to \mathcal{C}_n .

For a given $i \leq n$, let π_i be the projection in the i -th coordinate of a tuple. In order to choose wisely, we write a technical issue:

Choice of generators: Let $r \in \mathcal{L}(P, X)$ where P is a finite rank Hilbert module. By assumption, there exists a natural number $n \in \mathbb{N}$ and tuples $x, y \in P^n$ such that $I_P = \Omega_y \Omega_x^*$. Notice that their existence depends on the domain of r . Therefore,

$$r = r \Omega_y \Omega_x^* = \sum_{i=1}^n r \pi_i(y) \langle \pi_i(x), \cdot \rangle$$

Hence, $\text{Im } r$ is contained in the submodule generated by $(r \pi_i(y))_{i \leq n}$.

In order to obey (i), our initial collection \mathcal{C}_0 must contain all elements of the form $r \pi_i(y(r))$ for r varying over the desired operators. Since $I_X - T_{XX}$ is a compact set, there exists tuple sequences $(\xi_n)_{n \in \mathbb{N}}, (\zeta_n)_{n \in \mathbb{N}}$ such that

$$I_X - T_{XX} = \lim_{n \rightarrow \infty} \Omega_{\zeta_n} \Omega_{\xi_n}^*.$$

Let \mathcal{C}_0 be given by:

$$\begin{aligned} \mathcal{C}_0 := & \{ r \pi_i(y) \mid r \in \{T_{XM}, S_{XN}, T_{NX}^*, S_{XM}^*\}, I_{\text{dom}(r)} = \Omega_y \Omega_x^* \}_{i \in \mathbb{N}} \\ & \cup \bigcup_{n \in \mathbb{N}} \{ \pi_i(\xi_n), \pi_i(\zeta_n) \}_{i \in \mathbb{N}} \end{aligned}$$

By construction, those properties are obeyed. Inductively, we set new collections in order to obey to above properties in terms of the operator T_{XX} and S_{XX} :

$$\begin{aligned} \mathcal{C}_{n+1} := & \mathcal{C}_n \cup T_{XX}(\mathcal{C}_n) \cup S_{XX}(\mathcal{C}_n) \\ & \cup T_{XX}^*(\mathcal{C}_n) \cup S_{XX}^*(\mathcal{C}_n) \end{aligned} \quad (n \in \mathbb{N})$$

in order that each \mathcal{C}_n satisfies (i) and (ii). Therefore, the union $\mathcal{C} := \bigcup_n \mathcal{C}_n$ is then obviously countable, and also obeys those same properties above. In addition, the following one belongs to package:

(iii) **\mathcal{C} is invariant under $T_{XX}, T_{XX}^{-1}, T_{XX}^*$ and $(T_{XX}^{-1})^*$.** Let $r \in \mathcal{L}(X)$ be one the operators. If it was the case that \mathcal{C} wasn't invariant over r , necessarily it would exists $w \in r(\mathcal{C}) \setminus \mathcal{C}$, hence, $r^{-1}(w) \in \mathcal{C}_n$ for some n . However,

$$w = r(r^{-1}(w)) \in \mathcal{C}_n \cup r(\mathcal{C}_n) \subset \mathcal{C}_{n+1} \subset \mathcal{C}$$

i.e., it cant be the case.

Let $X' := \overline{\langle \mathcal{C} \rangle}$ be the Hilbert submodule of X generated by \mathcal{C} . Because of (i) and (iii) we see that

$$T(M \oplus X') \subseteq N \oplus X' \quad \text{and} \quad T^*(N \oplus X') \subseteq M \oplus X'.$$

The restriction of T then gives an operator T' in $\mathcal{L}(M \oplus X', N \oplus X')$. The same reasoning applies to T^{-1} providing $(T^{-1})'$ in $\mathcal{L}(N \oplus X', M \oplus X')$ which is obviously the inverse of T' . In virtue of (ii) it is clear that T' satisfies the conditions of definition 3.6.1. \square

In terms of the Morita equivalence, this is a big deal since we are not restricting ourselves into C^* -algebras with countably approximate identities. After studying Fredholm operators between Hilbert modules, one shall construct group morphisms $K_*(A) \longrightarrow K_*(B)$ for generic C^* -algebras A and B . Those maps turn out to be isomorphisms when A and B are strongly Morita equivalent.

Theorem 3.6.3. Let M and N be quasi-stably-isomorphic finite rank Hilbert modules over a C^* -algebra A . Therefore, $\text{rank}(M) = \text{rank}(N)$.

Proof. Let $T \in \text{GL}(\mathcal{L}(M \oplus X, N \oplus X))$ with X being a countably generated Hilbert module (3.6.2) and $I_X - T_{XX} \in \mathcal{K}(X)$. By the countability condition, we can apply Kasparov's Stabilization Theorem 3.4.6 in order to obtain that $X \oplus \mathcal{H}_A \simeq \mathcal{H}_A$. Without loss of generality, we can assume that $X = \mathcal{H}_A$.

Since M is finitely generated as an A -module, by Kasparov's theorem again, there exists a isomorphism $\varphi : \mathcal{H}_A \longrightarrow M \oplus \mathcal{H}_A$. Now, we construct operators F and G given by the compositions:

$$\begin{aligned} F : \mathcal{H}_A &\xrightarrow{\varphi} M \oplus \mathcal{H}_A \xrightarrow{T} N \oplus \mathcal{H}_A \twoheadrightarrow \mathcal{H}_A \hookrightarrow M \oplus \mathcal{H}_A \xrightarrow{\varphi^{-1}} \mathcal{H}_A \\ G : \mathcal{H}_A &\xrightarrow{\varphi} M \oplus \mathcal{H}_A \twoheadrightarrow \mathcal{H}_A \hookrightarrow N \oplus \mathcal{H}_A \xrightarrow{T^{-1}} M \oplus \mathcal{H}_A \xrightarrow{\varphi^{-1}} \mathcal{H}_A \end{aligned}$$

Those are the first moments we are dealing with the so called generalized Fredholm operators. We state that

- (i) **Both $I - FG$ and $I - GF$ are compact:** Let $\Pi_M = I_M \oplus 0$ and $\Pi_{\mathcal{H}_A} = 0 \oplus I$ be the coordinate projections. When composing, one can simplify:

$$\begin{aligned} FG : \mathcal{H}_A &\xrightarrow{\varphi} M \oplus \mathcal{H}_A \xrightarrow{\Pi_{\mathcal{H}_A}} M \oplus \mathcal{H}_A \xrightarrow{\varphi^{-1}} \mathcal{H}_A \\ GF : \mathcal{H}_A &\xrightarrow{\varphi} M \oplus \mathcal{H}_A \xrightarrow{T} N \oplus \mathcal{H}_A \xrightarrow{\Pi_{\mathcal{H}_A}} N \oplus \mathcal{H}_A \xrightarrow{T^{-1}} M \oplus \mathcal{H}_A \xrightarrow{\varphi^{-1}} \mathcal{H}_A \end{aligned}$$

Meaning that

$$\begin{aligned} I - FG &= I - \varphi^{-1} \Pi_{\mathcal{H}_A} \varphi & \text{and} & & I - GF &= I - \varphi^{-1} T^{-1} \Pi_{\mathcal{H}_A} T \varphi \\ &= \varphi^{-1} \Pi_M \varphi & & & &= (T\varphi)^{-1} \Pi_N (T\varphi) \end{aligned}$$

Therefore, $I - FG$ and $I - GF$ are unitarily equivalent to Π_M and Π_N , which are compact operators.

(ii) **$I - FG$ and $I - GF$ are idempotents:** Notice that $F = \varphi^{-1}(0_M \oplus \Pi_{\mathcal{H}_A} T \varphi(\cdot))$. Therefore:

$$\begin{aligned} FGF &= \varphi^{-1}(0_M \oplus \Pi_{\mathcal{H}_A} T \varphi GF(\cdot)) \\ &= \varphi^{-1}(0_M \oplus \Pi_{\mathcal{H}_A} T \varphi \varphi^{-1} T^{-1} \Pi_{\mathcal{H}_A} T \varphi(\cdot)) \\ &= \varphi^{-1}(0_M \oplus \Pi_{\mathcal{H}_A} T \varphi(\cdot)) = F. \end{aligned}$$

Similarly, one can check that $GFG = G$. Hence, $I - FG$ and $I - GF$ are idempotents.

(iii) **F and G are compact perturbations of the identity:** The operators which ignores φ and φ^{-1} in the constructions of F and G , are represented by

$$F' := \begin{pmatrix} 0 & 0 \\ T_{\mathcal{H}_A M} & T_{\mathcal{H}_A \mathcal{H}_A} \end{pmatrix} \quad \text{and} \quad G' := \begin{pmatrix} 0 & S_{M \mathcal{H}_A} \\ 0 & S_{\mathcal{H}_A \mathcal{H}_A} \end{pmatrix}$$

where $S := T^{-1}$. Since $T_{\mathcal{H}_A \mathcal{H}_A}$ is a compact perturbation of the identity and both M and N are finite rank modules (and so any operator having either M or N as domain or codomain must be compact), one concludes that F' and G' , hence, F and G are indeed, compact perturbations of the identity.

For sake of notation, let $\mathcal{Q}(E) := \mathcal{L}(E)/\mathcal{K}(E)$ for a Hilbert A -module E . Since the compact set is an ideal of the adjointable maps, one can consider the following exact sequence of C^* -algebras:

$$(3.13) \quad 0 \longrightarrow \mathcal{K}(\mathcal{H}_A) \hookrightarrow \mathcal{L}(\mathcal{H}_A) \xrightarrow{\pi} \mathcal{Q}(\mathcal{H}_A) \longrightarrow 0$$

Notice that we can consider the index map ∂ induced by (3.13). As stated in item (iii), one concludes that $\pi(F) = 1_{\mathcal{Q}(\mathcal{H}_A)}$. Since I_2 is a lift of $\text{diag}(1_{\mathcal{Q}(\mathcal{H}_A)}, 1_{\mathcal{Q}(\mathcal{H}_A)}^{-1})$, one can obtain that $\partial([1_{\mathcal{Q}(\mathcal{H}_A)}]_1) = [I_2 p_2 I_2^{-1}]_0 - [p_2]_0 = 0$.

We can extract more information by writing down who is the index in terms of F and G . By (i), is easy to see that $\pi(F)$ and $\pi(G)$ are each others inverse in $\mathcal{Q}(\mathcal{H}_A)$. In order to compute the index, the element

$$w := \begin{pmatrix} F & I - FG \\ I - GF & G \end{pmatrix} \in \text{GL}_2^{(0)}(\mathcal{Q}(\mathcal{H}_A))$$

is a lift of $\text{diag}(\pi(F), \pi(F)^{-1})$, and its inverse just swaps F and G places. Therefore:

$$\begin{aligned} 0 &= \partial([\pi(F)]_1) \\ &= [w p_2 w^{-1}]_0 - [p_2]_0 \\ &= \left[\begin{pmatrix} F & I - FG \\ I - GF & G \end{pmatrix} \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} G & I - GF \\ I - FG & F \end{pmatrix} \right]_0 - \left[\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \right]_0 \\ &= \left[\begin{pmatrix} FG & 0 \\ 0 & I - GF \end{pmatrix} \right]_0 - \left[\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \right]_0 \\ &= [I - GF]_0 - [I - FG]_0 \end{aligned}$$

Hence $[I - FG]_0 = [I - GF]_0$ and by the corollary 3.5.9, one obtains that $\text{rank}(I - FG) = \text{rank}(I - GF)$. We had seen that $I - FG = \varphi^{-1}\Pi_M\varphi$, i.e., $\text{Im}(I - FG) \simeq M$. Therefore:

$$\text{rank}(M) = [M]_0 = [\text{Im}(I - FG)]_0 = \text{rank}(I - FG).$$

Similarly, one obtains that $\text{rank}(N) = \text{rank}(I - GF)$ and consequentially, $\text{rank}(M) = \text{rank}(N)$ as we were looking. \square

Chapter 4

Fredholm Operators

In regular Fredholm theory, one would define Fredholm operators as those with both kernel and cokernel were finite dimensional. Here, our concepts involving “finiteness” depends on a fixed C^* -algebra A , so, talking about dimension is kinda tricky. But classical Atkinson’s theorem characterizes Fredholm operators in terms only of compactness, which we fully characterize in Theorem 3.6.3. That points towards a generalization.

Definition 4.0.1 (A -Fredholm). A given $T \in \mathcal{L}(E, F)$ is said to be A -Fredholm if it is invertible modulo $\text{FR}(E, F)$, i.e., exists $S \in \mathcal{L}(F, E)$ such that both $I_E - ST$ and $I_F - TS$ are A -finite rank operators.

If adjointable operators S , S' and T are such that both TS and $S'T$ are Fredholm, one can obtain operators R and R' such that $I - T(SR)$ and $I - (R'S')T$ are finite-rank ones. Hence T is Fredholm.

Proposition 4.0.2. If T is invertible modulo $\mathcal{K}(E, F)$, then it is A -Fredholm. Hence, definition 4.0.1 really fits the sentence “Atkinson’s theorem trivialization”.

Proof. Let us digress a little bit:

Remark 4.0.3. Let $J \triangleleft A$ be a non-closed ideal of a unital C^* -algebra, and suppose that $a \in A$ is invertible modulo \overline{J} (the topological closure), i.e., there exists $b \in A$ such that $(1-ab) \in \overline{J}$. Therefore, there exists a sequence $(j_n)_n \subset J$ such that $j_n \rightarrow (1-ab)$, which means that at least one of them, say j , obeys $\|1-ab-j\| < 1$. Since $ab+j = 1-(1-ab-j)$, $ab+j$ is invertible by $??$. Therefore, some algebraic manipulation shows that

$$a \underbrace{b(ab+j)^{-1}}_{x_1} - 1 = -j(ab+j)^{-1} \in J$$

i.e., a is right invertible modulo J . But since b also ensure that $(1-ba) \in \overline{J}$, hence it exist x_2 such that $(x_2a - 1) \in J$. Notice

that

$$x_2(ax_1 - 1) = ((x_2a)x_1 - x_2) = (x_1 - x_2) \in J,$$

i.e., $[x_1] = [x_2]$ in the quotient algebra A/J . Therefore, there is a representative of the class of x_1 and x_2 , say x , such that both $ax - 1$ and $xa - 1$ belong to J , i.e., a is invertible modulo J . \blacksquare

With this digression in mind, notice that if $(I_E - TS) \in \mathcal{K}(E, F)$, there exist an element S such that both $I_E - TS$ and $I_F - ST$ are finite-rank operators, i.e., T is Fredholm. \square

Remark 4.0.4. Let H and W be complex Hilbert spaces. The set of classical Fredholm operators coincide with the \mathbb{C} -Fredholm ones. \blacksquare

Example 4.0.5 ([21]). Let $X \in \mathbf{CHaus}$ and $\mathcal{F}(\ell^2(\mathbb{N}))$ the set of \mathbb{C} -Fredholm classical operators over the separable Hilbert space $\ell^2(\mathbb{N})$. We avoid writing $\ell^2(\mathbb{N})$ everywhere, so \mathcal{B} , \mathcal{K} and \mathcal{F} will respectively denote the bounded, compact and Fredholm operators over $\ell^2(\mathbb{N})$.

For any continuous family of operators¹ $T : X \longrightarrow \mathcal{B}$, is possible to see T as a $C(X)$ -endomorphism over the standard Hilbert $C(X)$ -module $\mathcal{H}_{C(X)}$:

$$\begin{aligned} \widehat{T} : \mathcal{H}_{C(X)} &\longrightarrow \mathcal{H}_{C(X)} \\ \xi &\longmapsto (x \mapsto T_x \xi(x)) \end{aligned}$$

Each $T_x := T(x)$ is a classical operator in a Hilbert space, so $T^* \in C(X, \mathcal{B})$. Hence \widehat{T} is adjointable. In order to show that continuous families of Fredholm operators extent to $C(X)$ -Fredholm ones, we need the following claim:


- (i) **\widehat{T} is $C(X)$ -compact whenever T is continuous family of compact operators:** Let $\varepsilon > 0$ and $T \in C(X, \mathcal{K})$. For each $x \in X$, pick a finite rank operator R_x so that $\|R_x - T_x\| \leq \varepsilon/3$, and pick a neighborhood U_x such that $\|T_x - T_y\| \leq \varepsilon/3$ for $y \in U_x$. Since X is compact, extract a finite subcover U_1, \dots, U_n of $(U_x)_{x \in X}$, and a partition of unity $\lambda_1, \dots, \lambda_n$. Then

$$\|T_x - \sum_{i=1}^n \lambda_i(x) R_{x_i}\| \leq \varepsilon.$$

Therefore, is a finite rank the operator $\sum_i \lambda_i(\cdot) R_{x_i}$ and, \widehat{T} is compact.

Suppose that the range of T is constituted only by Fredholm operators, i.e., \mathcal{F} . In order to see the extension of T to the standard Hilbert $C(X)$ -module, $\widehat{T} \in \mathcal{L}_{C(X)}(\mathcal{H}_{C(X)})$ is a $C(X)$ -Fredholm operator, we must conjure the following:

¹For example, $T : [0, 1] \longrightarrow \mathcal{F}$, $T_x(\xi) := (x\xi_n)_n$.

Summoning 4.0.6 (Classical Bartle-Graves theorem - Corollary 17.67 [1]). Every surjective continuous linear operator between Banach spaces^a admits continuous right inverse, but not necessarily a linear one. 


^aMore generally, completely metrizable locally convex spaces, i.e., Fréchet spaces.

Bartle-Graves theorem has significant improvements and different versions (e.g., [5]), but for our needs, we are fine with the above.

Consider the Calkin algebra given by the quotient of compact operators: $\mathcal{Q} := \mathcal{B}/\mathcal{K}$ and π be the quotient map. Bartle-Graves theorem offers a continuous section $\sigma : \mathcal{Q} \rightarrow \mathcal{B}$ such that $\pi\sigma A = A$. Let S be given by the composition:

$$\begin{array}{ccc} X & \xrightarrow{\quad S \quad} & \mathcal{F} \\ T \downarrow & & \uparrow \sigma \\ \mathcal{F} & \xrightarrow[\pi]{} \text{GL } \mathcal{Q} \xrightarrow[(\cdot)^{-1}]{} & \text{GL } \mathcal{Q} \end{array}$$


Since $T_x \in \mathcal{F}$ if and only if $\pi(T_x)$ is invertible in \mathcal{Q} (by Atkinson's theorem), S is well defined and continuous (each arrow above is), hence defines $\hat{S} : C(X, \ell^2(\mathbb{N})) \rightarrow C(X, \ell^2(\mathbb{N}))$. We are left to show that both $I - \hat{S}\hat{T}$ and $I - \hat{T}\hat{S}$ are $C(X)$ -compact operators in $\mathcal{H}_{C(X)}$.

Since $1_{C(X)} - T \cdot S$ and $1_{C(X)} - S \cdot T$ are continuous families with compact range, \hat{T} is indeed $C(X)$ -Fredholm by (i). 

In the classical Fredholm theory between Hilbert spaces, one only requires that $\ker T$ and $\text{coker } T$ are finite dimensional. Those assumptions are sufficient to guarantee that every classical Fredholm operator has closed range, hence orthogonal decompositions are abundant in the proofs. Only if we could extend it so naturally, maybe we wouldn't be here with a slightly different definition.

Example 4.0.7 (Non closed range A -Fredholm operator). $C[0, 1]$ is a unital C^* -algebra, that we shall consider as a Hilbert C^* -module. Choose T to be


$$\begin{aligned} T : C[0, 1] &\longrightarrow C[0, 1] \\ f &\longmapsto (x \mapsto xf(x)) \end{aligned}$$

Since the algebra is unital, any adjointable operator is $C[0, 1]$ -compact: $\mathcal{L}(C[0, 1]) \simeq \mathcal{K}(C[0, 1])$ (for instance, see Example 3.3.4(ii)), hence must be $C[0, 1]$ -Fredholm as well. Unfortunately, the square root $\sqrt{\cdot}$ doesn't belong to the range of T , but it can be approximated by the Bernstein polynomials $B_n(x) := \sum_{k \leq n} \binom{n}{k} \sqrt{k/n} x^k (1-x)^{n-k}$. 

Therefore, in order to develop the theory of Fredholm operator between Hilbert modules and, in some extent, try to obtain a correspondence with the classical theory, we shall dodge that closure of the Fredholm range. Hence, we will focus on a smaller class of operators: those which admit pseudo-inverse, henceforth, the *regular* ones. Later, we shall extent our results to general Fredholm operators, showing that each and everyone is, in some extent, regularizable.

4.1 Regular Fredholm operators

Definition 4.1.1 (Regular operators). It is said to be *regular* any operator $T \in \mathcal{L}(E, F)$ that admits a *pseudo-inverse*, i.e., there exists $S \in \mathcal{L}(F, E)$ such that $TST = T$ and $STS = S$.

Example 4.1.2. The operators F and G constructed in the proof of Theorem 3.6.3 are regular Fredholm operators whose $FGF = F$ and $GFG = G$. 

For a regular Fredholm operator T , such a pseudo-inverse S fits the Fredholm criteria of T : If S' is such that $I_E - S'T$ and $I_F - TS'$ are finite-rank operators,

$$\begin{aligned} (I_E - S'T)(I_E - ST) &= (I_E - S'T) - (I_E - S'T)ST \\ &= I_E - S'T - ST + \underbrace{S'TST}_{S'T} = I_E - ST \end{aligned}$$

Since $\text{FR}(E, F)$ is an ideal, the above manipulation shows that $I_E - ST$ is indeed a finite-rank operator (and similarly for $I_F - TS$).


Remark 4.1.3. When S is such that TS and ST are idempotents, it is called a *Moore-Penrose* inverse. To motivate the study of regular Fredholm operators as some way to deal with a weaker version of "the range is closed", we exhibit the following theorem:

Theorem. For a Hilbert space H , a bounded operator $T \in \mathcal{B}(H)$ admits a Moore-Penrose pseudo-inverse S if, and only if, $\text{Im } T$ is closed.

One way is disrespectfully trivial: If there exists a Moore-Penrose pseudo-inverse S , $\text{Im } T = \text{Im } TS$. Since TS is a orthogonal projection by hypothesis, $\text{Im } T$ is closed.

Conversely, consider the following decompositions

$$H = \ker T \oplus \text{Im } T^* = \ker T^* \oplus \text{Im } T.$$

Therefore, $T|_{\text{Im } T^*}$ is an injective bounded operator, which posses a bounded inverse S . Similarly, $T^*|_{\text{Im } T}$ contains a bounded inverse R . Those inverses can be extended to all the space, by setting it to zero (which is fine, since the kernels are all there is left in each case). One can verify that $R = S^*$ and that S induces a Moore-Penrose inverse. 

Proposition 4.1.4. Let $T \in \mathcal{L}(E, F)$ be a A -Fredholm operator. If T admits a pseudo-inverse S , then:

- (i) $I_E - ST$ and TS are idempotents with ranges $\ker T$ and $\text{Im } T$.
- (ii) $\ker T$ and $\ker T^*$ are finite rank modules.

Proof. Notice that $(ST)^2 = S(TST) = ST$ and similarly for TS , i.e., they are idempotents. It is easy to see that $I_E - ST$ also has the idempotent badge, $\text{Im}(I_E - ST) = \ker T$ and $\text{Im } TS = \text{Im } T$.

Is easy to see that $I_{\ker T} = (I_E - ST)|_{\ker T}$. When supposing that T is A -Fredholm, let $x, y \in E^n$ be such that $I_E - ST = \Omega_y \Omega_x^*$. Remind that idempotent operators share their range with some projection by the remark 3.5.5. Since $I_E - ST$ is an idempotent, there exists a self-adjoint idempotent operator P such that $\text{Im}(I_E - ST) = \text{Im } P = \ker T$. Therefore, with $a = \Omega_y \Omega_x^*$ and $p = P$, 3.5.5.(vii) guarantee that

$$\Omega_y \Omega_x^*|_{\ker T} = P \Omega_y \Omega_x^* P \stackrel{P^* = P}{=} \Omega_{Py} \Omega_{Px}^*.$$

Since $Py, Px \in (\ker T)^n$, it follows that $I_{\ker T} = \Omega_{Py} \Omega_{Px}^*$ is a finite-rank operator over $\ker T$, i.e., $\ker T$ is a finite-rank module. Very much the same is sufficient to obtain that $\ker T^*$ also is a finite-rank module. \square

The rank of a finite rank module is well defined as seen before. Hence, the above proposition enable us to define the Index of regular Fredholm operators.

Definition 4.1.5. If T is a A -Fredholm operator who admits a pseudo-inverse (i.e., regular), set their *index* to be the $K_0(A)$ element given by

$$\text{ind } T := \text{rank}(\ker T) - \text{rank}(\ker T^*).$$

Proposition 4.1.6. Our elementary propositions of classical Fredholm operators stands nicely. If $T \in \mathcal{L}(E, F)$ is a regular Fredholm operator, then:

- (i) $\text{ind } T^* = -\text{ind } T$.
- (ii) For any pseudo-inverse S , $\text{rank}(\ker T^*) = \text{rank}(\ker S)$ and $\text{ind } S = -\text{ind } T$.
- (iii) If there are invertible operators U and V between Hilbert modules such that:

$$\begin{array}{ccccccc} & & & & VTU & & \\ & & & \curvearrowright & & & \\ X & \xrightarrow[\cong]{U} & E & \xrightarrow{T} & F & \xrightarrow[\cong]{V} & Y \end{array}$$

Therefore VTU is Fredholm and $\text{ind}(VTU) = \text{ind } T$.

- (iv) If $T_i \in \mathcal{L}(E_i, F_i)$ is a regular Fredholm operator for $i \in \{1, 2\}$, the direct sum $T_1 \oplus T_2$ is also regular Fredholm and $\text{ind}(T_1 \oplus T_2) = \text{ind } T_1 + \text{ind } T_2$.

Proof.

- (i) Clear.
- (ii) Since S and T^* are Fredholm operators, $\ker T^*$ and $\ker S$ are finite rank modules (4.1.4). In what comes next, keep in mind that $(\text{Im } T)^\perp = \ker T^*$. Visiting again the remark 3.5.5, one can conclude that for any idempotent $Q : F \rightarrow E$, $F = \text{Im } Q \oplus (\text{Im } Q)^\perp$. Since TS is an idempotent, we obtain the following diagram of equality's:

$$\begin{array}{ccc} \text{Im}(I_F - TS) \oplus \text{Im } TS & \xlongequal{\quad} & F \xrightarrow{\text{3.5.5}} (\text{Im } TS)^\perp \oplus \text{Im } TS \\ \parallel & & \parallel \\ \ker S \oplus \text{Im } TS & & \ker T^* \oplus \text{Im } TS \end{array}$$

$(\text{Im } TS)^\perp = (\text{Im } T)^\perp = \ker T^*$

Therefore, $\ker S$ and $\ker T^*$ are quasi-stably-isomorphic. Therefore, 3.6.3 guarantee that $\text{rank}(\ker S) = \text{rank}(\ker T^*)$. Consequentially, $\text{ind } S = -\text{ind } T$.

- (iii) Notice that $\ker VT = \ker T$ since V is an invertible one, hence $\text{rank}(\ker VT) = \text{rank}(\ker T)$. Analysing $U|_{\ker TU}$, one obtains that $\ker TU \simeq \ker T$, thus $\text{rank}(\ker TU) = \text{rank}(\ker T)$. The exact same roll goes for the adjoints. Therefore, the indexes coincide.

- (iv) It is the case that $\Omega_{\xi_1 \oplus \xi_2} = \Omega_{\xi_1} \oplus \Omega_{\xi_2}$ for any $\xi_1 \oplus \xi_2 \in E_1 \oplus E_2$, which is sufficient to infer that $T_1 \oplus T_2$ is a Fredholm operator.

Since $\ker(T_1 \oplus T_2) = \ker T_1 \oplus \ker T_2$, $\ker(T_1 \oplus T_2)$ is a finite rank module. If $\text{rank } T_i = [p_i]_0$, it is clear that

$$\begin{aligned} \text{rank}(\ker(T_1 \oplus T_2)) &= [\text{diag}(p_1, p_2)]_0 \\ &= [p_1]_0 + [p_2]_0 \\ &= \text{rank}(\ker T_1) + \text{rank}(\ker T_2). \end{aligned}$$

Therefore, the desired index relation follows. \square

Since our compact operators aren't necessarily the same as in Hilbert space case, the index invariance under compact perturbations needs to be handed carefully.

Proposition 4.1.7. If $T \in \mathcal{L}(E)$ is a regular Fredholm operator such that $(I - T) \in \mathcal{K}(E)$, then $\text{ind } T = 0$.

Proof. Let S be a pseudo-inverse of T . Since $I - T$ is a compact operator and the compact operators is an ideal, notice that S is a compact perturbation of the identity:

$$S = I + S(I - T) - (I - ST)$$

Considering the isomorphism map $U : \ker T \oplus \operatorname{Im} S \longrightarrow \ker S \oplus \operatorname{Im} S$ given by

$$U := \begin{pmatrix} I - TS & I - TS \\ S & S \end{pmatrix} \quad U^{-1} = \begin{pmatrix} I - ST & (I - ST)T \\ ST & STT \end{pmatrix}$$

with the fact that $I - S = I_{\operatorname{Im} S} - U_{\operatorname{Im} S \operatorname{Im} S}$ is compact, the modules $\ker T$ and $\ker S$ are quasi-stably-isomorphic. By 3.6.3, $\operatorname{rank}(\ker T) = \operatorname{rank}(\ker S)$, hence

$$\begin{aligned} \operatorname{ind} T &= \operatorname{rank}(\ker T) - \operatorname{rank}(\ker T^*) \\ &\stackrel{4.1.6(ii)}{=} \operatorname{rank}(\ker T) - \operatorname{rank}(\ker S) = 0. \quad \square \end{aligned}$$

Theorem 4.1.8. If $T_1, T_2 \in \mathcal{L}(E, F)$ are regular Fredholm operators such that $T_1 - T_2$ is compact, then $\operatorname{ind} T_1 = \operatorname{ind} T_2$.

Proof. The action plan for the proof will be as follows: Build accessory operators U and R in function of the given maps, such that U is invertible and $\operatorname{ind} R = \operatorname{ind} T_2 - \operatorname{ind} T_1$. Hence, $\operatorname{ind}(UR)$ will be a compact perturbation of the identity, so we can use the previous theorem and obtain that $\operatorname{ind} R = \operatorname{ind}(UR) = 0$.

Let S_1 and S_2 be pseudo inverses for T_1 and T_2 . Define operators U and R in $\mathcal{L}(E \oplus F)$ by

$$U := \begin{pmatrix} I_E - S_1 T_1 & S_1 \\ T_1 & I_F - T_1 S_1 \end{pmatrix} \quad \text{and} \quad R := \begin{pmatrix} 0 & S_1 \\ T_2 & 0 \end{pmatrix}.$$

(i) $\operatorname{ind} R = \operatorname{ind} T_2 - \operatorname{ind} T_1$: Using the coordinate switch operator (which has index zero since it is a invertible one), one obtains that:

$$\begin{array}{ccc} E \oplus F & \xrightarrow{R} & E \oplus F \\ \searrow (x,y) \mapsto (y,x) & & \nearrow S_1 \oplus T_2 \\ & F \oplus E & \end{array} \quad \begin{aligned} \operatorname{ind} R &\stackrel{4.1.6(iii)}{=} \operatorname{ind}(S_1 \oplus T_2) \\ &\stackrel{4.1.6(iv)}{=} \operatorname{ind} S_1 + \operatorname{ind} T_2 \\ &\stackrel{4.1.6(ii)}{=} -\operatorname{ind} T_1 + \operatorname{ind} T_2. \end{aligned}$$

(ii) **U is invertible:** We'll show even more: it is a order 2 nilpotent element. Since $I_E - S_1 T_1$ and $I_F - T_1 S_1$ are idempotents,

$$\begin{aligned} U^2 &= \begin{pmatrix} I_E - S_1 T_1 & S_1 \\ T_1 & I_F - T_1 S_1 \end{pmatrix}^2 \\ &= \begin{pmatrix} (I_E - S_1 T_1)^2 + S_1 T_1 & (I_E - S_1 T_1)S_1 + S_1(I_F - T_1 S_1) \\ T_1(I_E - S_1 T_1) + (I_F - T_1 S_1)T_1 & T_1 S_1 + (I_F - T_1 S_1)^2 \end{pmatrix} \\ &= \begin{pmatrix} I_E & 0 \\ 0 & I_F \end{pmatrix} = I_{E \oplus F}. \end{aligned}$$

Therefore U is invertible.

(iii) **UR is a compact perturbation of identity:** First, we obtain UR :

$$UR = \begin{pmatrix} I_E - S_1 T_1 & S_1 \\ T_1 & I_F - T_1 S_1 \end{pmatrix} \begin{pmatrix} 0 & S_1 \\ T_2 & 0 \end{pmatrix} = \begin{pmatrix} S_1 T_2 & 0 \\ (I_F - T_1 S_1) T_2 & T_1 S_1 \end{pmatrix}$$

Bravely evaluating the difference, we must determine if is compact the following operator:

$$I_{E \oplus F} - UR = \begin{pmatrix} I_E - S_1 T_2 & 0 \\ (T_1 S_1 - I_F) T_2 & I_F - T_1 S_1 \end{pmatrix}$$

Notice that all operators in the second row are compact since T_1 is Fredholm. From the hypothesis, $T_1 - T_2$ is compact, hence:

$$\begin{aligned} I_E - S_1 T_2 &= I_E - S_1 T_1 + S_1 T_1 - S_1 T_2 \\ &= (I_E - S_1 T_1) + S_1 (T_1 - T_2) \in \mathcal{K}(E). \end{aligned}$$

Since each entry of $I_{E \oplus F} - UR$ is a compact operator, the claim is proved.

Using 4.1.6(iii) again, we have that $\text{ind}(UR) = \text{ind } R$. Since UR is a compact perturbation of the identity, it follows that $\text{ind}(UR) = 0$ by Proposition 4.1.7. \square

4.2 Regularization of Fredholm operators

Time to extend our concepts to general Fredholm operators. A change in algebras will be necessary, so we write our next lemma with new a C^* -algebra notation.

Lemma 4.2.1. Let B be a unital C^* -algebra and $T \in \mathcal{L}_B(E, F)$ a B -Fredholm but not necessarily regular. There exists a natural n and some $x \in E^n$ such that

$$\begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} : E \oplus B^n \longrightarrow F \oplus B^n$$

is a regular B -Fredholm operator.

Proof. Let S be a pseudo-inverse of T such that both $I_E - ST$ and $I_F - TS$ are finite rank operators, and $I_E - ST = \Omega_y \Omega_x^*$ for some $y \in F^n$, $x \in E^n$. We will construct operators \tilde{T} and \tilde{S} that are regular Fredholm. Define the following operators:

$$\tilde{T} := \begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} \quad \text{and} \quad \tilde{S} := \begin{pmatrix} S & \Omega_y \\ 0 & 0 \end{pmatrix}.$$

(i) **\tilde{T} and \tilde{S} are pseudo-inverses of each other, hence regular:** In what follows, we need the expressions:

- (a) $T\Omega_y\Omega_x^* = T(I_E - ST) = T - TST = 0.$
- (b) $\Omega_x^*(ST + \Omega_y\Omega_x^*) = \Omega_x^*(ST + I_E - ST) = \Omega_x^*.$

Notice that

$$\begin{aligned}
 \tilde{T}\tilde{S}\tilde{T} &= \begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} \begin{pmatrix} S & \Omega_y \\ 0 & 0 \end{pmatrix} \begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} \\
 &= \begin{pmatrix} TS & T\Omega_y \\ \Omega_x^*S & \Omega_x^*\Omega_y \end{pmatrix} \begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} \\
 &= \begin{pmatrix} TST + T\Omega_y\Omega_x^* & 0 \\ \Omega_x^*(ST + \Omega_y\Omega_x^*) & 0 \end{pmatrix} \stackrel{(a)+(b)}{=} \begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} = \tilde{T}
 \end{aligned}$$

Similarly, one can obtain that $\tilde{S}\tilde{T}\tilde{S} = \tilde{S}$, hence \tilde{T} and \tilde{S} are regular due to the fact that they are each others pseudo-inverses.

(ii) **\tilde{T} and \tilde{S} are Fredholm operators:** Notice that:

$$\begin{aligned}
 (4.1) \quad I_{E \oplus B^n} - \tilde{S}\tilde{T} &= \begin{pmatrix} I_E & 0 \\ 0 & I_{B^n} \end{pmatrix} - \begin{pmatrix} ST + \Omega_y\Omega_x^* & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & I_{B^n} \end{pmatrix} \\
 I_{F \oplus B^n} - \tilde{T}\tilde{S} &= \begin{pmatrix} I_F & 0 \\ 0 & I_{B^n} \end{pmatrix} - \begin{pmatrix} TS & T\Omega_y \\ \Omega_x^*S & \Omega_x^*\Omega_y \end{pmatrix} = \begin{pmatrix} I_F - TS & -T\Omega_y \\ -\Omega_x^*S & I_{B^n} - \Omega_x^*\Omega_y \end{pmatrix}
 \end{aligned}$$

Lets check that every entry in those matrices are compact:

- (a) **I_E is finite rank:** Since B is unital, $I_{B^n} = \Omega_{(1_E, \dots, 1_E)}\Omega_{(1_E, \dots, 1_E)}^*$.
- (b) **$I_F - TS$ is finite rank:** By assumption.
- (c) **$-\Omega_x^*S, -T\Omega_y$ and $I_{B^n} - \Omega_x^*\Omega_y$ are compact:** This is due to the fact that Ω_y and Ω_x^* are compact (proposition 3.3.5) and the set of compact operators is an ideal.

Therefore, $I_{E \oplus B^n} - \tilde{S}\tilde{T}$ and $I_{F \oplus B^n} - \tilde{T}\tilde{S}$ are compact operators. Finally, proposition 4.0.2 guarantee that both \tilde{T} and \tilde{S} are regular Fredholm operators. \square

Definition 4.2.2 (Regularization of a Fredholm operator). Given a A -Fredholm $T \in \mathcal{L}_A(E, F)$, the *regularization of T* is the \tilde{A} -Fredholm $\tilde{T} \in \mathcal{L}_{\tilde{A}}(E \oplus \tilde{A}^n, F \oplus \tilde{A}^n)$ constructed using lemma 4.2.1 for $B = \tilde{A} := A \oplus \mathbb{C}$ being the unitization of A .

Proposition 4.2.3. For any A -Fredholm operator T , despite the fact that the regularization \tilde{T} is a \tilde{A} -Fredholm operator, the index of \tilde{T} lies in $K_0(A)$.

Proof. Let $\varepsilon : \tilde{A} \longrightarrow \mathbb{C}$ the complex projection. Since $K_0(A)$ is the kernel of ε_0 , we seek to obtain that $\varepsilon_0(\text{ind } \tilde{T}) = 0$. We borrow notations and results from the proof of 4.2.1, i.e., $I_E - ST = \Omega_y\Omega_x^*$ and

$$\tilde{T} := \begin{pmatrix} T & 0 \\ \Omega_x^* & 0 \end{pmatrix} \quad \text{and} \quad \tilde{S} := \begin{pmatrix} S & \Omega_y \\ 0 & 0 \end{pmatrix}$$

are regular \tilde{A} -Fredholm operators and pseudo-inverses of each other. To compute the index of \tilde{T} , first we obtain that $\mathbf{rank}(\ker \tilde{T}) = n \cdot 1_{K_0(A)}$; in order to obtain $\mathbf{rank}(\ker \tilde{T}^*) = \mathbf{rank}(\ker \tilde{S})$, we will introduce two new operators P and Q , such that the rank of $\ker \tilde{S}$ will coincide with the embedding of trace of $\varepsilon(Q)$, which will be equal to n .

Now, we look to verify those claims:

- (i) $\mathbf{rank}(\ker \tilde{T}) = n \cdot 1_{K_0(A)}$: In the proof of Lemma 4.2.1 (4.1), we saw that $I_{E \oplus \tilde{A}^n} - \tilde{S}\tilde{T} = 0 \oplus I_{\tilde{A}^n}$. Since $\ker \tilde{T} = \text{Im}(I_{E \oplus \tilde{A}^n} - \tilde{S}\tilde{T})$, it follows that

$$\mathbf{rank}(\ker \tilde{T}) = [0 \oplus I_{\tilde{A}^n}]_0 = n \cdot [1_{\tilde{A}}]_0 = n \cdot 1_{K_0(A)}.$$

For notation sake, let:

$$(4.2) \quad P := I_{F \oplus \tilde{A}^n} - \tilde{T}\tilde{S} \stackrel{(4.1)}{=} \begin{pmatrix} I_F - TS & -T\Omega_y \\ -\Omega_x^*S & I_{B^n} - \Omega_x^*\Omega_y \end{pmatrix}$$

- (ii) $\ker \tilde{S} = \text{Im } P$. Lets check that the two sets coincide: In one direction, $\tilde{S} - \tilde{S}\tilde{T}\tilde{S} = 0$ since \tilde{S} and \tilde{T} are pseudo-inverses of each other. Hence $\ker \tilde{S} \supset \text{Im } P$. Conversely, the elements of the range of P can be written as:

$$\begin{aligned} P(\zeta + a) &= \begin{pmatrix} I_F - TS & -T\Omega_y \\ -\Omega_x^*S & I_{B^n} - \Omega_x^*\Omega_y \end{pmatrix} \begin{pmatrix} \zeta \\ a \end{pmatrix} \\ &= \begin{pmatrix} \zeta - T(S\zeta + \Omega_y a) \\ a - \Omega_x^*(\Omega_y a + S\zeta) \end{pmatrix} \end{aligned}$$

whenever $\zeta \in F$ and $a \in \tilde{A}^n$. If $(\zeta + a) \in \ker \tilde{S}$, then $P(\zeta + a) = \zeta + a$, hence $\zeta \oplus a$ is in the range of P , i.e., $\ker \tilde{S} \subset \text{Im } P$.

Hence, we shall compute $\mathbf{rank}(\text{Im } P)$. Since \tilde{S} is a regular Fredholm operator, $\text{Im } P = \ker \tilde{S}$ is a finite-rank module (4.1.4), i.e., $I_{\text{Im } P}$ can be written as $\Omega_\phi \Omega_\psi^*$ for some $m \in \mathbb{N}$ and a pair of tuples $\phi, \psi \in (F \oplus B^n)^m$, hence

$$I_{\text{Im } P} = \Omega_\phi \Omega_\psi^* \Rightarrow P = \Omega_\phi \Omega_\psi^* P$$

Replacing if necessary each coordinate ϕ_i with $P\phi_i$ if necessary, we can assume that $P\Omega_\phi = \Omega_\phi$. This will lead us to the next claim:

- (iii) $Q := \Omega_\psi^* \Omega_\phi \in \mathcal{L}(\tilde{A}^n)$ is an idempotent operator: Indeed:

$$Q^2 \stackrel{P\Omega_\phi = \Omega_\phi}{=} (\Omega_\psi^* P \Omega_\phi)^2 = \Omega_\psi^* P \underbrace{(\Omega_\phi \Omega_\psi^*)}_P P \Omega_\phi = \Omega_\psi^* \Omega_\phi = Q.$$

Therefore, Q is an idempotent operator in $\mathcal{L}(\tilde{A}^n)$ which corresponds to left multiplication by the matrix $(\langle \phi_i, \psi_j \rangle)_{i,j}$, and $\text{Im } Q \simeq \text{Im } P$ as \tilde{A} -modules.

(iv) $\text{Tr } \varepsilon(Q) = n$: Let $(e_r)_r$ be the canonical basis of \tilde{A}^n . We shall write the coordinates of ϕ and ψ as:

$$\phi_i = \zeta_i + a_i \quad \text{and} \quad \psi_i = \xi_i + b_i$$

for $\zeta_i, \xi_i \in F$ and $a_i, b_i \in \tilde{A}^n$. Hence $\varepsilon(\langle \psi_i, \phi_i \rangle) = \varepsilon(\langle b_i, a_i \rangle)$ which enables us to expand in the following way:

$$\begin{aligned} \text{Tr } \varepsilon(Q) &= \sum_{i=1}^m \varepsilon(\langle \psi_i, \phi_i \rangle) \\ &= \sum_{i=1}^m \varepsilon(\langle b_i, a_i \rangle) \\ &= \varepsilon\left(\sum_{i=1}^m \sum_{r=1}^n \langle b_i, e_r \rangle \langle e_r, a_i \rangle\right) \\ &= \varepsilon\left(\sum_{i=1}^m \sum_{r=1}^n \langle e_r, a_i \rangle \langle b_i, e_r \rangle\right) \\ &= \varepsilon\left(\sum_{r=1}^n \langle (0, e_r), P(0, e_r) \rangle\right) \end{aligned}$$

Using the definition of P in (4.2), the term $\sum_{r=1}^n \langle (0, e_r), P(0, e_r) \rangle$ can be expressed as

$$\sum_{r=1}^n \langle e_r, (I_{B^n} - \Omega_x^* \Omega_y) e_r \rangle = n \cdot 1_A - \sum_{r=1}^n \langle x_r, y_r \rangle$$

hence $\text{Tr } \varepsilon(Q) = n$.

With all these steps, we conclude that

$$\text{rank}(\ker \tilde{S}) = \text{rank}(\text{Im } P) = \text{rank}(\text{Im } Q) = \text{Tr } \varepsilon(Q) \cdot 1_{K_0(A)} = n \cdot 1_{K_0(A)}$$

and finally that $\varepsilon_0(\text{ind } \tilde{T}) = 0$. \square

The statement of 4.2.3 is meant to refer to the specific construction of \tilde{T} obtained in 4.2.2. But note that any regular Fredholm operator in $\mathcal{L}_{A^u}(E \oplus \tilde{A}^n, F \oplus \tilde{A}^n)$, which has T in the upper left corner, will differ from the \tilde{T} above, by a A -compact operator. Therefore its index will coincide with that of \tilde{T} by 4.2.1, and so will be in $K_0(A)$ as well.

Definition 4.2.4. If T is a Fredholm operator in $\mathcal{L}(E, F)$, then the Fredholm index of T , denoted $\text{ind } T$, is defined to be the index of the regular Fredholm operator \tilde{T} constructed in proposition 4.2.3.

It is clear that all properties listed in 4.1.6 are naturally extended to general Fredholm operators;

As consequence of the Atkinson's theorem in the classical theory, one can obtain that the original index is locally constant. Since it is now our definition, we can extract the same proof.

Proposition 4.2.5. Let $(E, F)Q := \mathcal{L}(E, F)/\mathcal{K}(E, F)$ be the Calkin algebra and $\pi : \mathcal{L}(E, F) \longrightarrow \mathcal{Q}(E, F)$ be the quotient projection. The set of Fredholm operators $\mathcal{F}(E, F) := \pi^{-1} \text{GL } \mathcal{Q}(E, F) \subset \mathcal{L}(E, F)$ is an open subset and $\text{ind} : \mathcal{F}(E, F) \longrightarrow K_0(A)$ is locally constant.

Proof. The fact that $\mathcal{F}(E, F)$ is an open set follows from the continuity of π on the the invertible elements of a unital C^* -algebra. To check the continuity of the index, let T be a Fredholm operator and S be one of its pseudo-inverses. If R is a Fredholm operator in the open ball around T of radius $\|S\|^{-1}$,

$$\|TS - RS\| \leq \|T - R\|\|S\| \leq 1.$$

Hence $I - (TS - RS)$ is a invertible Fredholm operator, which means that it has index 0. Notice that $(I - (TS - RS))T = RST$. Therefore:

$$\begin{aligned} \text{ind } T &= \text{ind } ((I - (TS - RS))T) \\ &= \text{ind } (RST) = \text{ind } R + \text{ind } S + \text{ind } T \end{aligned}$$

hence $\text{ind } R = -\text{ind } S$. Since R was an arbitrary element of the open ball, it follows necessarily that $\text{ind} \upharpoonright_{B(T, \|S\|^{-1})}(R) = -\text{ind } S$, i.e., the index is locally constant. \square

Proposition 4.2.6. Choose A -Fredholm operators such that

$$\begin{array}{ccccc} & & T_2 T_1 & & \\ & \curvearrowright & & \curvearrowleft & \\ E & \xrightarrow{T_1} & F & \xrightarrow{T_2} & G \end{array}$$

Therefore, $T_2 T_1$ is a Fredholm operator and $\text{ind } (T_2 T_1) = \text{ind } T_1 + \text{ind } T_2$.

Proof. Assume beforehand that $E = F = \mathcal{H}_A$ and consider $H_t : \mathcal{H}_A \oplus \mathcal{H}_A \longrightarrow \mathcal{H}_A \oplus \mathcal{H}_A$ be a continuous path of Fredholm operators given by

$$H_t := \begin{pmatrix} T_1 & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} \cos(t) & -\sin(t) \\ \sin(t) & \cos(t) \end{pmatrix} \begin{pmatrix} I & 0 \\ 0 & T_2 \end{pmatrix} \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$$

for $t \in [0, \pi/2]$, which connects $T_1 \oplus T_2$ to $T_2 T_1 \oplus I$. Therefore

$$\text{ind } (T_1 T_2) = \text{ind } (H_{\pi/2}) = \text{ind } (H_0) = \text{ind } (T_1 \oplus T_2) = \text{ind } T_2 + \text{ind } T_1.$$

The general case follows by using the Kasparov stabilization theorem 3.4.6. \square


Remark 4.2.7. In light of example 4.0.5, let $T : X \longrightarrow \mathcal{F}(\ell^2(\mathbb{N}))$ be a continuous family of classical Fredholm operators. We saw that $\hat{T} : \xi \longmapsto T_{(\cdot)} \xi(\cdot)$ for $\xi \in \mathcal{H}_{C(X)}$ was a $C(X)$ -Fredholm operator.

However, since $\ell^2(\mathbb{N})$ is a separable Hilbert space, one can consider an orthogonal basis $(e_n)_n \subset \ell^2(\mathbb{N})$ and let H_n be the closed subspace spanned by e_0, e_1, \dots, e_{n-1} , and let p_n be the associated projection. Let $\text{Vect}(X)$

denote the abelian semi-group of isomorphism classes of all (complex) vector bundles over X .

Using topological K -theory, one can define a analytical index given by

$$\overline{\text{ind}} T := [\ker p_n \circ T]_0 - [X \times H_n^\perp]_0$$

for a sufficiently large n , where $[\cdot]_0$ is the isomorphism class in $\text{Vect}(X) \hookrightarrow K(X)$. 

Proposition 4.2.8. For any $\alpha \in K_0(A)$, there exists a Fredholm operator T with $\text{ind}(T) = \alpha$.

Proof. Write $\alpha = [p]_0 - [q]_0$ with $\varepsilon_0([p]_0 - [q]_0) = 0$, for self-adjoint idempotent matrices. Hence $\varepsilon(p)$ and $\varepsilon(q)$ are similar matrix. After performing a conjugation of, say q , by a complex unitary matrix, we may assume that $\varepsilon(p)$ and $\varepsilon(q)$ are in fact equal, hence $(p - q) \in \mathbb{M}_n(A)$. With bricks in hands, choose

$$\begin{aligned} T : pA^n &\longrightarrow qA^n \\ x &\longmapsto qx \end{aligned}$$

which is

- (i) **T is a Fredholm operator:** Let $S : qA^n \longrightarrow pA^n$ be the similar operator given by $Sy = py$. Let $(u_\lambda)_\lambda$ be an approximate identity for A . Therefore, consider the tuples ξ and η^λ , where their coordinates are given by:

$$\xi_i = p(p - q)p_i \quad \text{and} \quad \eta_i^\lambda = p_i u_\lambda \quad (1 \leq i \leq n)$$

With the tuples defined, remember that $\langle a, b \rangle_A = a^*b$, hence:

$$\begin{aligned} \Omega_\xi \Omega_{\eta^\lambda}^* x &= \sum_{i=1}^n \xi_i \langle \eta_i^\lambda, x \rangle_A \\ &= \sum_{i=1}^n \xi_i ((p_i u_\lambda)^* x) \\ &= \sum_{i=1}^n p(p - q)p_i u_\lambda p_i^* x \end{aligned}$$

for each $x \in pA^n$. Therefore, the following converges uniformly:

$$\begin{aligned} \lim_{\lambda} \Omega_\xi \Omega_{\eta^\lambda}^* x &= \sum_{i=1}^n p(p - q)p_i p_i^* x \\ &= p(p - q)px \\ &= x - pqx = (I - ST)x \end{aligned} \quad (\|x\| \leq 1)$$

The above shows that $I_{pA^n} - ST$ is compact, and the same conclusions can be drawn for $I_{qA^n} - TS$ also. By applying 4.0.2, the claim is proved.

(ii) **ind** $T = \alpha$: In order to compute the index of T , consider the operators

$$T' = \begin{pmatrix} qp & q(I-p) \\ (I-q)p & (I-q)(I-p) \end{pmatrix} \quad \text{and} \quad S' = \begin{pmatrix} pq & p(I-q) \\ (I-p)q & (I-p)(I-q) \end{pmatrix}$$

Direct computation shows that

$$S'T' = \begin{pmatrix} I_{pA^n} & 0 \\ 0 & I_{\tilde{A}^n - p} \end{pmatrix} \quad \text{and} \quad T'S' = \begin{pmatrix} I_{qA^n} & 0 \\ 0 & I_{\tilde{A}^n - q} \end{pmatrix},$$

from which it follows that S' is a pseudo-inverse for T' and hence that T' is a regular A^u -Fredholm operator.

By construction, $I - S'T'$ and $I - T'S'$ are compact self-adjoint idempotents, hence their ranges are finite rank modules (3.5.4). Moreover, we already know that

$$\text{Im}(I - S'T') = \ker T' \quad \text{and} \quad \text{Im}(I - T'S') = \ker S'$$

which turns possible the index calculation:

$$\begin{aligned} \text{ind } T &= \text{ind } T' \\ &= \text{rank}(\ker T') - \text{rank}(\ker S') \\ &= \text{rank} \text{Im}(I - S'T') - \text{rank} \text{Im}(I - T'S') \\ &= [p]_0 - [q]_0 = \alpha \end{aligned}$$

as desired. □

4.3 Fredholm Picture of $K_0(A)$

We already saw that every element of $K_0(A)$ is the index of some Fredholm operator.

In order to avoid pissing any set-theoretical reader, choose ω to be one of your favorite cardinal numbers, as long as it is greater than the cardinality of each and every A^n for every integer n . Denote by $F_0(A)$ the family of all A -Fredholm operators whose domain and codomain are Hilbert modules with cardinality no larger than ω . We will introduce an equivalence relation in $F_0(A)$, by characterizing whenever two operators T_1 and T_2 contains the same index.

Notice that whenever $T \oplus I_{A^n}$ is a compact perturbation of an invertible operator, one can conclude by 4.1.8 that **ind** $T = 0$. This is a good indicator for an equivalence relation:

Proposition 4.3.1. Let $T \in \mathcal{L}(E, F)$ be Fredholm operator with **ind** $T = 0$. Therefore, there exists some integer n such that $T \oplus I_{A^n}$ is a compact perturbation of an invertible operator.

Proof. Let \tilde{T} be the regularization of T and, as always, there is an integer n and an operator S such that $I - \tilde{S}\tilde{T} = 0 \oplus I_{\tilde{A}^n}$ described in the proof of Lemma 4.2.1 (4.1).

- (i) **There exists n such that $\text{Im}(I - \tilde{T}\tilde{S}) \cong \tilde{A}^n$:** The hypothesis of null index is equivalent to $\text{rank Im}(I - \tilde{T}\tilde{S}) = \text{rank Im}(I - \tilde{T}\tilde{S})$, hence

$$\text{rank Im}(I - \tilde{T}\tilde{S}) = \text{rank Im}(0 \oplus I_{\tilde{A}^n}) = n \cdot 1_{K_0(A)}$$

i.e., $\text{Im}(I - \tilde{T}\tilde{S})$ is stably isomorphic to \tilde{A}^n as \tilde{A} -modules, meaning that for some integer r , $\text{Im}(I - \tilde{T}\tilde{S}) \oplus \tilde{A}^r \simeq \tilde{A}^{n+r}$. Remember that $\Omega_x^* = (\langle x_i, \cdot \rangle)_{i \leq n}$, hence, for $0 \in E^m$, $\Omega_{(x,0)}^*(\cdot) = (\Omega_x^*(\cdot), 0)$ and the regularization \tilde{T} can be updated to

$$\tilde{T} = \begin{pmatrix} (T, 0) & 0 \\ \Omega_{(x,0)}^* & 0 \end{pmatrix}$$

As seen, n can be increased without essentially changing \tilde{T} . Therefore, there is no danger in assuming that $\text{Im}(I - \tilde{T}\tilde{S}) \simeq \tilde{A}^n$.

- (ii) **There is a orthonormal generating set $((\zeta_i + a_i))_{i \leq n} \subset \text{Im}(I - \tilde{T}\tilde{S})$ such that $\varepsilon((a_1, \dots, a_n)) = I_{\mathbb{M}_n(\tilde{A})}$:** Let $(p_i)_{i \leq n} \subset \text{Im}(I - \tilde{T}\tilde{S})$ with $p_i = (\zeta_i + a_i) \in F \oplus \tilde{A}^n$. The elements p_i can be chosen so that they generate the module and $\langle p_i, p_j \rangle = \delta_{i,j}$, i.e., they are orthonormal. Since each $a_i \in \tilde{A}^n$, one can write $a_i = (a_{i,r})_{r \leq n}$. Hence, orthonormality can be written as:

$$\begin{aligned} \delta_{i,j} &= \langle p_i, p_j \rangle_{F \oplus \tilde{A}^n} \\ &= \langle \zeta_i, \zeta_j \rangle_F + \langle a_i, a_j \rangle_{\tilde{A}^n} \\ &= \langle \zeta_i, \zeta_j \rangle_F + \sum_{r=1}^n \langle a_{i,r}, a_{j,r} \rangle_{\tilde{A}} \\ &= \langle \zeta_i, \zeta_j \rangle_F + \sum_{r=1}^n a_{i,r}^* a_{j,r} \end{aligned}$$

The projected matrix $u := \varepsilon((a_{i,r})_{i,r})$ is unitary, i.e., $uu^* = u^*u = I_{\mathbb{M}_n(\tilde{A})}$. Whence, setting $q_i := \sum_j u_{ij}^* p_j$, we obtain $q_i = \xi_i + b_i$ in which $\varepsilon(b_{i,j}) = \delta_{i,j}$, i.e., $\varepsilon((b_1, \dots, b_n)) = I_{\mathbb{M}_n(\tilde{A})}$. At the end of the day, one can suppose that $(p_i)_{i \leq n}$ attends the required condition, otherwise, replace p by q .

With this simplifications, we have in hands the following isomorphism:

$$\begin{aligned} U : E \oplus \tilde{A}^n &\longrightarrow F \oplus \tilde{A}^n \\ \xi + b &\longmapsto \begin{pmatrix} T & \Omega_\zeta \\ \Omega_x^* & \Omega_a \end{pmatrix} \begin{pmatrix} \xi \\ b \end{pmatrix} \end{aligned}$$

The complex component can be forget, so that $(E \oplus \tilde{A}^n) \cdot A = A^n$ for any Hilbert A -module E . Thus, U can be restricted to an element in $\text{GL } \mathcal{L}(E \oplus A^n, F \oplus A^n)$. With this in mind, notice that the difference operator

$$U - T \oplus I_{A^n} = \begin{pmatrix} 0 & \Omega_\zeta \\ \Omega_x^* & ((a_{ij} - \delta_{ij}))_{i,j} \end{pmatrix}$$

is compact, since the right lower entry $((a_{ij} - \delta_{ij}))_{i,j}$ is compact. But this matrix was seen to be in $\mathbb{M}_n(A)$, since its projection by ε is zero. \square

As consequence, proposition 4.3.1 immediately characterizes whenever two Fredholm operators between different Hilbert modules have the same index.

Corollary 4.3.2. Whenever two Fredholm operators $T_i \in \mathcal{L}(E_i, F_i)$ ($i \in \{1, 2\}$) share the same index $\text{ind } T_1 = \text{ind } T_2$, there exists a integer n such that

$$T_1 \oplus T_2^* \oplus I_{A^n} : E_1 \oplus F_2 \oplus A^n \longrightarrow E_2 \oplus F_1 \oplus A^n$$

is a A -compact perturbation of an invertible operator.

Declare two operators in $T_1, T_2 \in F_0(A)$ to be equivalent whenever $T_1 \oplus T_2^* \oplus I_{A^n}$ is a compact perturbation of an invertible operator, i.e., $\text{ind } T_1 = \text{ind } T_2$. Denote $F(A)$ the be the induced set of equivalence classes, which is an abelian group when equipped with the direct sum operation \oplus , where $(\cdot)^{-1} : T \longmapsto T^*$.

Considering the index map between $F(A)$ and $K_0(A)$ is the most natural think up to this point, since 4.2.8 already shows that it is a surjective map, and the equivalence relation ensures the injectivity. More over, since $[\text{diag}(x, y)]_0 = [x]_0 + [y]_0$ in $K_0(A)$, we had produced the following Atiyah-Jänich analogue:

Corollary 4.3.3. The index map

$$\text{ind} : F(A) \longrightarrow K_0(A)$$

is a group isomorphism.

Not only analogue, equivalent: Let $K^0(X)$ be the K -group of a compact Hausdorff space $X \in \mathbf{CHaus}$, in the realm of topological K -theory.

Chapter 5

Application to Morita Equivalence

5.1 Preliminars on Hilbert C^* -bimodules

Let A and B be two C^* -algebras. A Hilbert (A, B) -bimodule X is a space with two inner products:

$$(\cdot \mid \cdot) : X \times X \longrightarrow A \quad \text{and} \quad \langle \cdot, \cdot \rangle : X \times X \longrightarrow B$$

where $(X, (\cdot \mid \cdot))$ is a *left* Hilbert A -module and $(X, \langle \cdot, \cdot \rangle)$ a *right* Hilbert B -module, besides satisfying the following transition relation:

$$(x \mid y)z = z\langle y, z \rangle$$

In order to operations make sense, it is required that $(\cdot \mid \cdot)$ to linear in the first entry, and sesquilinear in the second.

Definition 5.1.1. A Hilbert (A, B) -bimodule X is said to be *left-full* (resp. *right-full*) if $(X \mid X)$ coincides with A (resp. if $\langle X, X \rangle$ coincides with B).

Definition 5.1.2. A bimodule X is said to be an (A, B) -*imprimitivity bimodule* if X is left-full Hilbert A -module and a right-full Hilbert B -module.

Examples 5.1.3.

- (i) Every C^* -algebra A is an (A, A) -imprimitivity bimodule with $(a \mid b) = ab^*$ and $\langle a, b \rangle = a^*b$.
- (ii) Let H be a Hilbert space, which is a right Hilbert \mathbb{C} -module, with some inner product $\langle \cdot, \cdot \rangle$. If $\mathcal{K} := \mathcal{K}(H)$ is the usual C^* -algebra of compact operators, consider the \mathcal{K} -valued inner product

$$(\cdot \mid \cdot) : H \times H \longrightarrow \mathcal{K}$$

$$(x, y) \longmapsto \left(z \xmapsto{x \otimes \overline{y}} \langle z, y \rangle x \right)$$


One can check that $(x \otimes \bar{y})^* = y \otimes \bar{x}$, $(x \otimes \bar{y})(z \otimes \bar{w}) = \langle z, y \rangle (x \otimes \bar{w})$ and the operator norm of $x \otimes \bar{x}$ is $\|x\|^2$, which will lead to $(H, (\cdot | \cdot))$ being a Hilbert \mathcal{K} -module.

Notice that $(H | H) = \overline{\text{Span}\{\langle \cdot, y \rangle x\}} = \mathcal{K}$ since each $(x | y)$ is a finite-rank 1 operator. Since $\langle H, H \rangle = \mathbb{C}$, one concludes that H is a $(\mathcal{K}, \mathbb{C})$ -imprimitivity bimodule.



5.2 K -theory and Hilbert C^* -bimodules

Throughout this section, suppose that A and B are C^* -algebras and X is a left-full Hilbert (A, B) -bimodule.

Summoning 5.2.1 ([6], Corollary 4.3). If $T \in \mathcal{L}_A(E, F)$ is an A -Fredholm operator, the induced operator $(T \otimes I_X) \in \mathcal{L}_B(E \otimes_A X, F \otimes_A X)$ is B -Fredholm. 

Definition 5.2.2. Let X be a left-full Hilbert (A, B) -bimodule. For $\alpha \in K_0(A)$, choose T to be a Fredholm operator in which $\text{ind}(T) = \alpha$ using the isomorphism given by the corollary 4.3.3. This induces a morphism X_* which commutes the following diagram:


$$\begin{array}{ccc} F(A) & \xrightarrow{\cdot \otimes I_X} & F(B) \\ \text{ind} \downarrow & & \downarrow \text{ind} \\ K_0(A) & \xrightarrow{X_*} & K_0(B) \end{array}$$

This application is well defined since

$$\text{ind}(T_1) = \text{ind}(T_2) \Rightarrow \text{ind}(T_1 \otimes I_X) = \text{ind}(T_2 \otimes I_X)$$

as stated in [6], Proposition 4.4. Since tensor products are associative, one concludes that $Y_* \circ X_* = (X \otimes_B Y)_*$.

Consider X^* to be the conjugated module by $*$, which is (B, A) -imprimitivity bimodule. Considering the composition law and the summoning 5.2.3, we can consider the inverse of X_* given by $(X^*)_*: K_0(B) \longrightarrow K_0(A)$.

Summoning 5.2.3 ([6], Proposition 4.13). For any left-full Hilbert (A, B) -bimodule X , the tensor product $X \otimes_B X^*$ is a Hilbert (A, A) -bimodule isomorphic to $(A | A)$. 

Therefore, we can extract the following theorem:

Theorem 5.2.4. If A and B are strongly Morita equivalent, and X is the (A, B) -imprimitivity bimodule, then $X_* : K_0(A) \longrightarrow K_0(B)$ is an isomorphism.

Suppose that A and B are strongly Morita equivalent, and X is the certifying (A, B) -imprimitivity bimodule. If $SA := C_0(\mathbb{R}) \otimes A$ denotes the suspension C^* -algebra of A , one can induce a suspension of X , given by $SX := C_0(\mathbb{R}) \otimes X$. Since $K_1(A) \simeq K_0(SA)$ and SA and SB are Morita equivalent, one can induce an isomorphism $(SX)_* : K_1(A) \longrightarrow K_1(B)$.

Bibliography

- [1] Ch D Aliprantis and KC Border. Infinite dimensional analysis, 1994.
- [2] Armand Borel and Jean-Pierre Serre. Le théorème de riemann-roch. *Bulletin de la Société mathématique de France*, 86:97–136, 1958.
- [3] Lawrence Brown, Philip Green, and Marc Rieffel. Stable isomorphism and strong morita equivalence of C^* -algebras. *Pacific Journal of Mathematics*, 71(2):349–363, 1977.
- [4] J. Dixmier. *C^* -algebras*. North-Holland mathematical library. North-Holland, 1982.
- [5] Asen L Dontchev. Bartle-graves theorem revisited. *Set-Valued and Variational Analysis*, 28(1):109–122, 2020. <https://link.springer.com/article/10.1007/s11228-019-00524-1>.
- [6] Ruy Exel. A fredholm operator approach to morita equivalence. *K-Theory*, 7(3). <http://mtm.ufsc.br/~exel/papers/morita.pdf>.
- [7] Aweygan (<https://math.stackexchange.com/users/234668/aweygan>). Strictly positive element in unital c^* -algebra is invertible. Mathematics Stack Exchange, 2020. URL:<https://math.stackexchange.com/q/3926191> (version: 2020-11-28).
- [8] Kjeld Knudsen Jensen and Klaus Thomsen. *Elements of KK -theory*. Springer Science & Business Media, 2012.
- [9] Irving Kaplansky. Modules over operator algebras. *American Journal of Mathematics*, 75(4):839–858, 1953.
- [10] Max Karoubi. *K -theory: An introduction*, volume 226. Springer Science & Business Media, 2008.
- [11] G. G. KASPAROV. Hilbert C^* -modules : Theorems of stinespring and voiculescu. *Journal of Operator Theory*, 4(1):133–150, 1980.

- [12] E Christopher Lance. *Hilbert C^* -modules: a toolkit for operator algebraists*, volume 210. Cambridge University Press, 1995.
- [13] V. M. Manuilov and E. V. Troitsky. *Hilbert C^* -Modules*. American Mathematical Society, 2001.
- [14] James A Mingo and William J Phillips. Equivariant triviality theorems for hilbert C^* -modules. *Proceedings of the American Mathematical Society*, 91(2):225–230, 1984. <https://www.ams.org/journals/proc/1984-091-02/S0002-9939-1984-0740176-0/S0002-9939-1984-0740176-0.pdf>.
- [15] William L Paschke. Inner product modules over B^* -algebras. *Transactions of the American Mathematical Society*, 182:443–468, 1973. <https://www.ams.org/journals/tran/1973-182-00/S0002-9947-1973-0355613-0/S0002-9947-1973-0355613-0.pdf>.
- [16] Iain Raeburn and Dana P Williams. *Morita equivalence and continuous-trace C^* -algebras*. Number 60. American Mathematical Soc., 1998.
- [17] Michael Reed. *Methods of modern mathematical physics: Functional analysis*. Elsevier, 2012.
- [18] Marc Rieffel. C^* -algebras associated with irrational rotations. *Pacific Journal of Mathematics*, 93(2):415–429, 1981. <https://msp.org/pjm/1981/93-2/pjm-v93-n2-p12-s.pdf>.
- [19] Marc A Rieffel. Induced representations of C^* -algebras. *Advances in Mathematics*, 13(2):176–257, 1974. <https://www.sciencedirect.com/science/article/pii/0001870874900681>.
- [20] Alan D. Sokal. A really simple elementary proof of the uniform boundedness theorem. *The American Mathematical Monthly*, 118(5):450–452, 2011. <https://www.tandfonline.com/doi/abs/10.4169/amer.math.monthly.118.05.450>.
- [21] Johannes Ebert (<https://mathoverflow.net/users/9928/johannes-ebert>). Operators on hilbert C^* -module and families of fredholm operators. MathOverflow. URL:<https://mathoverflow.net/q/239982> (version: 2016-05-28).