

10pt Bravo!

4pt 1.1 Calculation in finite fields

Evaluate the following polynomials:

1.1.1 Evaluate the polynomial $r(X)$

$$\begin{aligned} r(X) &= 3 \cdot X^3 + 2 \cdot X^2 + X && \in GF(5)[X] \text{ at } X = 2 \\ r(X) &= (3 \cdot 2^3 + 2 \cdot 2^2 + 2) \bmod 5 && \text{because } p = 5 \text{ is prime} \\ &= (24 + 8 + 2) \bmod 5 \\ &= 34 \bmod 5 \\ &\equiv 4 \quad \checkmark \end{aligned}$$


1.1.2 Evaluate the polynomial $s(X)$

$$\begin{aligned} s(X) &= (1 + \alpha) \cdot X^3 + \alpha \cdot X^2 + X && \in GF(4)[X] \text{ at } X = \alpha \\ s(X) &= (1 + \alpha) \cdot \alpha^3 + \alpha \cdot \alpha^2 + \alpha \\ &= (1 + \alpha) \cdot \alpha \cdot \alpha \cdot \alpha + \alpha \cdot \alpha \cdot \alpha + \alpha \\ &= 1 \cdot \alpha \cdot \alpha + (1 + \alpha) \cdot \alpha + \alpha && , \text{because } (1 + \alpha) \cdot \alpha = 1 \text{ and } \alpha \cdot \alpha = 1 + \alpha \\ &= \alpha \cdot \alpha + 1 + \alpha && , \text{because } 1 \cdot \alpha = \alpha \text{ and } (1 + \alpha) \cdot \alpha = 1 \\ &= (1 + \alpha) + 1 + \alpha && , \text{because } \alpha \cdot \alpha = (1 + \alpha) \\ &= \alpha + \alpha && , \text{because } (1 + \alpha) + 1 = \alpha \\ &\equiv 0 \quad \checkmark \end{aligned}$$

3pt 1.2 Trivial functions for secure computation

- a) is trivial ✓
- b) is non-trivial, because if the value of x is greater than the value of y , e.g. $x = 5$ in $\mathbb{GF}(5)$, we cannot determine whether y was, e.g. 2 or 3. ✓
- c) is non-trivial, because if the value of x is 0, then no matter which value was assigned to y , there is no way to determine that value after the computation, because f will return 0. ✓
- d) is trivial ✓
- e) is non-trivial, because there can be more than one value which is greater or less than the chosen value x in the finite field $\mathbb{GF}(5)$, e.g. $x = 3$ then y could be 2 or 1 to produce the output 1. ✓

1.3 Non-trivial functions and an embedded OR

 b) The corresponding table looks as follows:

$\max(x, y)$	0	1	2	3	4
0	0	1	2	3	4
1	1	1	2	3	4
2	2	2	2	3	4
3	3	3	3	3	4
4	4	4	4	4	4

As one can see, when the value 4 is picked for x we cannot determine whether y was e.g. 2 or 3, because the output will always be 4. ✓

c) The corresponding table looks as follows:

$x \cdot y$	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

As one can see, when the value 0 is picked for x we cannot determine whether y was e.g. 2 or 3, because the output will always be 0. ✓

e) The corresponding table looks as follows:

$e(x, y)$	0	1	2	3	4
0	1	1	1	1	1
1	0	1	1	1	1
2	0	0	1	1	1
3	0	0	0	1	1
4	0	0	0	0	1

As one can see, when the value 2 is picked for x and the output of the function is 1, we cannot determine whether the value y was 0 or 1 (or even 2). ✓