

Suspicious Strawberries

Carolyn Wright, Rachel Donahue, Ranfei Xu, Zara Waheed



Introduction


This project is based on the analysis of Strawberry yield data that was sourced from the USDA. For the purpose of this presentation, we chose to focus on chemical usage trends over time between 1990 and 2018. We started by analysing multiple states but as we made progress, we ultimately chose to focus on California and the different types of chemicals that were prevalent in the strawberry crops.

Research Question:

How has toxic chemical usage changed over time and does this vary across different states?

Overall Toxin Instances


After cleaning out the data and arranging it in an appropriate format, we decided to look at toxins as a whole and compare the amount of the different types of chemicals detected in the strawberry yield.



The figure above shows the toxin counts across all years and states. This chart shows that Bee Toxins are at the highest detected levels. However, we wanted to assess Human toxins for this project so we decided to dig deeper into Carcinogens, as they were the second highest detected toxin.

Carcinogens per State


Map of the US with Level of Carcinogens Used in Strawberry Farming



This is a view to demonstrate where it seems Carcinogens are most prevalent. As California is a shade of deep red on this map, this indicates that they are the state out of all the 6 states included in this dataset (California, Florida, New York, North Carolina, Oregon and Washington) that has the highest amount of Carcinogens at first glance.

Carcinogens per State Over Time

We decided to compare the levels of carcinogens in the different states by their known, possible and probable quantities overtime. The scatterplots below show our findings.



We can see from the figures that we do not have a large number of data points for most of the states and the information may be insufficient for a detailed analysis. The amount of carcinogens present in most states is also very small compared to others. We chose to focus on an area with the highest level of carcinogens so that the study will have the largest impact. Based on this we chose to mainly focus on California and try to dig deeper into the types of Carcinogens detected in strawberries and their trend over time.

A Deeper Dive into Carcinogens within California


California Chemical and Toxicity Levels Counts

	known	possible	probable
BIFENTHRIN	0	10	0
BOSCALID	0	7	0
CAPTAN	13	0	0
CARBARYL	12	0	0
DIMETHOATE	0	1	0
HEXYTHIAZOX	0	0	9
IPRODIONE	9	0	0
MALATHION	0	13	0
PIPERONYL BUTOXIDE	0	7	0
PROPICONAZOLE	0	5	0
PYRIMETHANIL	0	6	0

This table shows the counts of chemicals that are considered Carcinogens along with their levels of toxicity for the state of California.


Carcinogenic Chemical Instances in California Over Time

California had 11 main Carcinogenic chemicals detected in strawberry yields. They fall into the known, possible and probable categories. The figure below shows a scaled plot of the distribution of Carcinogens detected in strawberries in California over time.



We then took a closer look at the different categories and how they are distributed among them. We can see from the figures below that most of the data we have is on known carcinogens and is also found in higher levels as compared to the possible and probable ones.

Carcinogenic Chemical Instances in California Over Time by Severity



When we centered our attention on the known toxins, there seemed to be some patterns in the data. There are three main types of known carcinogens in California; Captan, Carbaryl and Iprodione. According to the figure, Captan seemed to have an increasing trend, apart from two major dips that can be seen on the graph. Carbaryl and Iprodione seemed to have died down completely after about 2002.

We looked further into why these patterns existed and came across other sources such as the California Department of Pesticide Control (CDPC) showing similar trends in the usage of Captan over time, that verified our results. The California Office of Environmental Health Hazard Assessment (OEHHA) had Proposition 65, officially known as the Safe Drinking Water and Toxic Enforcement Act of 1986. This states Captan as a toxic chemical causing cancer but other sources such as [insert source here] state that research shows that there is no direct link between cancer and the use of Captan. It is, however, harmful if ingested in large amounts over a long period of time and Proposition 65 also states a maximum limit of Captan allowed to crops.

Even though we found no specific information explaining the patterns in Carbaryl and Iprodione, the reason could be that they were found to be toxic and eliminated completely from pesticides for crops.

Additional research and elaboration is required on the trends of known carcinogens used in Strawberry crops and our next steps would be to dig deeper into why Captan is still used as widely as it is if it is toxic, if the extreme low points can be due to a specific event or policy and whether the trends for other chemicals can be explained by further exploration.