

# MATH 263: Section 003, Tutorial 13

Mohamed-Amine Azzouz  
mohamed-amine.azzouz@mail.mcgill.ca

November 29<sup>th</sup> 2021

## 1 (Review) The Laplace Transform

The **Laplace Transform** is defined as:

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt$$

where the integral is defined.

Note that the **Laplace Transform** is an **integral Transform**, meaning that it is linear (due to the properties of the integral). Therefore, for all  $a, b \in \mathbb{R}$ :

$$\mathcal{L}\{a f(t) + b g(t)\} = a \mathcal{L}\{f(t)\} + b \mathcal{L}\{g(t)\}$$

Example: Find  $\mathcal{L}\{\cos t\}$  in two ways:

Using Taylor Series:

(*Will be discussed in the tutorial*)

Another way is by using the definition and the periodicity of  $\cos t$ :  $\cos t = \cos t + 2\pi$  :

(*Will be discussed in the tutorial*)

## 2 (Review) Solution of Initial Value Problems using The Laplace Transform

Using integration by parts, we can find the Laplace Transform of a derivative:

$$\mathcal{L}\{f'(t)\} = s\mathcal{L}\{f(t)\} - f(0)$$

note that this Laplace Transform is also in terms of the function's initial conditions. Now, from induction, we can find the Laplace Transform of the  $n$ 'th derivative of a function:

$$\mathcal{L}\{f^{(n)}(t)\} = s^n \mathcal{L}\{f(t)\} - s^{n-1} f(0) - \dots - s f^{(n-2)}(0) - f^{(n-1)}(0)$$

This Laplace Transform has  $n - 1$  initial conditions. Note that this way, we can solve first order constant coefficient ODE of the form:

$$ay''(t) + by'(t) + cy(t) = g(t)$$

By taking the Laplace Transform on both sides, using the formula to take the Laplace Transform of a derivative, solve for the solution  $Y(s)$ , and apply the Inverse Laplace Transform ( $\mathcal{L}^{-1}\{F(s)\}$ ) to find the solution to the IVP. To find inverse Laplace Transforms in this course, one should decompose  $Y(s)$  into simpler terms (usually using partial fractions) and use a Laplace Transform table to find the solution  $y(t)$ .

### 3 Solution of Initial Value Problems using The Laplace Transform

Unit step function:

$$u(x - x_0) = \begin{cases} 1 & \text{when } x \geq x_0 \\ 0 & \text{when } x < x_0 \end{cases}$$

Dirac-Delta function:

$$\delta(x - x_0) = \begin{cases} \infty & \text{when } x = x_0 \\ 0 & \text{when } x \neq x_0 \end{cases}$$

Convolution integral:

$$f(t) * g(t) = \int_0^t f(t - \tau)g(\tau)d\tau$$

, where

$$\mathcal{L}\{f(t) * g(t)\} = F(s)G(s)$$

**Problem 3a.** From Boyce and DiPrima, 10th edition (6.4, exercise 9, p.340):  
Solve this IVP using the Laplace Transform:

$$y'' + y = g(t); \quad y(0) = 0, \quad y'(0) = 1,$$

where

$$g(t) = \begin{cases} \frac{t}{2} & \text{when } 0 \leq t < 6 \\ 3 & \text{when } t \geq 6 \end{cases}$$

**Problem 3b.** From Boyce and DiPrima, 10th edition (6.5, exercise 4, p.348):  
Solve this IVP using the Laplace Transform:

$$y'' - y = -20\delta(t - 3); \quad y(0) = 1, \quad y'(0) = 0,$$