

MATH 263: Section 003, Tutorial 6

Mohamed-Amine Azzouz
mohamed-amine.azzouz@mail.mcgill.ca

October 14th 2021

1 Complex Numbers, Euler and DeMoivre's Formulas

The ODE $ay'' + by' + cy = 0$ gives the characteristic equation $ak^2 + bk + c = 0$, which does not have solutions in \mathbb{R} when $b^2 < 4ac$. Therefore, we can define a number, called i (the imaginary unit), such that $i^2 = -1$ ($i = \sqrt{-1}$). This lets us work with a new kind of numbers, called the **complex numbers**, denoted as \mathbb{C} .

$$\mathbb{C} = \{z : z = a + bi, a \in \mathbb{R}, b \in \mathbb{R}\}$$

Where a is the **real part** of z , and b is the **imaginary part** of z .

$$\text{Re}(z) = a, \text{Im}(z) = b$$

Note: In Electrical Engineering, j is used instead as a complex unit. This is to not confuse the imaginary unit i with the current variable i . Complex numbers can be added, subtracted, multiplied, and divided the same way as if i were an algebraic variable, while keeping in mind that $i^2 = -1$. Examples:

$$(3 + 2i) + (2 - 4i) = (3 + 2) + (2 - 4)i = 5 - 2i$$

$$(3 + 2i) - (2 - 4i) = (3 - 2) + (2 + 4)i = 1 + 6i$$

$$(3 + 2i)(2 - 4i) = 6 - 12i + 4i - 8i^2 = 6 - 12i + 4i - 8(-1) = 14 - 8i$$

$$\frac{3 + 2i}{2 - 4i} = \frac{-2 + 16i}{2^2 - (4i)^2} = \frac{-2 + 16i}{2^2 + 4^2} = \frac{-2 + 16i}{20} = \frac{1}{10}(-1 + 8i).$$

Note that the division process consists of multiplying the numerator and denominator by the complex conjugate of the denominator. In general, the complex conjugate of $z = a + bi$, often denoted as $z^* = a - bi$. Complex numbers can also be written in a polar form, $z = a + bi = r[\cos(\theta) + i \sin(\theta)]$, where $r = \sqrt{a^2 + b^2}$ is the norm, and $\theta = \arctan(\frac{b}{a})$ is the argument. Those two representations can be illustrated using the **complex plane**.

Defining exponentiation for complex numbers as $e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, we can let:

$$e^{ix} = \sum_{n=0}^{\infty} \frac{(ix)^n}{n!} = \sum_{\text{even } n} \frac{i^n x^n}{n!} + \sum_{\text{odd } n} \frac{i^n x^n}{n!}$$

For even numbers, let $n = 2k \Rightarrow i^n = i^{2k} = (-1)^k$. For odd numbers, $n = 2k + 1 \Rightarrow i^n = i^{2k+1} = i \cdot i^{2k} = i(-1)^k$.

$$e^{ix} = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!} + i \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(2k+1)!}$$
$$e^{ix} = \cos x + i \sin x,$$

which is **Euler's Formula**. Therefore, the polar form can be written as

$$z = a + bi = r[\cos(\theta) + i \sin(\theta)] = re^{i\theta}.$$

Similarly, **DeMoivre's Formula** is:

$$(e^{ix})^n = e^{inx} = \cos(nx) + i \sin(nx).$$

This formula can be used to multiply and find powers of complex numbers:

$$z^n = re^{in\theta} = r^n [\cos(n\theta) + i \sin(n\theta)].$$

Problem 1a. Using Euler's Formula, compute:

$$(1 - i)^{12}$$

Problem 1b. Find the general solution of:

$$\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 13x = 0$$

2 The Wronskian and Abel's Theorem

Given two solutions of a second order linear ODE, $y_1(x)$, $y_2(x)$, they are independent if their **Wronskian** is not 0, which is given by:

$$W = \begin{vmatrix} y_1(x) & y_2(x) \\ y_1'(x) & y_2'(x) \end{vmatrix} = y_1(x)y_2'(x) - y_1'(x)y_2(x)$$

Given a differential equation of the form:

$$y'' + p(x)y' + q(x)y = 0$$

Abel's Theorem states that:

$$W = c \exp\left[-\int p(x) dx\right]$$

where the constant can be found with a initial condition on the Wronskian.

Problem 2a. From Boyce and DiPrima, 10th edition (3.2, exercise 29, p.157):
Given the ODE:

$$t^2 y'' - t(t+2)y' + (t+2)y = 0$$

Find the general form of the Wronskian.

3 Euler's Equation

Euler's Equations are of the form:

$$ax^2y'' + bxy' + cy = 0$$

this is solved by making the substitution $y = x^r$, $x > 0$. The characteristic polynomial becomes:

$$ar^2 + (b-a)r + c = 0.$$

For complex roots, the solution would be of the form:

$$y(x) = c_1 x^{r_1} + c_2 x^{r_2}, \quad x > 0$$

In the case where $x < 0$, one can make the substitution $t = -x > 0$ and $y(x) = u(t)$, which would give the same ODE. Therefore, for all $x \neq 0$, the solution is:

$$y(x) = c_1 |x|^{r_1} + c_2 |x|^{r_2}$$

For complex roots $r_{1,2} = \lambda \pm i\mu$, the solution would be of the form:

$$y(x) = c_1 x^{\lambda+i\mu} + c_2 x^{\lambda-i\mu}, \quad x > 0$$

Knowing that $x^{i\mu} = e^{i\mu \ln x} = \cos(\mu \ln x) + i \sin(\mu \ln x)$, the final real solution would be:

$$y(x) = x^\lambda [k_1 \cos(\mu \ln |x|) + k_2 \sin(\mu \ln |x|)]$$

Given a double root r_1 , reduction of order gives us a solution of:

$$y(x) = (c_1 + c_2 \ln |x|)|x|^{r_1}$$

Problem 3. Find the general solution of:

$$x^2 y'' - xy' + y = 0$$

Show the two solutions are linearly independent for $x > 0$ and solve the IVP : $y(1) = 2, y'(1) = 0$.

Problem 4. Find the general solution of:

$$4x^2 y'' + 8xy' + 17y = 0$$