

### Theorem 3.21

**WTS.** *The Lebesgue Differentiation Theorem. Suppose  $f \in L^1_{loc}$ , and for every  $x \in \mathcal{L}_f$ , (so that  $x \in \mathbb{R}^n$  a.e). We have*

1.  $\lim_{r \rightarrow 0} \frac{1}{m(E_r)} \int_{E_r} |f(y) - f(x)| dy = 0,$
2.  $\lim_{r \rightarrow 0} \frac{1}{m(E_r)} \int_{E_r} f(y) dy = f(x),$

For every family  $\{E_r\}_{r>0}$  that shrinks nicely to  $x \in \mathbb{R}^{n'}$ .

*Proof.* Since the family  $\{E_r\}_{r>0}$  shrinks nicely, we have

$$m(E_r) \gtrsim m(B(r, x)) \implies m(E_r) > \alpha \cdot m(B(r, x))$$

for some  $\alpha > 0$ , independent on  $r$ . Rearranging gives

$$m^{-1}(E_r) < \alpha^{-1} m^{-1}(B(r, x))$$

And monotonicity of the integral

$$\int_{E_r} |f(y) - f(x)| dy \leq \int_{B(r, x)} |f(y) - f(x)| dy$$

Combining the last two results, for every  $\varepsilon > 0$ , if  $0 < r < \varepsilon$ , then

$$m^{-1}(E_r) \int_{E_r} |f(y) - f(x)| dy \leq m^{-1} B(r, x) \int_{B(r, x)} |f(y) - f(x)| dy$$

Taking the supremum on both sides,

$$\sup_{0 < r < \varepsilon} m^{-1}(E_r) \int_{E_r} |f(y) - f(x)| dy \leq \sup_{0 < r < \varepsilon} m^{-1} B(r, x) \int_{B(r, x)} |f(y) - f(x)| dy$$

and sending  $\varepsilon \rightarrow 0$ , proves the first claim. The second claim is immediate upon applying the  $L^1$  inequality.

Fix any  $\varepsilon > 0$ , and

$$\begin{aligned}
\lim_{r \rightarrow 0} m^{-1}(E_r) \int_{E_r} f(y) dy = f(x) &\iff \lim_{\varepsilon \rightarrow 0} \sup_{0 < r < \varepsilon} \left| m^{-1}(E_r) \int_{E_r} f(y) dy - f(x) \right| \\
&\iff \lim_{\varepsilon \rightarrow 0} \sup_{0 < r < \varepsilon} \left| m^{-1}(E_r) \int_{E_r} [f(y) - f(x)] dy \right| \\
&\leq \lim_{\varepsilon \rightarrow 0} \sup_{0 < r < \varepsilon} m^{-1}(E_r) \int_{E_r} |f(y) - f(x)| dy \\
&= \lim_{r \rightarrow 0} m^{-1}(E_r) \int_{E_r} |f(y) - f(x)| dy \\
&= 0
\end{aligned}$$

□