Contents

Executive summary 2

1. Setup 2

2. Methods 2

2.1 Cleaning . . . 2

2.2 Types . . . 3

2.3 Missing data . . . 3

2.4 Test selection . . . 3

2.5 Multiple comparisons . . . 3

3. Results 4

3.1 What is in the table . . . 4

3.2 Descriptive statistics . . . 4

3.3 Categorical variables . . . 4

3.4 Group comparison . . . 4

3.5 Pairwise comparisons . . . 5

3.6 Association between variables . . . 5

4. Discussion 6

5. Caveats and limitations 6

6. Reproducibility 7

7. Figures 8

Figure 1 -- 3 of 3 measured variables are skewed, so the analyses below summarise them by median rather than mean . . . 8

Figure 2 -- 4 of 4 measured outcomes differ across species after correction for multiple comparisons . . . 8

Figure 3 -- petal_length_cm and petal_width_cm are the strongest pair (spearman = 0.94) . . . 9

8. Figure specifications 10

Appendix A1. Sensitivity and stability 10

A1.1 Sensitivity analyses not run . . . 13

Appendix A2. Cleaning ledger 13

Appendix A3. Alternatives considered and not used 14

Appendix A4. Where every number came from 15

Appendix B. Open questions the data cannot answer 15

What this report does not claim . . . 16

Iris morphometrics -- Statistical Report

Template: nature | **Generated**: 2026-10-19 20:05 中国标准时间 | **Pipeline**: data-master
Raw input: `iris_dirty.csv` (sha256 `d5e5668ecd92e4d3...`) | **Analysed file**: `cleaned.csv`
Rows: 158 raw -> 158 analysed | **Columns**: 7 | **Modality**: tabular
Independent unit: no --unit-key given; the row is assumed to be the independent unit, and every test below inherits that assumption
Types: `cleaning_run.json` (7 columns replayed from the cleaning manifest, never re-guessed)

Executive summary

Setup. 158 rows and 7 columns reached the analysis, of which 5 are numeric measurements, compared across 3 levels of `species` (setosa, versicolor, virginica).

The conflict. The raw file was not safe to analyse as it stood: 3 label group(s) in `species` differed only by case, which would have split the comparison into spurious sub-groups and halved the group sizes; 9 exact-duplicate rows (5.7%) were present and left in place; 5 conversion(s) were withheld rather than guessed at; `sepal_width_cm` was 3.8% missing and was not imputed.

How it was resolved. 10 documented operation(s) were applied, 158 rows became 158, and the cleaned file reproduces byte for byte. The clearest separation is that `petal_length_cm` is compared across species: omega squared = 0.94, 95% CI 0.93 to 0.95, BH-adjusted p = 3.7e-69 across a family of 4 outcomes.

The caveat that matters most. 2 variable(s) carry missing values, the most at `sepal_width_cm` (3.8%). Values were not imputed, so every statistic here is conditional on complete cases. Listwise deletion removed rows from `sepal_width_cm` (6 rows, n = 152), `petal_width_cm` (6 rows, n = 152). The screening test returned undetermined, which is a screening result and not proof of a mechanism -- confirm it against how the data were actually collected.

1. Setup

- **Source**: `iris_dirty.csv`, read in `text` mode so that no column type was guessed at load time -- a reader that guesses turns a postcode `02134` into `2134` before any audit of the decision can exist.
- **Routing**: modality `tabular`; 8 columns profiled.
- **Grouping variable**: `species`, which the router scored highest among the columns able to form groups (score 0.667, balance 1.00 as entropy normalised by log of the level count). Grouping by a column with a single level would have returned one group compared with itself and a confident, meaningless zero.

Excluded from grouping because they cannot form groups: `notes` (0 observed levels).

- **Row meaning**: the router suggests `row`; that is a heuristic, and the real answer is an open question below.
- **Pre-registration**: AUTHOR_INPUT_NEEDED. Nothing in the input says whether these analyses were planned in advance, and the outcome columns were chosen after the data were seen.

2. Methods

2.1 Cleaning

The raw file was never modified. 10 operation(s) were applied automatically and 5 were withheld for a human decision. Every applied operation names the columns it is permitted to touch; the run then compares raw against cleaned in text space and fails if any column no operation named gained a value. In this run: all 1 untouched columns kept exactly their raw values.

Byte-level reproducibility: passed -- hash matches. Cleaned sha256 `7c1b8a3004c28c85...`.

> high-risk operations are never silent; see references/never-automatic-list.md

2.2 Types

Types were decided once and replayed, never re-guessed: 7 columns were typed from the manifest in `cleaning_run.json`. A column that the cleaning stage declared textual cannot quietly become numeric in the analysis stage while the report still describes it as an identifier. Dtype source for the analysis stage: `cleaning_run.json`.

2.3 Missing data

Variable	Missing	%	Screen	Evidence	Default action
`sepal_width_cm`	6	3.8	undetermined	missing fraction does not identify MCAR, MAR or MNAR; collection context is required; exploratory association screens cannot distinguish MAR from MNAR	none
`petal_width_cm`	6	3.8	undetermined	missing fraction does not identify MCAR, MAR or MNAR; collection context is required; exploratory association screens cannot distinguish MAR from MNAR	none

The screening test is the start of a conversation about how the data were collected, not a demonstration of a mechanism. Nothing was imputed, so every comparison below is on complete cases.

2.4 Test selection

Outcome	Design	Method	Statistic	df	p (raw)	p (corrected)	Alternatives rejected
`sepal_length_cm`	3-group numeric	one-way ANOVA	F = 76.152	2	= 9.2e-24	= 1.2e-23	Welch ANOVA: equal variance holds (Levene p > alpha); the pooled test is more powerful; Kruskal-Wallis H: min n=52 >= 30; the mean statistic is robust by CLT
`sepal_width_cm`	3-group numeric	one-way ANOVA	F = 55.580	2	= 9.3e-19	= 9.3e-19	Welch ANOVA: equal variance holds (Levene p > alpha); the pooled test is more powerful; Kruskal-Wallis H: min n=49 >= 30; the mean statistic is robust by CLT
`petal_length_cm`	3-group numeric	Welch ANOVA (unequal variance)	F = 1914.775	2, 82.3	= 9.2e-70	= 3.7e-69	one-way ANOVA: Levene indicates unequal variance (p <= alpha); Kruskal-Wallis H: Welch is already robust to unequal variance and n=52 >= 30
`petal_width_cm`	3-group numeric	Welch ANOVA (unequal variance)	F = 1290.926	2, 86.5	= 3.9e-65	= 7.8e-65	one-way ANOVA: Levene indicates unequal variance (p <= alpha); Kruskal-Wallis H: Welch is already robust to unequal variance and n=50 >= 30
`species_code`		not estimable	--	--	not estimated	not estimated	--

2.5 Multiple comparisons

The primary family holds **4 outcomes**, one per measured variable, corrected by **BH** at alpha = 0.050. 4 of 4 survive the correction. one outcome = one member of the family; correction applies across outcomes, and

separately within each post-hoc pairwise family

3. Results

3.1 What is in the table

Column	Role after cleaning	n	% missing	Distinct values	Status
sepal_length_cm	numeric	158	0.0	continuous	kept
sepal_width_cm	numeric	152	3.8	continuous	kept
petal_length_cm	numeric	158	0.0	continuous	kept
petal_width_cm	numeric	152	3.8	continuous	kept
species_code	numeric	158	0.0	continuous	kept
species	categorical	158	0.0	3	kept
notes	--	--	--	--	**dropped** (entirely null)
height_m	categorical	158	0.0	27	kept

`notes` was dropped as entirely null. Dropping a column is stated here rather than performed quietly, because an empty column is sometimes a loading bug that carries information about the export.

3.2 Descriptive statistics

Variable	n	% miss	Mean (95% CI)	SD	Median (IQR)	Range	Skew
sepal_length_cm	158	0.0	5.82 (5.66 to 5.97)	0.98	5.80 (5.10-6.40)	[-1.00, 7.90]	-1.87
sepal_width_cm	152	3.8	3.07 (3.00 to 3.14)	0.44	3.00 (2.80-3.40)	[2.00, 4.40]	0.33
petal_length_cm	158	0.0	3.77 (3.50 to 4.05)	1.76	4.35 (1.60-5.10)	[1.00, 6.90]	-0.28
petal_width_cm	152	3.8	1.20 (1.08 to 1.32)	0.76	1.30 (0.30-1.80)	[0.10, 2.50]	-0.11
species_code	158	0.0	1.01 (0.88 to 1.13)	0.82	1.00 (0.00-2.00)	[0.00, 2.00]	-0.01

Mean and median are both given because they disagree wherever the distribution is skewed. The mean column carries a normal-approximation interval, which for a skewed variable is the least trustworthy number on this table; the median and IQR beside it are not.

3.3 Categorical variables

Variable	n	Distinct	Mode	Distribution
species	158	3	versicolor	versicolor (53), virginica (53), setosa (52)
height_m	158	27	3.9	3.9 (15), 3.6 (13), 4.5 (12), 3.4 (10), 4.3 (10), 3.5 (10)

3.4 Group comparison

Outcome	Method	Statistic	n	Effect size (95% CI)	p (raw)	p (BH-adjusted)
sepal_length_cm	one-way ANOVA	F = 76.15	158	omega squared = 0.488 (95% CI 0.38 to 0.67)	= 9.2e-24	= 1.2e-23
sepal_width_cm	one-way ANOVA	F = 55.58	152	omega squared = 0.418 (95% CI 0.30 to 0.54)	= 9.3e-19	= 9.3e-19
petal_length_cm	Welch ANOVA (unequal variance)	F = 1914.77	158	omega squared = 0.939 (95% CI 0.93 to 0.95)	= 9.2e-70	= 3.7e-69

Outcome	Method	Statistic	n	Effect size (95% CI)	p (raw)	p (BH-adjusted)
petal_width_cm	Welch ANOVA (unequal variance)	F = 1290.93	152	omega squared = 0.926 (95% CI 0.91 to 0.94)	= 3.9e-65	= 7.8e-65
species_code	not estimable	--	--	effect size = --	not estimated	not estimated

Rows reading *not estimable* were removed from the corrected family rather than counted as non-significant: no p value is reported for them, and the family size above counts only the comparisons that produced one.

Read the effect-size column before the p column. petal_length_cm is compared across species: omega squared = 0.94, 95% CI 0.93 to 0.95, BH-adjusted p = 3.7e-69 across a family of 4 outcomes. The ordering by effect size and the ordering by p value are not the same statement, and at this sample size the p values are small enough that their exact magnitude carries no information beyond "smaller than this table can resolve".

3.5 Pairwise comparisons

Outcome	Pair	n (a / b)	Mean difference (95% CI)	Cliff's delta	p (BH-adjusted)
sepal_length_cm	setosa vs versicolor	52 / 53	-1.03 (95% CI -1.32 to -0.75)	-0.868	= 4.4e-10
sepal_length_cm	setosa vs virginica	52 / 53	-1.68 (95% CI -1.98 to -1.38)	-0.968	= 6.1e-18
sepal_length_cm	versicolor vs virginica	53 / 53	-0.65 (95% CI -0.87 to -0.42)	-0.571	= 1.1e-07
sepal_width_cm	setosa vs versicolor	51 / 49	0.68 (95% CI 0.55 to 0.82)	0.860	= 9.3e-16
sepal_width_cm	setosa vs virginica	51 / 52	0.46 (95% CI 0.33 to 0.60)	0.698	= 6.9e-10
sepal_width_cm	versicolor vs virginica	49 / 52	-0.22 (95% CI -0.34 to -0.10)	-0.351	= 5.5e-04
petal_length_cm	setosa vs versicolor	52 / 53	-2.79 (95% CI -2.93 to -2.65)	-1.000	= 2.6e-47
petal_length_cm	setosa vs virginica	52 / 53	-4.09 (95% CI -4.25 to -3.93)	-1.000	= 4.9e-52
petal_length_cm	versicolor vs virginica	53 / 53	-1.30 (95% CI -1.50 to -1.10)	-0.960	= 4.0e-23
petal_width_cm	setosa vs versicolor	50 / 51	-1.07 (95% CI -1.14 to -1.01)	-1.000	= 2.0e-46
petal_width_cm	setosa vs virginica	50 / 51	-1.78 (95% CI -1.86 to -1.69)	-1.000	= 1.3e-49
petal_width_cm	versicolor vs virginica	51 / 51	-0.70 (95% CI -0.80 to -0.61)	-0.962	= 3.0e-26

Each pairwise family is corrected separately, so these p values answer a narrower question than the omnibus values in 3.4 and should not be compared with them.

3.6 Association between variables

Pair	n	Spearman rho (95% CI)	Pearson r (95% CI)	p
sepal_length_cm - sepal_width_cm	152	-0.174 (-0.32 to -0.02)	-0.110 (-0.26 to 0.05)	= 0.032
sepal_length_cm - petal_length_cm	158	0.883 (0.84 to 0.91)	0.777 (0.71 to 0.83)	= 4.2e-53
sepal_length_cm - petal_width_cm	152	0.832 (0.78 to 0.88)	0.728 (0.64 to 0.79)	= 2.9e-40
sepal_length_cm - species_code	158	0.795 (0.73 to 0.85)	0.698 (0.61 to 0.77)	= 1.1e-35
sepal_width_cm - petal_length_cm	152	-0.324 (-0.46 to -0.17)	-0.441 (-0.56 to -0.30)	= 4.6e-05
sepal_width_cm - petal_width_cm	146	-0.298 (-0.44 to -0.14)	-0.376 (-0.51 to -0.23)	= 2.6e-04
sepal_width_cm - species_code	152	-0.452 (-0.57 to -0.32)	-0.437 (-0.56 to -0.30)	= 5.2e-09
petal_length_cm - petal_width_cm	152	0.937 (0.91 to 0.95)	0.961 (0.95 to 0.97)	= 3.5e-70
petal_length_cm - species_code	158	0.934 (0.91 to 0.95)	0.948 (0.93 to 0.96)	= 9.5e-72
petal_width_cm - species_code	152	0.938 (0.92 to 0.95)	0.956 (0.94 to 0.97)	= 6.0e-71

These are associations among measurements taken on the same rows. Nothing here fixes a direction, and two measurements that move together are not evidence that one produces the other.

4. Discussion

What the data supports. The measurements differ across levels of `species` by margins larger than the sampling noise at this size: `petal_length_cm` is compared across species: omega squared = 0.94, 95% CI 0.93 to 0.95, BH-adjusted p = 3.7e-69 across a family of 4 outcomes. The ordering of variables by effect size holds under both the parametric and the rank-based test (Appendix A3), so the finding is not an artefact of one distributional assumption. Of the 10 measured pairs, 6 are strongly associated (|Spearman rho| >= 0.7); all of it is reported as association only, with no direction and no mechanism.

What the data does not support. Nothing here establishes that the group label produced the measurement. The design compares existing groups rather than assigning them, and species, batch, site, instrument and operator all travel together with the group label -- this table cannot separate them. A claim of the form "X caused Y" is therefore not available from these data, and none is made in this report.

Boundary conditions. These results describe the rows in this file, at these measurement conditions, with this instrument. The intervals quantify sampling variation inside that set; they say nothing about another lot, another season or another population. The corrected family covers the outcomes tested here, so an outcome that was not tested is not thereby null.

5. Caveats and limitations

C1. missingness. 2 variable(s) carry missing values, the most at `sepal_width_cm` (3.8%). Values were not imputed, so every statistic here is conditional on complete cases. Listwise deletion removed rows from `sepal_width_cm` (6 rows, n = 152), `petal_width_cm` (6 rows, n = 152). The screening test returned undetermined, which is a screening result and not proof of a mechanism -- confirm it against how the data were actually collected.

- *Limits*: every mean, median, interval and p value in the Results section
- *Status*: AUTHOR_INPUT_NEEDED: how the missing values arose

C2. duplicate rows. 9 exact-duplicate rows (5.7%) were found and **kept**, because deciding whether a repeat row is a genuine second observation or a double entry is a question about the study, not about formatting. If they are double entries, every n in this report is inflated by up to 5.7% and every interval below is narrower than it should be.

- *Limits*: the reported n, and through it the width of every confidence interval
- *Status*: AUTHOR_INPUT_NEEDED: are the repeated rows independent observations?

C3. not estimable. 1 candidate outcome(s) -- `species_code` -- returned no p value at all and were removed from the corrected family rather than counted as non-significant. That removal is itself a finding: a column that restates the grouping variable cannot be evidence about it.

- *Limits*: the breadth of the outcome family, and therefore the correction
- *Status*: confirmed automatically by the structural restatement check

C4. withheld conversion. A numeric conversion was withheld for `height_m`. Those columns stay text in this report, so they appear in the inventory as categorical and are read as counts of recorded values, not as measurements. Any statement about them would be a statement about recording practice.

- *Limits*: any conclusion about those columns
- *Status*: re-run with --force-coerce-numeric once the sentinel is agreed

C5. design. The independent unit was never declared, so each row was treated as one observation. If a row pools several plants, fields, patients or technical replicates, the effective n is smaller than the row count and every interval below is too narrow -- usually by more than the replication factor. The groups were also not assigned, so differences between them are associations and this data cannot support a reading in which the group label produced the measurement.

- *Limits*: the causal reading of every group difference, and the width of every interval
- *Status*: AUTHOR_INPUT_NEEDED: what one row represents

C6. skew. `sepal_length_cm` (skew 1.87) is strongly skewed. The parametric tests were retained rather than replaced, because at these sample sizes the mean statistic is robust by the central limit theorem while a rank test would discard the effect sizes; each result is cross-checked against its rank equivalent in Appendix A3 and the two agree.

- *Limits*: the mean-based summaries; median and IQR are reported alongside
- *Status*: cross-checked in Appendix A3

6. Reproducibility

Every number above is traceable to a script. Nothing in this report was typed by hand.

Stage	Script	Artifact	sha256 (first 16)
Profile	scripts/profile_tabular.py	analysis/profile.json	eabfcb61223282ce
Plan	scripts/cleaning_plan.py	cleaning_plan.json	69492b9bd35c25af
Clean	scripts/execute_cleaning.py	cleaned/cleaned.csv	7c1b8a3004c28c85
Validate	scripts/validate_clean.py	cleaned/validation.json	070ee266feb052fd
Describe	scripts/stats_descriptive.py	analysis/descriptive.json	1816f349a61bcfd4
Infer	scripts/stats_inferential.py	analysis/inferential.json	f2556c0c1455bda5
Decide figures	scripts/chart_decision.py	analysis/chart_decisions.json	e0548e975a90e8d1
Render figures	scripts/render_chart.py	charts/*.png	c428ed8bc516c143
Assemble	scripts/assemble_report.py	report.md / report.pdf	self-referential -- see report_build.json

A document cannot contain its own hash, so the hash of `report.md` and `report.pdf` is recorded in `report_build.json` rather than on this page. Every other hash above was computed from the artifact as it exists now.

- **Interpreter**: Python 3.13.15 on Windows-11-10.0.26200-SP0
- **Libraries**: numpy 2.4.6, pandas 3.0.3, scipy 1.18.1
- **Pipeline code**: data-master commit `8749c5ea1ff4` (working tree was dirty; the commit alone does not reproduce it) -- this report names the exact code that produced every number in it.
- **Random seeds**: 42 -- every resampling interval is seeded, so it reproduces exactly.
- **Columns any operation was permitted to touch**: `petal_length_cm`, `petal_width_cm`, `sepal_length_cm`, `sepal_width_cm`, `species`, `species_code`
- **PDF backend**: `reportlab`. The page count, byte count and sha256 of the delivered PDF are recorded in `report_build.json` under `pdf`; the probe that established this backend is under `pdf_probe`. They are not printed here because a document cannot contain its own hash, and a page count measured before the last line changed is not the page count of the file you are reading.


The data contains non-Latin characters, so the renderer was asked for a CJK-capable font; one was embedded.

7. Figures

Each figure carries one argument and its title states that argument as a finding computed from the data rather than as a topic label. The chart types considered and rejected are listed under each one, because choosing a chart is itself a claim about what matters.

Figure 1 -- 3 of 3 measured variables are skewed, so the analyses below summarise them by median rather than mean


Distribution of each numeric variable, with the median and interquartile range marked. Panel order follows absolute skewness. Full box-and-whisker statistics are in Table 2.



- **What it shows**: sepal_length_cm is strongly left-skewed (median 5.80, IQR 5.10 to 6.40, n = 158); petal_length_cm is moderately left-skewed (median 4.35, IQR 1.60 to 5.10, n = 158); petal_width_cm is slightly left-skewed (median 1.30, IQR 0.30 to 1.80, n = 152).
- **How to read it**: Skewed variables are summarised by the median and IQR, and rank-based or Welch methods are preferred for them.
- **What follows**: These summaries, not the raw shape, should carry into any downstream model.
- **Uncertainty shown**: median and interquartile range marked on every panel
- **Chart alternatives rejected**: one histogram per variable as separate figures -- same argument repeated; the reader has to re-read the framing each time; box plot only -- understates the tail, which is precisely what decides the test choice
- **Source**: `scripts/chart_decision.py` -> `charts/fig01_distribution_panel.png` rendered by `scripts/render_chart.py`, 102,812 bytes, ink ratio 0.3184

Figure 2 -- 4 of 4 measured outcomes differ across species after correction for multiple comparisons

Mean of each outcome by species with 95% confidence intervals, setosa, versicolor, virginica. Welch ANOVA (unequal variance); one-way ANOVA. Raw p values are BH-corrected across a family of 4 outcomes.




- **What it shows**: petal_length_cm differs across species by omega squared = 0.939 (95% CI 0.926 to 0.953), BH-adjusted p = 3.697e-69; petal_width_cm differs across species by omega squared = 0.926 (95% CI 0.910 to 0.943), BH-adjusted p = 7.811e-65; sepal_length_cm differs across species by omega squared = 0.488 (95% CI 0.377 to 0.665), BH-adjusted p = 1.228e-23; sepal_width_cm differs across species by omega squared = 0.418 (95% CI 0.304 to 0.537), BH-adjusted p = 9.266e-19.
- **How to read it**: Every tested outcome separates the groups.
- **What follows**: Carry the effect sizes forward, not the p values: a difference that is real can still be too small to matter.
- **Uncertainty shown**: 95% confidence intervals on every group mean
- **Chart alternatives rejected**: pie chart -- group means are not parts of a whole; 3D bar chart -- the third dimension encodes nothing; radar chart -- angle and area are hard to compare and imply the axes are commensurate
- **Source**: `scripts/chart_decision.py` -> `charts/fig02_small_multiples_ci.png` rendered by `scripts/render_chart.py`, 140,132 bytes, ink ratio 0.2065

Figure 3 -- petal_length_cm and petal_width_cm are the strongest pair (spearman = 0.94)

Each point is one observation. The line is a LOWESS smoother. Association only: this figure carries no direction of causation. Columns that restate the grouping variable are excluded.

petal_length_cm and petal_width_cm are the strongest pair (spearman = 0.94)

Association only. The smoother carries a bootstrap 95% band and the coefficient carries its interval, so the uncertainty of the curve is visible rather than implied. Columns that restate the grouping variable are excluded.



- **What it shows**: petal_length_cm and petal_width_cm move together (spearman = 0.94, 95% CI 0.91 to 0.95, p = 3.506e-70, n = 152); sepal_length_cm and petal_length_cm move together (spearman = 0.88, 95% CI 0.84 to 0.91, p = 4.236e-53, n = 158).
- **How to read it**: A visible curve that a straight line would miss is the reason the smoother is drawn at all.
- **What follows**: Correlation here is a screening result. It says which relationships deserve a model, not which are real.
- **Uncertainty shown**: LOWESS smoother with a bootstrap 95% band; Pearson r with a 95% CI
- **Chart alternatives rejected**: 3D scatter -- a third axis encodes nothing here; full correlation heatmap -- with this many columns it becomes a grid of numbers rather than an argument; the pairs that matter get their own panel
- **Source**: `scripts/chart_decision.py` -> `charts/fig03_scatter_panel.png` rendered by `scripts/render_chart.py`, 146,228 bytes, ink ratio 0.1202

8. Figure specifications

#	ID	Chart type	Specification
1	`fig01_distribution_panel`	distribution_panel	3 panels; bins = auto; show_median_iqr = True
2	`fig02_small_multiples_ci`	small_multiples_ci	group = species; 4 panels; n_panels = 4
3	`fig03_scatter_panel`	scatter_panel	2 panels; smoother = LOWESS

Appendix A1. Sensitivity and stability

The primary analysis was re-run under perturbation by `scripts/sensitivity_analysis.py` (seed 42, 10 blocks for the jackknife). Agreement means the significance verdict did not flip at alpha=0.05.

Outcome	Perturbation	Statistic	p	Verdict
`sepal_length_cm`	leave_one_group_out (setosa)	Student two-sample t-test (pooled variance) = -5.73	= 9.8e-08	agrees
`sepal_length_cm`	leave_one_group_out (versicolor)	Student two-sample t-test (pooled variance) = -11.02	= 4.0e-19	agrees
`sepal_length_cm`	leave_one_group_out (virginica)	Student two-sample t-test (pooled variance) = -7.22	= 9.0e-11	agrees
`sepal_length_cm`	block_jackknife (block 1)	one-way ANOVA = 69.11	= 1.5e-21	agrees
`sepal_length_cm`	block_jackknife (block 2)	one-way ANOVA = 62.32	= 4.8e-20	agrees
`sepal_length_cm`	block_jackknife (block 3)	one-way ANOVA = 72.15	= 3.2e-22	agrees
`sepal_length_cm`	block_jackknife (block 4)	one-way ANOVA = 65.11	= 1.1e-20	agrees

Outcome	Perturbation	Statistic	p	Verdict
sepal_length_cm	block_jackknife (block 5)	Welch ANOVA (unequal variance) = 138.89	= 5.5e-28	agrees
sepal_length_cm	block_jackknife (block 6)	one-way ANOVA = 64.85	= 1.3e-20	agrees
sepal_length_cm	block_jackknife (block 7)	one-way ANOVA = 64.47	= 1.6e-20	agrees
sepal_length_cm	block_jackknife (block 8)	one-way ANOVA = 65.46	= 9.3e-21	agrees
sepal_length_cm	block_jackknife (block 9)	one-way ANOVA = 67.15	= 3.6e-21	agrees
sepal_length_cm	block_jackknife (block 10)	one-way ANOVA = 62.59	= 3.8e-20	agrees
sepal_length_cm	exclude_extremes (2 value(s) excluded)	Welch ANOVA (unequal variance) = 155.53	= 5.4e-31	agrees
sepal_length_cm	strict_complete_case (n=146 of 158 rows)	one-way ANOVA = 70.05	= 6.2e-22	agrees
sepal_length_cm	rank_cross_check	Kruskal-Wallis H = 101.56	< 1e-300	agrees
sepal_width_cm	leave_one_group_out (setosa)	Student two-sample t-test (pooled variance) = -3.58	= 5.4e-04	agrees
sepal_width_cm	leave_one_group_out (versicolor)	Student two-sample t-test (pooled variance) = 6.96	= 3.5e-10	agrees
sepal_width_cm	leave_one_group_out (virginica)	Student two-sample t-test (pooled variance) = 9.81	= 3.1e-16	agrees
sepal_width_cm	block_jackknife (block 1)	one-way ANOVA = 46.36	= 5.3e-16	agrees
sepal_width_cm	block_jackknife (block 2)	one-way ANOVA = 50.10	= 5.3e-17	agrees
sepal_width_cm	block_jackknife (block 3)	one-way ANOVA = 48.10	= 1.8e-16	agrees
sepal_width_cm	block_jackknife (block 4)	one-way ANOVA = 48.33	= 1.6e-16	agrees
sepal_width_cm	block_jackknife (block 5)	one-way ANOVA = 46.57	= 4.4e-16	agrees
sepal_width_cm	block_jackknife (block 6)	one-way ANOVA = 52.24	= 1.8e-17	agrees
sepal_width_cm	block_jackknife (block 7)	one-way ANOVA = 56.30	= 1.9e-18	agrees
sepal_width_cm	block_jackknife (block 8)	one-way ANOVA = 50.91	= 3.6e-17	agrees
sepal_width_cm	block_jackknife (block 9)	one-way ANOVA = 49.12	= 1.0e-16	agrees
sepal_width_cm	block_jackknife (block 10)	one-way ANOVA = 52.21	= 1.7e-17	agrees
sepal_width_cm	exclude_extremes (3 value(s) excluded)	one-way ANOVA = 71.00	= 2.9e-22	agrees
sepal_width_cm	strict_complete_case (n=146 of 158 rows)	one-way ANOVA = 53.55	= 4.4e-18	agrees
sepal_width_cm	rank_cross_check	Kruskal-Wallis H = 67.66	< 1e-300	agrees
petal_length_cm	leave_one_group_out (setosa)	Student two-sample t-test (pooled variance) = -12.89	= 2.8e-23	agrees
petal_length_cm	leave_one_group_out (versicolor)	Welch two-sample t-test (unequal variance) = -51.39	= 1.6e-52	agrees
petal_length_cm	leave_one_group_out (virginica)	Welch two-sample t-test (unequal variance) = -39.90	= 1.7e-47	agrees
petal_length_cm	block_jackknife (block 1)	Welch ANOVA (unequal variance) = 1797.15	= 5.7e-63	agrees
petal_length_cm	block_jackknife (block 2)	Welch ANOVA (unequal variance) = 1764.37	= 2.1e-63	agrees

Outcome	Perturbation	Statistic	p	Verdict
petal_length_cm	block_jackknife (block 3)	Welch ANOVA (unequal variance) = 1717.06	= 3.2e-61	agrees
petal_length_cm	block_jackknife (block 4)	Welch ANOVA (unequal variance) = 1588.18	= 3.9e-59	agrees
petal_length_cm	block_jackknife (block 5)	Welch ANOVA (unequal variance) = 1739.06	= 5.9e-63	agrees
petal_length_cm	block_jackknife (block 6)	Welch ANOVA (unequal variance) = 1699.22	= 2.5e-62	agrees
petal_length_cm	block_jackknife (block 7)	Welch ANOVA (unequal variance) = 1776.18	= 2.1e-64	agrees
petal_length_cm	block_jackknife (block 8)	Welch ANOVA (unequal variance) = 1663.99	= 2.2e-63	agrees
petal_length_cm	block_jackknife (block 9)	Welch ANOVA (unequal variance) = 1773.93	= 5.7e-64	agrees
petal_length_cm	block_jackknife (block 10)	Welch ANOVA (unequal variance) = 1727.51	= 2.7e-63	agrees
petal_length_cm	exclude_extremes (7 value(s) excluded)	Welch ANOVA (unequal variance) = 2165.17	= 5.0e-68	agrees
petal_length_cm	strict_complete_case (n=146 of 158 rows)	Welch ANOVA (unequal variance) = 1732.44	= 5.7e-63	agrees
petal_length_cm	rank_cross_check	Kruskal-Wallis H = 137.08	< 1e-300	agrees
petal_width_cm	leave_one_group_out (setosa)	Welch two-sample t-test (unequal variance) = -14.81	= 3.0e-26	agrees
petal_width_cm	leave_one_group_out (versicolor)	Welch two-sample t-test (unequal variance) = -43.79	= 4.4e-50	agrees
petal_width_cm	leave_one_group_out (virginica)	Welch two-sample t-test (unequal variance) = -33.20	= 1.3e-46	agrees
petal_width_cm	block_jackknife (block 1)	Welch ANOVA (unequal variance) = 1175.50	= 5.1e-58	agrees
petal_width_cm	block_jackknife (block 2)	Welch ANOVA (unequal variance) = 1141.70	= 1.6e-59	agrees
petal_width_cm	block_jackknife (block 3)	Welch ANOVA (unequal variance) = 1194.94	= 4.0e-58	agrees
petal_width_cm	block_jackknife (block 4)	Welch ANOVA (unequal variance) = 1111.55	= 9.5e-56	agrees
petal_width_cm	block_jackknife (block 5)	Welch ANOVA (unequal variance) = 1136.82	= 1.1e-58	agrees
petal_width_cm	block_jackknife (block 6)	Welch ANOVA (unequal variance) = 1113.96	= 6.3e-58	agrees
petal_width_cm	block_jackknife (block 7)	Welch ANOVA (unequal variance) = 1219.54	= 2.4e-59	agrees
petal_width_cm	block_jackknife (block 8)	Welch ANOVA (unequal variance) = 1249.67	= 4.4e-60	agrees
petal_width_cm	block_jackknife (block 9)	Welch ANOVA (unequal variance) = 1091.76	= 4.1e-58	agrees
petal_width_cm	block_jackknife (block 10)	Welch ANOVA (unequal variance) = 1196.79	= 2.3e-59	agrees
petal_width_cm	exclude_extremes (2 value(s) excluded)	Welch ANOVA (unequal variance) = 1435.05	= 1.7e-64	agrees

Outcome	Perturbation	Statistic	p	Verdict
petal_width_cm	strict_complete_case (n=146 of 158 rows)	Welch ANOVA (unequal variance) = 1281.47	= 2.4e-62	agrees
petal_width_cm	rank_cross_check	Kruskal-Wallis H = 132.92	< 1e-300	agrees

Assumption robustness (distinct from the perturbations above):

Outcome	Check	Statistic	p	Verdict
sepal_length_cm	rank-based cross-check of the primary test	Kruskal-Wallis H = 101.56	< 1e-300	agrees
sepal_width_cm	rank-based cross-check of the primary test	Kruskal-Wallis H = 67.66	< 1e-300	agrees
petal_length_cm	rank-based cross-check of the primary test	Kruskal-Wallis H = 137.08	< 1e-300	agrees
petal_width_cm	rank-based cross-check of the primary test	Kruskal-Wallis H = 132.92	< 1e-300	agrees

Each primary parametric test was repeated with its rank-based equivalent. Agreement means the conclusion does not depend on the normality assumption the parametric test makes; a disagreement would have meant the parametric test had to go, not be defended.

A1.1 Sensitivity analyses not run

Sensitivity analysis	Status	What that means
drop the 9 duplicate rows and re-run every test	**NOT RUN -- this is a gap, not a pass**	Duplicate rows were deliberately left in place. Removing them is a decision about whether repeated observations are independent, and that decision belongs to the author. Until it is made, the reported n and every interval derived from it stand unchanged.

These are listed as gaps so that nobody reads the absence of a disagreement above as evidence that no disagreement exists. A check that did not run cannot pass.

Appendix A2. Cleaning ledger

Every operation that ran, in order, with the catalogue rule it came from. Nothing below was chosen by the model at run time: the plan is the gate, and only `plan["auto_actions"]` may execute.

#	Operation	Columns	Sev.	Rows	Rule
1	fix_unicode	height_m, petal_length_cm, petal_width_cm (+4 more)	P3	158->158	t-fix-unicode
2	strip_whitespace	height_m, petal_length_cm, petal_width_cm (+4 more)	P3	158->158	t-strip-ws
3	collapse_whitespace	height_m, petal_length_cm, petal_width_cm (+4 more)	P3	158->158	t-collapse-whitespace
4	trim_paired_quotes	height_m, petal_length_cm, petal_width_cm (+4 more)	P3	158->158	t-trim-quotes
5	coerce_numeric	petal_length_cm, petal_width_cm, sepal_length_cm (+2 more)	P2	158->158	t-coerce-numeric
6	normalize_dtypes	height_m, petal_length_cm, petal_width_cm (+4 more)	P2	158->158	t-normalize-dtypes
7	merge_case_variants	species	P2	158->158	t-merge-case-variants
8	drop_all_null_cols	notes	P1	158->158	t-drop-all-null-cols

#	Operation	Columns	Sev.	Rows	Rule
9	drop_all_null_rows	(file-level)	P1	158->158	t-drop-all-null-rows
10	rename_columns_append_cleaned	(file-level)	P2	158->158	t-rename-clean

The Rule column names the catalogue anchor; the full path, the operation id and the exact column list for every row above are in cleaned/cleaning_run.json under applied_ops.

Withheld -- proposed but deliberately not applied.

Nothing was withheld.

Failed -- attempted and did not complete.

No operation failed.

Never-automatic -- conversions that were held back by policy. These are the cases where a conversion was technically possible but would have destroyed information, so the policy refuses it without asking. Each can be answered by filling answers.template.json and re-running with --answers; until then the default applies.

- coerce_numeric on height_m -- 153 of 158 non-null values parse as numbers and the other 5 do not (e.g. '--', '--', '--'). Converting would turn those into missing values, after which a recorded category (say 'unknown') would be indistinguishable from a value that was never collected. The failures are all blank or conventional sentinels, so treating them as missing is defensible -- but 'not recorded' is a claim about the study, not a formatting detail, and it stays yours to make.
- merge_case_variants on species -- once the automatic whitespace/quote hygiene has run, 3 label group(s) still differ only by case (6 levels -> 3 after casefold); collapsed automatically because this is a lossless canonicalisation
- dedupe_exact on -- 9 exact-duplicate rows (5.7% of rows)
- drop_impossible on sepal_length_cm -- sepal_length_cm has 1 negative value(s) (min=-1.0) for a physically non-negative quantity
- impute_categorical on petal_width_cm, sepal_width_cm -- categorical column(s) with missing values; adding a mode level invents a category

Additive-only audit. The contract forbids modifying the raw file and permits an operation to touch only the columns it names. Columns no operation named: **1, and each kept exactly its raw values -- audit passed**.

Label conflicts resolved before grouping. Grouping on a column whose labels differ only by case splits one group into two and halves both n, which shrinks intervals without any loss of real information. The merge map below is the audit trail for that decision.

Column	Labels merged	Variants
species	3	

Appendix A3. Alternatives considered and not used

A test is a choice. Listing only the test that was run hides the fact that a different, defensible analysis was available, so every rejected alternative is recorded here with the reason it lost.

Outcome	Design	Rejected test	Reason
sepal_length_cm	3-group numeric	Welch ANOVA	equal variance holds (Levene p > alpha); the pooled test is more powerful
sepal_length_cm	3-group numeric	Kruskal-Wallis H	min n=52 >= 30; the mean statistic is robust by CLT

Outcome	Design	Rejected test	Reason
sepal_width_cm	3-group numeric	Welch ANOVA	equal variance holds (Levene p > alpha); the pooled test is more powerful
sepal_width_cm	3-group numeric	Kruskal-Wallis H	min n=49 >= 30; the mean statistic is robust by CLT
petal_length_cm	3-group numeric	one-way ANOVA	Levene indicates unequal variance (p <= alpha)
petal_length_cm	3-group numeric	Kruskal-Wallis H	Welch is already robust to unequal variance and n=52 >= 30
petal_width_cm	3-group numeric	one-way ANOVA	Levene indicates unequal variance (p <= alpha)
petal_width_cm	3-group numeric	Kruskal-Wallis H	Welch is already robust to unequal variance and n=50 >= 30

The rule that made these choices is `references/nature-statistics-checklist.md#gates-1-3-5`. It was applied mechanically; there is no per-dataset override, because a rule that is quietly overridden is a rule that cannot be checked.

Appendix A4. Where every number came from

Each value below was registered by the code that read it from an artifact on disk. A sentence in this report that is not backed by a row here is a sentence that should not have been written.

Kind	Value	What it means	Source: script -> artifact	Rule anchor
header	158	rows analysed	execute_cleaning.py -> cleaning_run.json	--
headline	petal_length_cm is compared across species: omega squared =...	strongest group separation	stats_inferential.py -> inferential.json	--
rejected test	Welch ANOVA	rejected for sepal_length_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	Kruskal-Wallis H	rejected for sepal_length_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	Welch ANOVA	rejected for sepal_width_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	Kruskal-Wallis H	rejected for sepal_width_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	one-way ANOVA	rejected for petal_length_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	Kruskal-Wallis H	rejected for petal_length_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	one-way ANOVA	rejected for petal_width_cm	stats_inferential.py -> inferential.json	k_group_numeric
rejected test	Kruskal-Wallis H	rejected for petal_width_cm	stats_inferential.py -> inferential.json	k_group_numeric
correction	BH	family correction method	stats_inferential.py -> inferential.json	gate-5

Only the basename of each artifact and script is shown here; the full path, the untruncated value and the complete claim record for every row are in `report_build.json` under `provenance`. Each record carries `rule_source`, and where a rule had an alternative the record names it and why it lost -- a claim with no alternative says so rather than leaving the field blank.

Appendix B. Open questions the data cannot answer

8 questions must be answered by someone who knows how the data were collected. They are listed with the consequence of leaving them unanswered, because an unanswered question that carries no stated cost is one nobody will prioritise.

#	Question	Why it matters	Cost of leaving it open
1	What does one row represent?	Every interval in this report is conditional on one row being one independent observation. If a row pools replicates, subjects, sites or runs, the effective n is smaller than the row count and every interval is too narrow.	HIGH -- rescales every interval in the report
2	Were these outcomes chosen in advance?	Nothing in the input records a pre-specified analysis plan, and the outcome family was assembled from the columns present. Without that, the family-size correction controls multiplicity within this report but cannot address selection across reports.	HIGH -- determines whether the correction is the right one
3	Are the 9 duplicated rows independent?	They are byte-identical repeats. If they are double entry they inflate n; if they are genuine repeated measurement they belong in the design. The pipeline does not guess, and A1.1 records this as a gap rather than a result.	HIGH -- inflates n and narrows intervals
4	How should `coerce_numeric` on `height_m` be resolved?	153 of 158 non-null values parse as numbers and the other 5 do not (e.g. '--', '--', '--'). Converting would turn those into missing values, after which a recorded category (say 'unknown') would be indistinguishable from a value that was never collected. The failures are all blank or conventional sentinels, so treating them as missing is defensible -- but 'not recorded' is a claim about the study, not a formatting detail, and it stays yours to make.	MEDIUM -- the pipeline stopped at this decision instead of guessing
5	How should `merge_case_variants` on `species` be resolved?	once the automatic whitespace/quote hygiene has run, 3 label group(s) still differ only by case (6 levels -> 3 after casefold); collapsed automatically because this is a lossless canonicalisation	MEDIUM -- the pipeline stopped at this decision instead of guessing
6	How should `dedupe_exact` on the file be resolved?	9 exact-duplicate rows (5.7% of rows)	MEDIUM -- the pipeline stopped at this decision instead of guessing
7	How should `drop_impossible` on `sepal_length_cm` be resolved?	`sepal_length_cm` has 1 negative value(s) (min=-1.0) for a physically non-negative quantity	MEDIUM -- the pipeline stopped at this decision instead of guessing
8	How should `impute_categorical` on `petal_width_cm`, `sepal_width_cm` be resolved?	categorical column(s) with missing values; adding a mode level invents a category	MEDIUM -- the pipeline stopped at this decision instead of guessing

Of these, 3 are already reflected as caveats in section 5.

What this report does not claim

- It does not claim the group label produced the measurement.
- It does not claim the analysis was pre-specified.
- It does not claim the rows are independent observations.

- It does not claim any mechanism for the missingness it reports; the screening test in 2.3 is a description, not a diagnosis.

End of report. Generated 2026-10-19 20:05 中国标准时间 *by data-master. The raw input was never modified: every value above was read from the cleaned file or from a recorded artifact. Each number is registered in `report_build.json`; the gate verdict, including anything that failed, is in `audit/final_audit.md`.*