

Cambridge International Examinations

Cambridge International Advanced Level

CANDIDATE NAME		
CENTRE NUMBER	CANDIDATE NUMBER	

BIOLOGY 9700/52

Paper 5 Planning, Analysis and Evaluation

May/June 2015
1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.



1 Many fungi are decomposer organisms which carry out extracellular digestion. To do this they secrete a number of enzymes.

A group of students made a solution of enzyme extract from a fungus. The extract contained the enzyme amylase. They wanted to find out the concentration of amylase in the extract.

They were provided with:

- 0.5 g dm⁻³ stock solution of amylase
- starch agar plates with wells into which enzyme solutions can be placed. Starch agar plates are Petri dishes containing agar mixed with starch.

Fig. 1.1 shows how the students used the plates to find the concentration of amylase.




Fig. 1.1

The students thought that the area stained brown was proportional to the amylase concentration.

(a) Identify the independent and dependent variables in this investigation.

independent variable	
dependent variable	[2]

© UCLES 2015 9700/52/M/J/15

		ıld be deta	9			
• • • • • • • • • • • • • • • • • • • •			 •	 	 	
	••••••		 	 	 	

- **(c)** There are different types of amylase enzyme. They hydrolyse starch in different ways. Two of these enzymes are:
 - β -amylase hydrolyses every second α -1,4 glycosidic bond in starch molecules
 - γ -amylase hydrolyses all α -1,6 glycosidic bonds and all α -1,4 glycosidic bonds in starch molecules.

In a second investigation, the students were provided with two beakers, **A** and **B**. One contained β -amylase and the other contained γ -amylase. They used these solutions to hydrolyse $25\,\text{cm}^3$ samples of $0.5\,\text{g}\,\text{dm}^{-3}$ starch solution.

Suggest and explain how the students could identify which beaker contained β -amylase and which contained γ -amylase.
[2]

(d) Humans produce the enzyme α -amylase in their salivary glands. There may be many copies of the gene coding for α -amylase on chromosome 1. The concentration of the α -amylase in the saliva is positively correlated with the number of copies of this gene.

In a third investigation, the students obtained saliva from six people, **A** to **F**. Equal volumes of saliva were added to wells in agar plates similar to those shown in Fig. 1.1. The plates were incubated for the same length of time and the area of the brown zone for each sample of saliva was calculated.

Table 1.1 shows results of this investigation.

Table 1.1

onzymo ovtract	area of brown zone/mm ²							
enzyme extract	plate 1	plate 2	plate 3	plate 4	plate 5	plate 6		
Α	3632	3848	3632	3632	3632	3848		
В	2827	2827	2642	2463	1963	2827		
С	2124	1963	1963	2124	1963	2124		
D	1385	1257	1809	1257	1257	1385		
E	656	707	707	656	707	656		
F	298	298	314	314	298	298		

© UCLES 2015 9700/52/M/J/15

					ir results using
		s =	$\sqrt{\frac{\sum (x - \overline{x})^2}{n - 1}}$		
K	Key to Symbols	·	, , ,		
S	= standard deviation	on $x = a res$	sult $\overline{x} = \text{mean}$	$\Sigma = \text{sum of}$	n = sample s
	plate	х	$x-\overline{x}$	$(x-\overline{x})^2$	
			able 1.2	2	
	-		x - x	$(x-x)^{-}$	
		298			
	1				
	2	298			
		298 314			
	2	298			
	2 3	298 314			
	2 3 4	298 314 314			
	2 3 4 5	298 314 314 298			

.....[3]

2 The speed at which an electrical impulse travels along a nerve can be determined by carrying out a nerve conduction velocity (NCV) test.

Surface electrodes are placed on the skin over nerves at various locations. They produce a very mild electrical charge, which stimulates the nerve.

The resulting electrical activity in the nerve is measured by a recording electrode. The distance between the electrodes and the time it takes for electrical impulses to travel between them are used to determine the nerve conduction velocity.

Fig. 2.1 shows how the NCV is measured in the ulnar nerve of the human forearm.




Fig. 2.1

An investigation to measure the NCV in the ulnar nerve in females of different ages was carried out on 394 individuals.

(a)	Suggest three variables which the investigators should have standardised.
	[3]

© UCLES 2015 9700/52/M/J/15

Table 2.1 shows the results of this investigation.

Table 2.1

age category/years	mean conduction	confidence limits			
	velocity $\pm S_M$	lower limit	upper limit		
30–39	54.3 ± 1.200	51.90	56.70		
40–49	54.7 ± 0.645	53.41	55.99		
50–59	52.4 ± 0.600	51.20	53.60		
60–69	52.2 ± 0.675				
70–79	49.0 ± 1.075	46.85	51.15		

 S_M = standard error

(b)	The	confidence	limit =	mean	±	2	S_M
-----	-----	------------	---------	------	---	---	-------

Use this formula to calculate the missing confidence limits. Use the space below for any working and enter your answers in Table 2.1.

[1]

One conclusion from these data is that mean conduction velocity in the ulnar nerve varies significantly with age.

(c)	(i)	Identify two age categories which appear to support this conclusion and give a reason
		for your choice.

age categories	. and
reason	
	[2]

(ii)	State which statistical test could have been used to confirm this conclusion and give a reason for your choice.
	test
	reason
	[2]
(iii)	State a null hypothesis for this test.
	[1]
(d) Sta	te one reason why the results of the investigation were considered to be reliable.
	[1]
	[Total: 10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.