

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

BIOLOGY 9700/51

Paper 5 Planning, Analysis and Evaluation

October/November 2009
1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use				
1				
2				
3				
Total				

This document consists of **7** printed pages and **5** blank pages.



An investigation into the effect of temperature on the rate of the light-dependent stage of photosynthesis was carried out using isolated chloroplasts. Samples of chloroplasts suspended in buffer were mixed with a coloured electron acceptor and exposed to light. The colour changes from blue to colourless as electrons are taken up by the electron acceptor. (a) (i) Sketch a graph to predict the results of the investigation. [2] Identify two key variables that must be controlled in this investigation. For each explain how it might be controlled.[4] (iii) Outline a procedure to find the rate of reaction for this investigation. (b) In a further investigation, small quantities of ADP and inorganic phosphate were added to the isolated chloroplasts before testing. Suggest an hypothesis being tested by this further investigation.

[Total: 9]

For

Examiner's Use A solution of substance Y, thought to be a growth hormone, was made by dissolving a known mass of Y in 10 cm³ of distilled water. This solution was added to samples from a culture of animal cells containing 3000 cells per mm³.


For Examiner's Use


- 25 mm³ of solution Y was added to a cell sample.
- 25 mm³ of distilled water was added to another cell sample.

After four days the number of cells per mm³ of each culture was estimated using a microscope slide with a counting grid.

(a)	Identify and explain the purpose of the control experiment used in this investigation.
	2]

(b) Fig. 2.1 shows a top view of a microscope slide with a counting grid. Fig. 2.2 shows a vertical section through the microscope slide and grid. Fig. 2.3 shows the detail of part of the grid viewed through a microscope.





© UCLES 2009

For Examiner's Use

c) Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture sample number 1 2 3 4 5 6 7 8 9 10 mean experimental culture 7.5 8.1 7.6 6.2 7.5 7.8 8.9 6.5 7.9 7.3 7.5 control culture 5.6 7.5 8.2 6.7 3.5 6.5 5.9 3.7 5.8 8.4 (i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the coculture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number — original number) × 100 original number Calculate the percentage increase in the control culture. Show your working.		ggest how this ap ture.	parat	us co	uld be	e use	d to e	estima	ite the	e num	nber d	of cells	s per mi	m ³
Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture sample number 1 2 3 4 5 6 7 8 9 10 mean experimental culture 7.5 8.1 7.6 6.2 7.5 7.8 8.9 6.5 7.9 7.3 7.5 control culture 5.6 7.5 8.2 6.7 3.5 6.5 5.9 3.7 5.8 8.4 (i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the coculture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number — original number) × 100 original number Calculate the percentage increase in the control culture.														
Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture ample number														
Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture ample number														
Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture ample number														
Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture ample number														
Table 2.1 shows estimated number of cells in the experimental and control cultures three days growth. Table 2.1 thousands of cells per mm³ of culture ample number														
three days growth. Table 2.1 thousands of cells per mm³ of culture ample number														
thousands of cells per mm³ of culture ample number 1 2 3 4 5 6 7 8 9 10 mean experimental culture 7.5 8.1 7.6 6.2 7.5 7.8 8.9 6.5 7.9 7.3 7.5 ontrol culture 5.6 7.5 8.2 6.7 3.5 6.5 5.9 3.7 5.8 8.4 (i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the co-culture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.	-		mated	l num	ber of	f cells	in the	e expe	erime	ntal a	nd co	ntrol	cultures	af
ample number 1 2 3 4 5 6 7 8 9 10 mean experimental culture 7.5 8.1 7.6 6.2 7.5 7.8 8.9 6.5 7.9 7.3 7.5 ontrol culture 5.6 7.5 8.2 6.7 3.5 6.5 5.9 3.7 5.8 8.4 (i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the co-culture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.					Tal	ble 2.	.1							
experimental culture 7.5 8.1 7.6 6.2 7.5 7.8 8.9 6.5 7.9 7.3 7.5 control culture 5.6 7.5 8.2 6.7 3.5 6.5 5.9 3.7 5.8 8.4 (i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the co-culture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.					thou	usand	ls of c	ells p	er mr	n ³ of	cultur	е		
(i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the coculture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.	ample	number	1	2	3	4	5	6	7	8	9	10	mean	
 (i) Complete Table 2.1 by calculating the mean number of cells per mm³ in the coculture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) x 100 original number Calculate the percentage increase in the control culture. 	experimental culture		7.5	8.1	7.6	6.2	7.5	7.8	8.9	6.5	7.9	7.3	7.5	
culture. Write your answer in Table 2.1. (ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.	ontrol	culture	5.6	7.5	8.2	6.7	3.5	6.5	5.9	3.7	5.8	8.4		
(ii) A student correctly calculated the percentage increase in the number of cells mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.	(i)		2.1 b	y calc	ulatin	g the	meai	n num	nber o	f cells	s per	mm ³ i	in the co	ont
mm³ in the experimental culture as 151% using the formula: (final number – original number) × 100 original number Calculate the percentage increase in the control culture.		Write your answ	er in	Table	2.1.									
original number Calculate the percentage increase in the control culture.	(ii)													
· · · · · · · · · · · · · · · · · · ·		<u>(1</u>	final r					ber)	× 100					
				age ir	ncrea	se in	the co	ontrol	cultui	re.				

(i) Identify evidence from the results that supports this hypothesis.

(ii) Identify evidence from the results that supports this hypothesis.

(iii) Identify evidence from the results that does not support this hypothesis.

[Total: 12]

For Examiner's Use

© UCLES 2009

QUESTION 3 STARTS ON PAGE 8

For Examiner's Use

3

(a)	ban	ychlorinated biphenyls (PCBs) are persistent organic pollutants. Their use has been ned in many countries. The effects of these pollutants on male fertility has been estigated by many scientists.
	(i)	One group of studies on fish indicates that some of these pollutants decrease the size of the testes in relation to body mass.
		Suggest a procedure by which the relative size of the testes of fish might be estimated.
		[3]
	(ii)	Name a statistical test that is suitable for determining if the decrease in the relative size of the testes is significant.
		Explain your choice.
		test
		explanation[2]
	(iii)	Suggest one way in which the decrease in relative size of the testes may lower fertility of fish.
		[1]

© UCLES 2009 9700/51/O/N/09

(b) Another group of studies tested the effect of one type of PCB, CB-153, on the DNA of human sperm.

For Examiner's Use

In this study, the concentration of CB-153 present in the lipid in the blood plasma of fishermen was measured.

The DNA of a sperm sample was labelled using a fluorescent marker. Undamaged DNA fluoresces green and damaged DNA fluoresces red. The proportion of damaged DNA can be calculated as a DNA fragmentation index.

The data was grouped into six equal sized groups and plotted in relation to the concentration of CB-153 in the lipid in the blood plasma.

Fig. 3.1 shows the results of this study.




Fig. 3.1

CB-153 on the DNA of human sperm.	
	[3]
	[Total: 9]

State the conclusions that can be drawn from this investigation about the effect of

© UCLES 2009 9700/51/O/N/09

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.