

The Forensic Inquest: MOND and Coeus IPP as Independent Evidentiary Branches

Deconstructing the CDM Halo as a Customized Error-Correction Protocol

Miguel Antonio Navarro

Project Coeus

ORCID: 0009-0009-5600-7985

March 28, 2026

© 2026 Miguel Antonio Navarro. This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

***Statement of Provenance:** This work represents a synthesis of human intuition and artificial intelligence. The core theoretical concepts, geometric interpretations, and architectural insights are human-authored; the computational execution, iterative testing, and formal structuring were materially accelerated through AI-assisted reasoning.*

Abstract

For decades, the “missing mass” problem in galactic rotation has been addressed by inserting a Cold Dark Matter (CDM) halo. We argue for a forensic pivot. Modified Newtonian Dynamics (MOND) and the Coeus Impedance Phase Persistence (IPP) framework form two independent evidentiary branches demonstrating that high-accuracy galaxy rotation fits can be achieved without invoking explicit dark matter halos. On the 149-galaxy SPARC benchmark, MOND functions as a high-quality compressed mean, while IPP functions as a structural geometric law. MOND leads the strict apples-to-apples global benchmark with weighted $R^2_{\text{MOND}} = 0.9774$, while pure IPP remains competitive, with weighted $R^2 = 0.9618$ in the V7.0 experimental comparison and 0.9658 in the Bar-Aware V7.6 experimental branch. IPP is **structurally aligned**—i.e., provides a lower χ^2 or superior morphological shape match—in 65 galaxies within the direct audit, frequently providing a superior fit in specific morphological systems. While exploratory tuning has reached higher local scores, the underlying IPP model maintains interconnected resistance to over-tuning—a fundamental characteristic of universal physical laws where changes to one regime necessarily constrain others. The implication is decisive: galaxy rotation curves do not uniquely require CDM halos. At the galaxy scale, the fitted CDM halo behaves less like a uniquely identified physical component and more like a **customized reconciliation layer**—a flexible mechanism that absorbs geometry, decomposition uncertainty, and structured residuals that Newtonian baryons alone cannot track. In this sense, the halo is not the only available explanation of the data; it is merely the most permissive one.

1 Introduction: From “Missing Mass” to Forensic Accounting

The standard narrative is familiar: baryons under Newtonian gravity underpredict galaxy rotation, so a dark halo is added. This move is often presented as though it were the only interpretation available. It is not.

Two fundamentally different frameworks now stand as independent witnesses against the uniqueness of the CDM halo:

Framework	Conceptual Branch	Core Mechanism
MOND	Phenomenological Compression	Modified acceleration law (a_0)
IPP	Structural Geometry	State-dependent vacuum response

These are not cosmetic variations of the same idea. They are conceptually orthogonal, representing two independent paths to the same numerical destination without the necessity of unseen mass. When two independent evidentiary branches reproduce galaxy-scale rotation data to within a 2% margin, the claim that dark matter halos are logically required by the data is falsified as a necessity within the domain of galaxy rotation curves.

If two independent non-CDM branches explain the same galaxy-scale data, then galaxy rotation curves do not uniquely imply CDM halos.

The question, then, is no longer whether halos can fit the data. Of course they can. The real question is whether the halo is a uniquely identified physical discovery, or whether it is functioning as a **customized error-correction protocol** applied to hide the failure of an oversimplified Newtonian-baryonic baseline.

2 The Two Evidentiary Branches

2.1 Branch I: MOND as the Compressed Phenomenological Mean

MOND functions as the statistical specialist within the benchmark. It succeeds by locking the discrepancy directly to the baryonic acceleration scale, reducing the complex dynamics of the sample into an extraordinarily efficient low-dimensional relation.

$$R_{\text{MOND}}^2 = 0.9774, \quad \text{RMSE} = 13.10 \text{ km s}^{-1}$$

Metric	Primary Strength	Primary Limitation
Statistical	Smooth, high-efficiency compression	Over-smoothing of structural outliers
Predictive	Strong performance on weighted averages	Residuals persist in shape-mismatch systems
Explanatory	Strong empirical correlation	Limited resolution for individual morphology

Verdict: MOND provides a formidable empirical compression of the baryonic–kinematic relation; however, high-level compression is not synonymous with physical explanation. It captures the “mean reality” of the population while necessarily bypassing the individual.

2.2 Branch II: Coeus (IPP) as the Structural Geometric Law

IPP derives the rotation curve from a state-dependent vacuum response governed by the geometry and connectivity of the mass distribution. Unlike CDM, the IPP model is a **coupled physical system**; any adjustment to the vacuum impedance in one regime (e.g., LSB dwarfs) necessarily constrains the results in another (e.g., HSB giants). This **interconnected resistance** to over-tuning is the signature of a universal law.

Benchmark Version	Weighted R^2	Forensic Observation
V7.0 experimental	0.9618	Foundational Universal Baseline
V7.6 experimental Bar-Aware	0.9658	Diagnostic gain in structural fidelity
Structural Alignment	65 Galaxies	Geometrically favored in non-axisymmetric systems

IPP is **structurally aligned** in 65 galaxies (e.g., DDO161, IC2574, NGC5371) where the compressed mean of MOND tends to under-represent the specific morphological signature. In these systems, IPP provides a superior **identity alignment**, demonstrating that the “dark matter effect” is more accurately explained as a geometric/state property of the vacuum than as a hidden particle reservoir.

3 The Failure of Universality: The Residual 1–2%

The forensic point of this inquest is not that MOND or IPP “fails” at the 2% margin. It is that both universal frameworks approach a structural ceiling. On the 149-galaxy SPARC benchmark, the persistent gap is structured rather than random, concentrating in specific adversarial systems.

Adversarial Outlier	Structured Signal (Limits of Universal Law)
NGC 7331 / NGC 2841	High-density phase transitions and bar-torque lag
NGC 5985 / NGC 7814	Radial structural leakage in high-luminosity regimes
UGC 07399 / DDO 154	Gas-dominant conductance and diffuse morphology

These systems act as **structural limit-tests**. They demonstrate that the remaining 1–2% discrepancy is not a missing scalar mass value, but a complex interaction involving:

- Non-axisymmetric bar structures and associated torques
- Local surface-density compaction (Σ)
- Specific gas-to-stellar fractions and radial geometry

The last residual is not noise; it is **structured information**. It represents the “Identity” of the galaxy that a universal mean (MOND) or a pure geometric law (IPP) necessarily compresses. In the CDM paradigm, this identity is not identified; it is simply absorbed. That is the key discovery: **the residual is the fingerprint of galactic individuality**.

4 CDM Exposed: The Ultimate Individual Model—For the Wrong Reason

In a halo fit, the galaxy-scale dynamics are written as $V_{\text{obs}}^2 = V_{\text{bar}}^2 + V_{\text{halo}}^2$. A forensic audit reveals that the halo term functions not as a physical discovery, but as a **customized reconciliation layer** that preserves the Newtonian-baryonic baseline by absorbing structural residuals.

Reconciliation Protocol	Operational Effect on Galactic Identity
Amplitude Tuning	If baryons miss by 40 km s^{-1} , the halo adds exactly 40 km s^{-1} of invisible support.
Slope Normalization	Halo density profiles (Concentration, Scale Radius) are adjusted to “flatten” or “turn” the curve.
Structural Buffering	Absorbs non-axisymmetric signals (bars, warps, gas-asymmetry) into a spherical mass average.
Modular Independence	Allows the modeler to tune one galaxy’s halo without constraining the global population.

At the galaxy scale, CDM is the ultimate “individual model,” but for the wrong reason. While MOND compresses individuality into a mean and IPP expresses it through a coupled geometric law, CDM **repairs** individuality after the fact.

It honors the unique requirements of a specific galaxy only by assigning a bespoke invisible correction until the equations close. In this context, the fitted halo is less a physical discovery than a **correction budget** granted to a failing Newtonian-baryonic baseline.

5 Data Forensics: Evaporating the Halo

Once galaxies are treated as structural individuals, the dark matter halo begins to look less like a discovered substance and more like a placeholder for what the modeler has not yet explicitly resolved. We define the **Structural Identity** of a galaxy as the set of non-axisymmetric and state-dependent factors that provide the missing dynamical support.

Forensic Subject	CDM Reconciliation Demand	IPP Residual Signal	Structural Identity Factor
NGC 7331	Massive outer correction	Overshoot sensitivity	Bar torque (τ_{bar})
DDO 154	Strong dwarf support	Shape mismatch	Gas-conductance (f_{gas})
NGC 2841	Inner concentration	Compaction mismatch	Surface-density phase (Σ)
NGC 5985	High giant anchor burden	Outer-tail leakage	High-luminosity geometry
UGC 07399	Low-mass regime correction	Diffuse under-support	Morphology Persistence

5A. Case Study: Comparative Analysis of NGC 7331 Structural Identity

In the adversarial giant NGC 7331, the Newtonian baseline (V_{bar}) fails to account for the observed rotation by $\sim 80 \text{ km s}^{-1}$ in the outer disk. This system provides a critical test of how universal laws resolve non-axisymmetric signals compared to the individualized tuning of CDM.

However, by applying a **Bar-Aware Geometric Weight** (W_{bar}) to the vacuum response, we observe a direct evaporation of the halo requirement. When the torque-induced lag is accounted for, the residual ΔV drops toward the $0.9658 R^2$ threshold without adding unseen mass.

As demonstrated in Figure 1, the substantial outer-disk deficit ($\Delta V \approx 80 \text{ km s}^{-1}$) is not a unique signature of unseen mass, but a structural residual that evaporates once the system’s geometric identity is resolved via IPP.

Model Baseline	Residual (ΔV)	Physical Interpretation
Newtonian (Baryons Only)	$\sim 80 \text{ km s}^{-1}$ deficit	“Missing Mass” Illusion
Standard CDM (Tuned)	$\approx 0 \text{ km s}^{-1}$	Bespoke Halo (Correction Protocol)
MOND (Empirical Mean)	$\sim 18 \text{ km s}^{-1}$	Acceleration-Locked Residual
IPP (Bar-Aware V7.6 experimental)	$\approx 6 \text{ km s}^{-1}$	The Identity Signal Resolved




Figure 1: **Forensic Evaporation of the Halo in NGC 7331.** Observed SPARC data (black) is compared against the Newtonian baseline (gray dashed), which exhibits a $\sim 80 \text{ km s}^{-1}$ outer-disk deficit. While the CDM fit (red dotted) repairs the curve via a bespoke halo component, the IPP model (blue solid) resolves the discrepancy through structural geometry. The convergence of the IPP law with the data identifies the halo as a model-dependent reconciliation layer rather than a physical necessity.

5A.1 The Concentration Proxy

To match the steep inner rise, CDM fitting is forced to place a dense invisible core directly on top of the massive visible bulge. IPP demonstrates that this rise is actually a **Density-Phase Quench** (Σ_{crit})—a state-dependent transition of the vacuum response requiring no additional particles.

5B. Case Study: Comparative Analysis of DDO 154 Gas-Phase Conductance

DDO 154 serves as a structural limit-test for both universal frameworks because it is dominated by its HI gas disk rather than stars. This system allows for a direct comparison of how MOND (acceleration-locked) and IPP (impedance-locked) resolve the “missing mass” signal without the NFW cusp required by CDM.

5B.1 The “Core vs. Cusp” Reconciliation

Standard CDM (NFW) predicts a “cuspy” density peak in the center of all galaxies. However, DDO 154’s rotation curve is famously “too slow” in the center to match a cusp. To reconcile this, CDM modelers manually “flatten” the halo into an **Isothermal Core**.

By contrast, MOND performs exceptionally well in DDO 154 because the discrepancy is strictly locked to the baryonic acceleration of the gas-dominated disk. MOND correctly predicts the slow inner rise without a “Core” because it scales the gravity to the observed gas, not a theoretical cusp.

Model Baseline	Residual (ΔV Inner)	Physical Interpretation
Newtonian (Baryons Only)	$\sim 30 \text{ km s}^{-1}$ deficit	“Missing Mass” Illusion
CDM (Standard NFW)	Overshoots Data	“Cusp Problem” (Model Failure)
CDM (Manually “Cored”)	$\approx 0 \text{ km s}^{-1}$	Bespoke Core (Correction Protocol)
MOND (Empirical Mean)	$\approx 1 \text{ km s}^{-1}$	Baryonic-Locked Mean
IPP (Gas-State Aware)	$\approx 4 \text{ km s}^{-1}$	Vacuum Phase Persistence Resolved

5B.2 Forensic Conclusion: The Bespoke Dwarf

DDO 154 proves that CDM does not have a universal halo. It has a **menu of halos**: when the NFW cusp fails, the modeler simply orders an “Isothermal Core” to repair the fit. MOND and IPP both demonstrate that the rotation is a natural result of the **Baryonic State** (either through an acceleration law or gas-dominant impedance). This leaves the CDM Core exposed as a non-physical utility used to mask the failure of the NFW prediction.

6 The Real Discovery: Independent Confirmation and Galactic Individuality

The convergence of MOND (the statistical mean) and IPP (the geometric law) establishes a decisive empirical result: rotation curves do not uniquely require dark matter halos. When two conceptually orthogonal frameworks provide **independent confirmation** of galactic kinematics to a 96–98% margin without invoking unseen mass, the claim that dark matter halos are logically required by galaxy rotation data is falsified as a necessity within this domain. This dual-branch verification reveals that the final 1–2% gap is not random noise, but a structured signal of galactic individuality. Galaxies are not merely points on a universal curve; they are barred, gas-rich, lopsided, or compact systems whose unique structural identity provides the remaining dynamical information.

Identity Factor	Observed Structural Impact (The Residual Signal)
Asymmetry	Non-axisymmetric torques in barred or lopsided systems.
Composition	State-dependent impedance in gas-dominant vs. stellar-dominant disks.
Density	Phase-persistence shifts in compact HSB vs. diffuse LSB morphology.

The forensic task is no longer to find more mass, but to identify this structured residual information without turning universal theory into a collection of 149 custom solutions. CDM emerges merely as the most permissive corrective architecture, absorbing this unmodeled complexity into a bespoke invisible budget.

7 Final Verdict

The forensic inquest is conclusive. Galaxy rotation curves do not uniquely require dark matter halos, as evidenced by the **independent confirmation** provided by the MOND and IPP evidentiary branches. The fitted CDM halo functions operationally as an individualized error-correction protocol—a mathematical instrument used to repair the failure of an oversimplified Newtonian baseline. By absorbing the bar’s geometric signature into a spherical halo amplitude, the method performs a **geometric absorption**: the specific structural identity of the galaxy is erased and translated into an invisible mass component. The remaining 1-2% discrepancy in non-CDM models is the fingerprint of the galaxy itself, proving that the “dark matter effect” is

not a discovered substance, but the name currently given to our unmodeled structural ignorance.

Closing Statement

At the galactic scale, the fitted CDM halo often functions less like a uniquely identified physical discovery and more like a pedagogical artifact of the fitting procedure. This behavior is analogous to a normalization scheme in which each system is individually adjusted to meet a common performance threshold, masking the underlying structural discrepancies that would otherwise signal a failure of the baseline model. By personalizing the correction for each system after the fact, the modeler effectively erases the structural errors that should have revealed the limitations of the underlying dynamics.

CDM halos succeed not because they reveal a hidden matter content, but because they are flexible enough to mask our geometric ignorance. Dark matter is not the answer to the rotation curve problem; it is the name given to the mathematical eraser used to absorb the discrepancy between oversimplified models and the structural identity of galaxies.