




NUCLEI

CHAPTER Thirteen

CLASS - XII

MANISH JOSHI

D.A.V. LOHAGHAT

September 12, 2025

Learning Objectives:

- Composition and size of nucleus, nuclear force
- Mass-energy relation, mass defect; binding energy per nucleon and its variation with mass number;
- Nuclear Fission, Nuclear Fusion.

Introduction

- In every atom, the **positive charge and mass** are **densely concentrated at the centre of the atom** forming its **nucleus**.

Introduction

- In every atom, the **positive charge and mass** are **densely concentrated at the centre of the atom** forming its **nucleus**.
- the **radius of a nucleus** was smaller than the radius of an atom by a factor of about **10^4** .

Introduction

- In every atom, the **positive charge and mass** are **densely concentrated at the centre of the atom** forming its **nucleus**.
- the **radius of a nucleus** was smaller than the radius of an atom by a factor of about **10^4** .
- This means the **volume of a nucleus** is about **10^{-12}** times the **volume of the atom**. In other words, an **atom is almost empty**.

Introduction

- In every atom, the **positive charge and mass** are **densely concentrated at the centre of the atom** forming its **nucleus**.
- the **radius of a nucleus** was smaller than the radius of an atom by a factor of about **10^4** .
- This means the **volume of a nucleus** is about **10^{-12}** times the **volume of the atom**. In other words, an **atom is almost empty**.
- Nevertheless, the **nucleus contains most** (more than **99.9%**) of the **mass of an atom**.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- The **mass of an atom is very small**, compared to a kilogram; for example, the **mass of a carbon atom, ^{12}C , is 1.992647×10^{-26} kg.**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- The **mass of an atom is very small**, compared to a kilogram; for example, the **mass of a carbon atom, ^{12}C , is 1.992647×10^{-26} kg.**
- **Kilogram is not a very convenient unit** to measure such small quantities.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- The **mass of an atom is very small**, compared to a kilogram; for example, the **mass of a carbon atom, ^{12}C , is 1.992647×10^{-26} kg.**
- **Kilogram is not a very convenient unit** to measure such small quantities.
- Therefore, **atomic mass unit (u)**, is **defined as 1/12th of the mass of the carbon (^{12}C) atom.** According to this definition

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- The **mass of an atom is very small**, compared to a kilogram; for example, the **mass of a carbon atom, ^{12}C , is 1.992647×10^{-26} kg.**
- **Kilogram is not a very convenient unit** to measure such small quantities.
- Therefore, **atomic mass unit (u)**, is **defined as 1/12th of the mass of the carbon (^{12}C) atom.** According to this definition

$$1u = \frac{\text{mass of one } ^{12}\text{C atom}}{12} = \frac{1.992647 \times 10^{-26}}{12} = 1.660539 \times 10^{-27} \text{ kg}$$

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- The **mass of an atom is very small**, compared to a kilogram; for example, the **mass of a carbon atom, ^{12}C , is 1.992647×10^{-26} kg.**
- **Kilogram is not a very convenient unit** to measure such small quantities.
- Therefore, **atomic mass unit (u)**, is **defined as 1/12th of the mass of the carbon (^{12}C) atom.** According to this definition

$$1u = \frac{\text{mass of one } ^{12}\text{C atom}}{12} = \frac{1.992647 \times 10^{-26}}{12} = 1.660539 \times 10^{-27} \text{ kg}$$

- **Accurate measurement of atomic masses** is carried out with a **mass spectrometer.**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- mass of proton = mass of H atom - mass of electron

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- mass of proton = mass of H atom - mass of electron

$$m_p = m_H - m_e = 1.00783 \text{ u} - 0.00055 \text{ u} = 1.00727 \text{ u} = 1.67262 \times 10^{-27} \text{ kg}$$

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- mass of proton = mass of H atom - mass of electron
$$m_p = m_H - m_e = 1.00783 \text{ u} - 0.00055 \text{ u} = 1.00727 \text{ u} = 1.67262 \times 10^{-27} \text{ kg}$$
- It was earlier thought that the nucleus may contain electrons, but this was ruled out later using arguments based on quantum theory. **All the electrons of an atom are outside the nucleus.**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- mass of proton = mass of H atom - mass of electron
$$m_p = m_H - m_e = 1.00783 \text{ u} - 0.00055 \text{ u} = 1.00727 \text{ u} = 1.67262 \times 10^{-27} \text{ kg}$$
- It was earlier thought that the nucleus may contain electrons, but this was ruled out later using arguments based on quantum theory. **All the electrons of an atom are outside the nucleus.**
- The **number of protons** in the nucleus of the atom is **Z, the atomic number.**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- mass of proton = mass of H atom - mass of electron
$$m_p = m_H - m_e = 1.00783 \text{ u} - 0.00055 \text{ u} = 1.00727 \text{ u} = 1.67262 \times 10^{-27} \text{ kg}$$
- It was earlier thought that the nucleus may contain electrons, but this was ruled out later using arguments based on quantum theory. **All the electrons of an atom are outside the nucleus.**
- The **number of protons** in the nucleus of the atom is **Z**, the **atomic number**.
- The **charge of the nucleus** is **(+Ze)**.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Atomic masses:

- mass of proton = mass of H atom - mass of electron
$$m_p = m_H - m_e = 1.00783 \text{ u} - 0.00055 \text{ u} = 1.00727 \text{ u} = 1.67262 \times 10^{-27} \text{ kg}$$
- It was earlier thought that the nucleus may contain electrons, but this was ruled out later using arguments based on quantum theory. **All the electrons of an atom are outside the nucleus.**
- The **number of protons** in the nucleus of the atom is **Z**, the **atomic number**.
- The **charge of the nucleus** is **(+Ze)**.
- since the atom is neutral total **charge of the atomic electrons** is thus **(-Ze)**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Discovery of Neutron:

- Since the **nuclei of deuterium and tritium are isotopes of hydrogen**, they must **contain only one proton** each.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Discovery of Neutron:

- Since the **nuclei of deuterium and tritium are isotopes of hydrogen**, they must **contain only one proton** each. But the **masses of the nuclei of hydrogen, deuterium and tritium** are in the **ratio of 1:2:3**. Therefore, the nuclei of deuterium and tritium **must contain**, in addition to a proton, some **neutral matter**.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Discovery of Neutron:

- Since the **nuclei of deuterium and tritium are isotopes of hydrogen**, they must **contain only one proton** each. But the **masses of the nuclei of hydrogen, deuterium and tritium** are in the **ratio of 1:2:3**. Therefore, the nuclei of deuterium and tritium **must contain**, in addition to a proton, some **neutral matter**.
- This hypothesis was verified in 1932 by **James Chadwick** who **observed emission of neutral radiation** when **beryllium nuclei were bombarded with alpha-particles**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Discovery of Neutron:

- Chadwick assume that the neutral radiation consists of a new type of neutral particles called **neutrons**.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

Discovery of Neutron:

- Chadwick assume that the neutral radiation consists of a new type of neutral particles called **neutrons**. From conservation of energy and momentum, he was able to determine the **mass of new particle as very nearly the same as mass of proton**
- mass of neutron $m_n = 1.00866u = 1.6749 \times 10^{-27} \text{ kg}$

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

ISOTOPES

- **Isotopes:** Atoms of same element (same Z) but different A .

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

ISOTOPES

- **Isotopes:** Atoms of same element (same Z) but different A .
- Example: Hydrogen has three isotopes:



ATOMIC MASSES AND COMPOSITION OF NUCLEUS

ISOTOPES

- **Isotopes:** Atoms of same element (same Z) but different A .
- Example: Hydrogen has three isotopes:



- Chlorine has isotopes $34.98u$ (75.4%) and $36.98u$ (24.6%). Average atomic mass $= 35.47u$.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

ISOTOPES

- **Isotopes:** Atoms of same element (same Z) but different A .
- Example: Hydrogen has three isotopes:



- Chlorine has isotopes $34.98u$ (75.4%) and $36.98u$ (24.6%). Average atomic mass $= 35.47u$.
- Chemical properties of isotopes are identical (same electronic configuration).

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

COMPOSITION OF NUCLEUS

- The composition of a nucleus can now be described using the following terms and symbols:

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

COMPOSITION OF NUCLEUS

- The composition of a nucleus can now be described using the following terms and symbols:
- **Z = atomic number = number of protons**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

COMPOSITION OF NUCLEUS

- The composition of a nucleus can now be described using the following terms and symbols:
- **Z = atomic number = number of protons**
- **N = number of neutrons**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

COMPOSITION OF NUCLEUS

- The composition of a nucleus can now be described using the following terms and symbols:
- **Z = atomic number = number of protons**
- **N = number of neutrons**
- **A = mass number = $Z + N$ = total no. of proton and neutron (or nucleons)**

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

COMPOSITION OF NUCLEUS

- The composition of a nucleus can now be described using the following terms and symbols:
- **Z = atomic number = number of protons**
- **N = number of neutrons**
- **A = mass number = $Z + N$ = total no. of proton and neutron (or nucleons)**
- Nuclear species or nuclides are shown by the notation A_ZX where X is the chemical symbol of the species.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

COMPOSITION OF NUCLEUS

- The composition of a nucleus can now be described using the following terms and symbols:
- **Z = atomic number = number of protons**
- **N = number of neutrons**
- **A = mass number = $Z + N$ = total no. of proton and neutron (or nucleons)**
- Nuclear species or nuclides are shown by the notation A_ZX where X is the chemical symbol of the species.
- For example, the nucleus of gold is denoted by ${}^{197}_{79}Au$. It contains 197 nucleons, of which 79 are protons and the rest 118 are neutrons.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

ISOBARS AND ISOTONES

- **Isobars:** Same mass number (A), different Z . Example: ${}^3_1\text{H}$ and ${}^3_2\text{He}$.

ATOMIC MASSES AND COMPOSITION OF NUCLEUS

ISOBARS AND ISOTONES

- **Isobars:** Same mass number (A), different Z . Example: ${}^3_1\text{H}$ and ${}^3_2\text{He}$.
- **Isotones:** Same neutron number (N), different Z . Example: ${}^{198}_{80}\text{Hg}$ and ${}^{197}_{79}\text{Au}$.

SIZE OF THE NUCLEUS

- Rutherford's suggestion, Geiger and Marsden performed their classic experiment: on the scattering of α -particles from thin gold foils. Their experiments revealed that the distance of closest approach to a gold nucleus of an α -particle of kinetic energy 5.5 MeV is about 4.0×10^{-14} m.

SIZE OF THE NUCLEUS


- Rutherford's suggestion, Geiger and Marsden performed their classic experiment: on the scattering of α -particles from thin gold foils. Their experiments revealed that the distance of closest approach to a gold nucleus of an α -particle of kinetic energy 5.5 MeV is about 4.0×10^{-14} m.
- the **actual size of the nucleus has to be less than 4.0×10^{-14} m.**

SIZE OF THE NUCLEUS

- Rutherford's suggestion, Geiger and Marsden performed their classic experiment: on the scattering of α -particles from thin gold foils. Their experiments revealed that the distance of closest approach to a gold nucleus of an α -particle of kinetic energy 5.5 MeV is about 4.0×10^{-14} m.
- the **actual size of the nucleus has to be less than 4.0×10^{-14} m.**
- By performing scattering experiments in which **fast electrons**, instead of α -particles, are projectiles that bombard targets made up of various elements, the **sizes of nuclei of various elements have been accurately measured.**

SIZE OF THE NUCLEUS


- The simplest model of the nucleus is a densely packed sphere of nucleons.




SIZE OF THE NUCLEUS

- The simplest model of the nucleus is a densely packed sphere of nucleons. **The volume V of the nucleus** is therefore **proportional to the number of nucleons A** , expressed by

$$V = \frac{4}{3}\pi R^3 = kA$$



SIZE OF THE NUCLEUS




- The simplest model of the nucleus is a densely packed sphere of nucleons. **The volume V of the nucleus is therefore proportional to the number of nucleons A , expressed by**

$$V = \frac{4}{3}\pi R^3 = kA$$

- where **R is the radius of a nucleus** and **k is a constant** with units of volume. Solving for R , we have

SIZE OF THE NUCLEUS



- The simplest model of the nucleus is a densely packed sphere of nucleons. **The volume V of the nucleus** is therefore **proportional to the number of nucleons A** , expressed by

$$V = \frac{4}{3}\pi R^3 = kA$$

- where **R is the radius of a nucleus** and **k is a constant** with units of volume. Solving for R , we have

$$R = R_0 A^{1/3}$$

DENSITY OF THE NUCLEUS

- $\text{Density} = \frac{\text{Mass of nucleus}}{\text{Volume of nucleus}} = \frac{A}{\frac{4}{3}\pi R^3} = \frac{A}{\frac{4}{3}\pi R_0^3 A}$

DENSITY OF THE NUCLEUS

- **Density** = $\frac{\text{Mass of nucleus}}{\text{Volume of nucleus}} = \frac{A}{\frac{4}{3}\pi R^3} = \frac{A}{\frac{4}{3}\pi R_0^3 A}$
- **Density** = $\frac{1}{\frac{4}{3}\pi R_0^3} = \text{Constant} = 2.3 \times 10^{17} \text{ kgm}^{-3}$

Example 1

Given the **mass** of iron nucleus as **55.85 u** and **A=56**, find the **nuclear density**?
($1u = 9.27 \times 10^{-26} \text{ kg}$)

Mass – Energy

- **Einstein** showed from his **theory of special relativity** that it is necessary to **treat mass as another form of energy**.

Mass – Energy

- **Einstein** showed from his **theory of special relativity** that it is necessary to **treat mass as another form of energy**.
- **Before** the advent of this **theory of special relativity** it was **presumed that mass and energy were conserved separately** in a reaction.

Mass – Energy

- **Einstein** showed from his **theory of special relativity** that it is necessary to **treat mass as another form of energy**.
- **Before** the advent of this **theory of special relativity** it was **presumed that mass and energy were conserved separately** in a reaction.
- However, **Einstein** showed that **mass is another form of energy** and **one can convert mass-energy into other forms of energy**, say kinetic energy and vice-versa.

Mass – Energy

- **Einstein** showed from his **theory of special relativity** that it is necessary to **treat mass as another form of energy**.
- **Before** the advent of this **theory of special relativity** it was **presumed that mass and energy were conserved separately** in a reaction.
- However, **Einstein** showed that **mass is another form of energy** and **one can convert mass-energy into other forms of energy**, say kinetic energy and vice-versa.
- Einstein gave the famous **mass-energy equivalence relation**

$$E = mc^2$$

Here, m is energy equivalent of mass and c is speed of light in vacuum.

Example 2

Calculate the energy equivalent of 1 g of substance.

Mass Defect

- We have seen that the **nucleus is made up of neutrons and protons.**

Mass Defect

- We have seen that the **nucleus is made up of neutrons and protons.**
- Therefore **it may be expected** that the **mass of the nucleus is equal to the total mass of its individual protons and neutrons.**

Mass Defect

- We have seen that the **nucleus is made up of neutrons and protons.**
- Therefore **it may be expected** that the **mass of the nucleus is equal to the total mass of its individual protons and neutrons.**
- However, According to nuclear particle experiments, the **total mass of a nucleus (M_{nuc}) is less than the sum of the masses of its constituent nucleons (protons and neutrons).**

Mass Defect

- For example, let us consider ${}^1_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.

Mass Defect

- For example, let us consider ${}^{16}_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.
- Mass of 8 neutrons $= 8 \times 1.00866 \text{ } u$

Mass Defect

- For example, let us consider ${}^{16}_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.
- Mass of 8 neutrons $= 8 \times 1.00866 \text{ } u$
- Mass of 8 protons $= 8 \times 1.00727 \text{ } u$

Mass Defect

- For example, let us consider ${}^{16}_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.
- Mass of 8 neutrons $= 8 \times 1.00866 \text{ } u$
- Mass of 8 protons $= 8 \times 1.00727 \text{ } u$
- Mass of 8 electrons $= 8 \times 0.00055 \text{ } u$

Mass Defect

- For example, let us consider ${}^{16}_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.
- Mass of 8 neutrons $= 8 \times 1.00866 \text{ } u$
- Mass of 8 protons $= 8 \times 1.00727 \text{ } u$
- Mass of 8 electrons $= 8 \times 0.00055 \text{ } u$
- Therefore the **expected mass ${}^{16}_8\text{O}$ nucleus** $= 8 \times 2.01593 \text{ } u = 16.12744 \text{ } u$.

Mass Defect

- For example, let us consider ${}^{16}_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.
- Mass of 8 neutrons $= 8 \times 1.00866 \text{ u}$
- Mass of 8 protons $= 8 \times 1.00727 \text{ u}$
- Mass of 8 electrons $= 8 \times 0.00055 \text{ u}$
- Therefore the **expected mass ${}^{16}_8\text{O}$ nucleus** $= 8 \times 2.01593 \text{ u} = 16.12744 \text{ u}$.
- The **atomic mass of ${}^{16}_8\text{O}$ found from mass spectroscopy** experiments is seen to be **15.99493 u**. Subtracting the mass of 8 electrons ($8 \times 0.00055 \text{ u}$) from this, we get the **experimental mass of ${}^{16}_8\text{O}$ nucleus to be 15.99053 u**.

Mass Defect

- For example, let us consider ${}^{16}_8\text{O}$; a nucleus which has **8 neutrons and 8 protons**.
- Mass of 8 neutrons $= 8 \times 1.00866 \text{ u}$
- Mass of 8 protons $= 8 \times 1.00727 \text{ u}$
- Mass of 8 electrons $= 8 \times 0.00055 \text{ u}$
- Therefore the **expected mass ${}^{16}_8\text{O}$ nucleus** $= 8 \times 2.01593 \text{ u} = 16.12744 \text{ u}$.
- The **atomic mass of ${}^{16}_8\text{O}$ found from mass spectroscopy** experiments is seen to be **15.99493 u**. Subtracting the mass of 8 electrons ($8 \times 0.00055 \text{ u}$) from this, we get the **experimental mass of ${}^{16}_8\text{O}$ nucleus to be 15.99053 u**.
- Thus, we find that the **mass of the ${}^{16}_8\text{O}$ nucleus is less than the total mass of its constituents by 0.13691 u**.

Mass Defect

- The **difference in mass of a nucleus and its constituents, ΔM , is called the mass defect**, and is given by

Mass Defect

- The **difference in mass of a nucleus and its constituents, ΔM , is called the mass defect**, and is given by

$$\Delta M = [Z M_p + (A - Z) M_n] - M_{nucl}$$


Mass Defect

- The **difference in mass of a nucleus and its constituents, ΔM , is called the mass defect**, and is given by

$$\Delta M = [Z M_p + (A - Z) M_n] - M_{nucl}$$


- where $Z M_p$ is the **total mass of the protons**, $(A - Z) M_n$ is the **total mass of the neutrons**, and M_{nuc} is the **mass of the nucleus**.

Binding Energy



- The **formation of a nucleus** from a system of isolated protons and neutrons is therefore an **exothermic reaction**-meaning that it **releases energy**. The **energy emitted**, or radiated, in this process is $(\Delta M)c^2$.

Binding Energy



- The **formation of a nucleus** from a system of isolated protons and neutrons is therefore an **exothermic reaction**-meaning that it **releases energy**. The **energy emitted**, or radiated, in this process is $(\Delta M)c^2$.
- Now **imagine this process occurs in reverse**. Instead of forming a nucleus, **energy is put into the system to break apart the nucleus**. The amount of energy required is called the **total binding energy (BE)**, $(Eb = \Delta Mc^2)$.

Example 13.3 Find the energy equivalent of one atomic mass unit, first in Joules and then in MeV. Using this, express the mass defect of $^{16}_8\text{O}$ in MeV/c^2 .

Solution

$$1\text{u} = 1.6605 \times 10^{-27} \text{ kg}$$

To convert it into energy units, we multiply it by c^2 and find that

$$\begin{aligned}\text{energy equivalent} &= 1.6605 \times 10^{-27} \times (2.9979 \times 10^8)^2 \text{ kg m}^2/\text{s}^2 \\ &= 1.4924 \times 10^{-10} \text{ J}\end{aligned}$$

$$= \frac{1.4924 \times 10^{-10}}{1.602 \times 10^{-19}} \text{ eV}$$

$$= 0.9315 \times 10^9 \text{ eV}$$

$$= 931.5 \text{ MeV}$$

$$\text{or, } 1\text{u} = 931.5 \text{ MeV}/c^2$$

$$\begin{aligned}\text{For } ^{16}_8\text{O}, \quad \Delta M &= 0.13691 \text{ u} = 0.13691 \times 931.5 \text{ MeV}/c^2 \\ &= 127.5 \text{ MeV}/c^2\end{aligned}$$

The energy needed to separate $^{16}_8\text{O}$ into its constituents is thus $127.5 \text{ MeV}/c^2$.

Graph of Binding Energy per Nucleon

- In nuclear physics, one of the most important experimental quantities is the binding energy per nucleon (E_{bn}), which is the **ratio of the binding energy** E_b of a nucleus **to the number of the nucleons**, A , in that nucleus:

Graph of Binding Energy per Nucleon

- In nuclear physics, one of the most important experimental quantities is the binding energy per nucleon (E_{bn}), which is the **ratio of the binding energy** E_b of a nucleus **to the number of the nucleons**, A , in that nucleus:

$$E_{bn} = \frac{E_b}{A}$$

Graph of Binding Energy per Nucleon

- In nuclear physics, one of the most important experimental quantities is the binding energy per nucleon (E_{bn}), which is the **ratio of the binding energy** E_b of a nucleus **to the number of the nucleons**, A , in that nucleus:

$$E_{bn} = \frac{E_b}{A}$$

- We can think of **Binding energy per nucleon** as the **average energy per nucleon needed to separate a nucleus into its individual nucleons**.

Graph of Binding Energy per Nucleon

- In nuclear physics, one of the most important experimental quantities is the binding energy per nucleon (E_{bn}), which is the **ratio of the binding energy** E_b of a nucleus **to the number of the nucleons**, A , in that nucleus:

$$E_{bn} = \frac{E_b}{A}$$

- We can think of **Binding energy per nucleon** as the **average energy per nucleon needed to separate a nucleus into its individual nucleons**.
- If the **Binding energy per nucleon** is relatively large, the **nucleus is relatively stable**.






Figure 10.7 In this graph of binding energy per nucleon for stable nuclei, the BEN is greatest for nuclei with a mass near ^{56}Fe . Therefore, fusion of nuclei with mass numbers much less than that of Fe, and fission of nuclei with mass numbers greater than that of Fe, are exothermic processes.

Graph of Binding Energy per Nucleon

- We notice the following **main features of the plot**:




Graph of Binding Energy per Nucleon




- We notice the following **main features of the plot**:
- the **binding energy per nucleon**, E_{bn} , is practically **constant**, i.e. practically independent of the atomic number of middle mass number nuclei ($30 < A < 170$).

Graph of Binding Energy per Nucleon




- We notice the following **main features of the plot**:
- the **binding energy per nucleon**, E_{bn} , is practically **constant**, i.e. practically independent of the atomic number of middle mass number nuclei ($30 < A < 170$). is a consequence of the fact that the **nuclear force is short-ranged**.
- The curve has a **maximum** of about 8.75 MeV **for Fe** $A = 56$ and has a value of 7.6 MeV

Graph of Binding Energy per Nucleon




- We notice the following **main features of the plot**:
- the **binding energy per nucleon**, E_{bn} , is practically **constant**, i.e. practically independent of the atomic number of middle mass number nuclei ($30 < A < 170$). is a consequence of the fact that the **nuclear force is short-ranged**.
- The curve has a **maximum** of about 8.75 MeV **for Fe** $A = 56$ and has a value of 7.6 MeV

Graph of Binding Energy per Nucleon




- typical E_{bn} values range from 6-10 MeV, with an average value of about 8 MeV.

Graph of Binding Energy per Nucleon




- typical E_{bn} values range from 6-10 MeV, with an average value of about 8 MeV.
- In other words, it takes **several million electron volts to pry a nucleon** from a typical nucleus, as compared to just **13.6 eV to ionize an electron** in the ground state of hydrogen.

Graph of Binding Energy per Nucleon




- typical E_{bn} values range from 6-10 MeV, with an average value of about 8 MeV.
- In other words, it takes **several million electron volts to pry a nucleon** from a typical nucleus, as **compared to just 13.6 eV to ionize an electron** in the ground state of hydrogen.
- This is why **nuclear force is referred to as the "strong" nuclear force.**

Graph of Binding Energy per Nucleon




- The graph rises at low A, peaks very near iron (Fe, $A = 56$), and then tapers off at high A.

Graph of Binding Energy per Nucleon




- The graph rises at low A , peaks very near iron (Fe, $A = 56$), and then tapers off at high A .
- The peak value suggests that the iron nucleus is the most stable nucleus in nature (it is also why nuclear fusion in the cores of stars ends with Fe).

Graph of Binding Energy per Nucleon




- The graph rises at low A , peaks very near iron (Fe , $A = 56$), and then tapers off at high A .
- The peak value suggests that the iron nucleus is the most stable nucleus in nature (it is also why nuclear fusion in the cores of stars ends with Fe).
- The reason the graph rises and tapers off has to do with **competing forces in the nucleus**.

Graph of Binding Energy per Nucleon



- At low values of A , attractive nuclear forces between nucleons **dominate over repulsive electrostatic forces** between protons.

Graph of Binding Energy per Nucleon



- At low values of A , attractive nuclear forces between nucleons **dominate over repulsive electrostatic forces** between protons.
- But at **high values of A** , **repulsive electrostatic forces** between forces **begin to dominate**, and **these forces tend to break apart the nucleus** rather than hold it together.

NUCLEAR FORCE

Why a new force is needed?

- Binding energy per nucleon ≈ 8 MeV, much larger than atomic binding energies (eV range).

NUCLEAR FORCE

Why a new force is needed?

- Binding energy per nucleon ≈ 8 MeV, much larger than atomic binding energies (eV range).
- Coulomb repulsion between protons cannot be balanced by gravity (too weak).

NUCLEAR FORCE

Why a new force is needed?

- Binding energy per nucleon ≈ 8 MeV, much larger than atomic binding energies (eV range).
- Coulomb repulsion between protons cannot be balanced by gravity (too weak).
- A new strong attractive force is required to hold nucleons together inside nucleus.

CHARACTERISTICS OF NUCLEAR FORCE

Main Properties:

- **Very Strong:** Stronger than both Coulomb and gravitational forces.

CHARACTERISTICS OF NUCLEAR FORCE

Main Properties:

- **Very Strong:** Stronger than both Coulomb and gravitational forces.
- **Short Range:** Effective only up to a few femtometres (fm); falls rapidly beyond.

CHARACTERISTICS OF NUCLEAR FORCE

Main Properties:

- **Very Strong:** Stronger than both Coulomb and gravitational forces.
- **Short Range:** Effective only up to a few femtometres (fm); falls rapidly beyond.
- **Saturation Property:** Each nucleon interacts only with its nearest neighbours.

CHARACTERISTICS OF NUCLEAR FORCE

Main Properties:

- **Very Strong:** Stronger than both Coulomb and gravitational forces.
- **Short Range:** Effective only up to a few femtometres (fm); falls rapidly beyond.
- **Saturation Property:** Each nucleon interacts only with its nearest neighbours.
- **Charge Independent:** Same for p–p, n–n, and p–n interactions.

CHARACTERISTICS OF NUCLEAR FORCE

Main Properties:

- **Very Strong:** Stronger than both Coulomb and gravitational forces.
- **Short Range:** Effective only up to a few femtometres (fm); falls rapidly beyond.
- **Saturation Property:** Each nucleon interacts only with its nearest neighbours.
- **Charge Independent:** Same for p–p, n–n, and p–n interactions.
- **Attractive–Repulsive Nature:** Attractive beyond 0.8 fm; strongly repulsive below 0.8 fm.

CHARACTERISTICS OF NUCLEAR FORCE

Main Properties:

- **Very Strong:** Stronger than both Coulomb and gravitational forces.
- **Short Range:** Effective only up to a few femtometres (fm); falls rapidly beyond.
- **Saturation Property:** Each nucleon interacts only with its nearest neighbours.
- **Charge Independent:** Same for p–p, n–n, and p–n interactions.
- **Attractive–Repulsive Nature:** Attractive beyond 0.8 fm; strongly repulsive below 0.8 fm.
- **No simple formula:** Unlike Coulomb's or Newton's law, its exact mathematical form is unknown.

POTENTIAL ENERGY CURVE




FIGURE 13.2 Potential energy of a pair of nucleons as a function of their separation.

For a separation greater than r_0 , the force is attractive and for separations less than r_0 , the force is strongly repulsive.

Nuclear Potential Energy vs Distance

- Potential energy minimum at ≈ 0.8 fm $\Rightarrow$ stable bound state.

POTENTIAL ENERGY CURVE




FIGURE 13.2 Potential energy of a pair of nucleons as a function of their separation.

For a separation greater than r_0 , the force is attractive and for separations less than r_0 , the force is strongly repulsive.

Nuclear Potential Energy vs Distance

- Potential energy minimum at ≈ 0.8 fm $\Rightarrow$ stable bound state.
- Attractive for $r > 0.8$ fm.

POTENTIAL ENERGY CURVE




FIGURE 13.2 Potential energy of a pair of nucleons as a function of their separation.

For a separation greater than r_0 , the force is attractive and for separations less than r_0 , the force is strongly repulsive.

Nuclear Potential Energy vs Distance

- Potential energy minimum at ≈ 0.8 fm $\Rightarrow$ stable bound state.
- Attractive for $r > 0.8$ fm.
- Strongly repulsive for $r < 0.8$ fm.

RADIOACTIVITY: Discovery

- **Discovered by Henri Becquerel (1896)** while studying uranium salts.

RADIOACTIVITY: Discovery

- **Discovered by Henri Becquerel (1896)** while studying uranium salts.
- Observed that uranium compounds emitted invisible rays that **darkened photographic plates**, even without sunlight.

RADIOACTIVITY: Discovery

- **Discovered by Henri Becquerel (1896)** while studying uranium salts.
- Observed that uranium compounds emitted invisible rays that **darkened photographic plates**, even without sunlight.
- This phenomenon was later named **Radioactivity**.

RADIOACTIVITY: Definition

- **Radioactivity:** Spontaneous decay of an unstable nucleus with emission of particles or radiation.

RADIOACTIVITY: Definition

- **Radioactivity:** Spontaneous decay of an unstable nucleus with emission of particles or radiation.
- It is a **nuclear phenomenon**, not influenced by temperature, pressure, or chemical state.

RADIOACTIVITY: Definition

- **Radioactivity:** Spontaneous decay of an unstable nucleus with emission of particles or radiation.
- It is a **nuclear phenomenon**, not influenced by temperature, pressure, or chemical state.
- Governed by **probability laws** (exponential decay).

Alpha (α) Decay


- Emission of a helium nucleus (${}^4_2\text{He}$).

Alpha (α) Decay

- Emission of a helium nucleus (${}^4_2\text{He}$).
- Reduces **mass number (A)** by 4 and **atomic number (Z)** by 2.

Alpha (α) Decay

- Emission of a helium nucleus (${}^4_2\text{He}$).
- Reduces **mass number (A)** by 4 and **atomic number (Z)** by 2.
- **Example:**



Beta (β) Decay


- Emission of an **electron** (β^-) or a **positron** (β^+).

Beta (β) Decay

- Emission of an **electron** (β^-) or a **positron** (β^+).
- **In β^- decay:** neutron $\rightarrow$ proton + electron + antineutrino.
Atomic number increases by 1.

Beta (β) Decay


- Emission of an **electron** (β^-) or a **positron** (β^+).
- **In β^- decay:** neutron $\rightarrow$ proton + electron + antineutrino.
Atomic number increases by 1.
- **Example:**



Beta (β) Decay

- Emission of an **electron** (β^-) or a **positron** (β^+).
- **In β^- decay:** neutron $\rightarrow$ proton + electron + antineutrino.
Atomic number increases by 1.

- **Example:**




- **In β^+ decay:** proton $\rightarrow$ neutron + positron + neutrino.
Atomic number decreases by 1.

Beta (β) Decay


- Emission of an **electron** (β^-) or a **positron** (β^+).
- **In β^- decay:** neutron $\rightarrow$ proton + electron + antineutrino.
Atomic number increases by 1.

- **Example:**



- **In β^+ decay:** proton $\rightarrow$ neutron + positron + neutrino.
Atomic number decreases by 1.

- **Example:**



Gamma (γ) Decay


- Emission of **high-energy photons (γ -rays)**.

Gamma (γ) Decay

- Emission of **high-energy photons (γ -rays)**.
- No change in **A or Z**. Only excess energy is released.

Gamma (γ) Decay

- Emission of **high-energy photons (γ -rays)**.
- No change in **A or Z**. Only excess energy is released.
- **Example:**



(* denotes excited nucleus).

Nuclear Energy: Binding Energy and Stability

- **Binding Energy (B.E.):** Energy needed to break a nucleus into free protons and neutrons.

Nuclear Energy: Binding Energy and Stability

- **Binding Energy (B.E.):** Energy needed to break a nucleus into free protons and neutrons.
- **Binding Energy per nucleon (B.E./A):** **Stability measure** of a nucleus.

Nuclear Energy: Binding Energy and Stability

- **Binding Energy (B.E.):** Energy needed to break a nucleus into free protons and neutrons.
- **Binding Energy per nucleon (B.E./A):** Stability measure of a nucleus.
- **Binding energy curve:**
 - Peaks at **Iron (Fe, A = 56)** → the **most stable nucleus**.
 - Lower for **very light (A < 30)** and **very heavy (A > 170)** nuclei.

Nuclear Energy: Binding Energy and Stability

- **Binding Energy (B.E.):** Energy needed to break a nucleus into free protons and neutrons.
- **Binding Energy per nucleon (B.E./A):** Stability measure of a nucleus.
- **Binding energy curve:**
 - Peaks at **Iron (Fe, A = 56)** → the **most stable nucleus**.
 - Lower for **very light (A < 30)** and **very heavy (A > 170)** nuclei.
- **Implication:**
 - **Heavy nuclei** release energy when split (**fission**).
 - **Light nuclei** release energy when combined (**fusion**).

Nuclear Fission




Figure 10.19 In the liquid drop model of nuclear fission, the uranium nucleus is split into two lighter nuclei by a high-energy neutron.




Figure 10.20 In a U-235 fission chain reaction, the fission of the uranium nucleus produces high-energy neutrons that go on to split more nuclei. The energy released in this process can be used to produce electricity.

Nuclear Fission



- **Definition:** Splitting of a **heavy nucleus** into two **medium-mass nuclei** with emission of **neutrons and large energy**.

Nuclear Fission

- **Definition:** Splitting of a **heavy nucleus** into two **medium-mass nuclei** with emission of **neutrons and large energy**.
- **Discovered:** Hahn and Strassmann (1939).



Nuclear Fission

- **Definition:** Splitting of a **heavy nucleus** into two **medium-mass nuclei** with emission of **neutrons and large energy**.
- **Discovered:** Hahn and Strassmann (1939).
- **Fission Reactions:**



Nuclear Fission

- **Definition:** Splitting of a **heavy nucleus** into two **medium-mass nuclei** with emission of **neutrons and large energy**.
- **Discovered:** Hahn and Strassmann (1939).
- **Fission Reactions:**



- **Energy released per fission** 200 MeV.

Features of Fission

- Fission triggered by absorption of a **slow neutron (thermal neutron)**.

Features of Fission

- Fission triggered by absorption of a **slow neutron (thermal neutron)**.
- Produces **fission fragments** which are usually **radioactive**.

Features of Fission

- Fission triggered by absorption of a **slow neutron (thermal neutron)**.
- Produces **fission fragments** which are usually **radioactive**.
- Emission of **2–3 neutrons** → can sustain a **chain reaction**.

Features of Fission

- Fission triggered by absorption of a **slow neutron (thermal neutron)**.
- Produces **fission fragments** which are usually **radioactive**.
- Emission of **2–3 neutrons** → can sustain a **chain reaction**.
- **Applications:**
 - **Controlled chain reaction:** Nuclear reactor (electricity).
 - **Uncontrolled chain reaction:** Atom bomb.

Nuclear Fusion

- **Definition:** Fusion is the process of two **light nuclei** combining to form a **heavier nucleus**, releasing energy.

Nuclear Fusion

- **Definition:** Fusion is the process of two **light nuclei** combining to form a **heavier nucleus**, releasing energy.
- Fusion reactions occur naturally in **stars (like the Sun)**.

Nuclear Fusion

- **Definition:** Fusion is the process of two **light nuclei** combining to form a **heavier nucleus**, releasing energy.
- Fusion reactions occur naturally in **stars (like the Sun)**.
- For fusion to take place, the two nuclei must come close enough so that attractive **short-range nuclear force** is able to affect them.

Nuclear Fusion

- **Definition:** Fusion is the process of two **light nuclei** combining to form a **heavier nucleus**, releasing energy.
- Fusion reactions occur naturally in **stars (like the Sun)**.
- For fusion to take place, the two nuclei must come close enough so that attractive **short-range nuclear force** is able to affect them.
- However, since they are both **positively charged**, they experience **Coulomb repulsion**. They, therefore, must have enough energy to **overcome the Coulomb barrier**.

Nuclear Fusion





- **Definition:** Fusion is the process of two **light nuclei** combining to form a **heavier nucleus**, releasing energy.
- Fusion reactions occur naturally in **stars (like the Sun)**.
- For fusion to take place, the two nuclei must come close enough so that attractive **short-range nuclear force** is able to affect them.
- However, since they are both **positively charged**, they experience **Coulomb repulsion**. They, therefore, must have enough energy to **overcome the Coulomb barrier**.
- Fusion requires **extremely high temperature** ($10^7 - 10^8 \text{ K}$) $\rightarrow$ **thermonuclear reaction**.

Nuclear Fusion


- **Definition:** Fusion is the process of two **light nuclei** combining to form a **heavier nucleus**, releasing energy.
- Fusion reactions occur naturally in **stars (like the Sun)**.
- For fusion to take place, the two nuclei must come close enough so that attractive **short-range nuclear force** is able to affect them.
- However, since they are both **positively charged**, they experience **Coulomb repulsion**. They, therefore, must have enough energy to **overcome the Coulomb barrier**.
- Fusion requires **extremely high temperature** ($10^7 - 10^8 \text{ K}$) $\rightarrow$ **thermonuclear reaction**.

Nuclear Fusion Reactions

- **Proton–Proton Chain (in the Sun):**



Net reaction: $2(i) + 2(ii) + 2(iii) + (iv)$



- Here $\nu \rightarrow$ a neutrino, $e^+ \rightarrow$ positron, $e^- \rightarrow$ electron and $\gamma \rightarrow$ gamma rays.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.
- When **hydrogen is depleted**, the core cools and collapses under gravity.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.
- When **hydrogen is depleted**, the core cools and collapses under gravity.
- Collapse raises the **core temperature**.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.
- When **hydrogen is depleted**, the core cools and collapses under gravity.
- Collapse raises the **core temperature**.
- At about 10^8 **K**, helium nuclei fuse into **carbon**.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.
- When **hydrogen is depleted**, the core cools and collapses under gravity.
- Collapse raises the **core temperature**.
- At about 10^8 **K**, helium nuclei fuse into **carbon**.
- Fusion can produce heavier elements up to the **binding energy peak (iron)**.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.
- When **hydrogen is depleted**, the core cools and collapses under gravity.
- Collapse raises the **core temperature**.
- At about 10^8 **K**, helium nuclei fuse into **carbon**.
- Fusion can produce heavier elements up to the **binding energy peak (iron)**.
- Elements more massive than iron **cannot be produced by fusion**.

Nuclear Fusion Reactions

- Hydrogen nuclei fuse into **helium** in the core of stars.
- When **hydrogen is depleted**, the core cools and collapses under gravity.
- Collapse raises the **core temperature**.
- At about 10^8 **K**, helium nuclei fuse into **carbon**.
- Fusion can produce heavier elements up to the **binding energy peak (iron)**.
- Elements more massive than iron **cannot be produced by fusion**.

Nuclear Fusion Reactions

- Age of the Sun: 5×10^9 **years**.

Nuclear Fusion Reactions

- Age of the Sun: 5×10^9 **years**.
- Hydrogen fuel will last another **5 billion years**.

Nuclear Fusion Reactions

- Age of the Sun: 5×10^9 **years**.
- Hydrogen fuel will last another **5 billion years**.
- After hydrogen burning stops, the **core contracts and heats up**.

Nuclear Fusion Reactions


- Age of the Sun: 5×10^9 **years**.
- Hydrogen fuel will last another **5 billion years**.
- After hydrogen burning stops, the **core contracts and heats up**.
- The outer envelope of the Sun will **expand**.

Nuclear Fusion Reactions

- Age of the Sun: 5×10^9 **years**.
- Hydrogen fuel will last another **5 billion years**.
- After hydrogen burning stops, the **core contracts and heats up**.
- The outer envelope of the Sun will **expand**.
- The Sun will turn into a **Red Giant**.

Nuclear Fusion Reactions

- Age of the Sun: 5×10^9 **years**.
- Hydrogen fuel will last another **5 billion years**.
- After hydrogen burning stops, the **core contracts and heats up**.
- The outer envelope of the Sun will **expand**.
- The Sun will turn into a **Red Giant**.
- **Deuterium–Tritium Fusion (on Earth, proposed):**



Controlled Thermonuclear Fusion

- **In Stars:** High pressure + very high temperature $\rightarrow$ fusion is sustained naturally.

Controlled Thermonuclear Fusion

- **In Stars:** High pressure + very high temperature $\rightarrow$ fusion is sustained naturally.
- **On Earth:** We try to reproduce this in reactors.

Controlled Thermonuclear Fusion

- **In Stars:** High pressure + very high temperature → fusion is sustained naturally.
- **On Earth:** We try to reproduce this in reactors.
- **Requirements:**
 - Temperature of $\approx 10^8$ K to **overcome Coulomb repulsion**.
 - At this temperature, fuel becomes **plasma (ionised gas)**.
 - Plasma must be **confined** so it does not touch container walls, since **no container can withstand such high temperature**.

Controlled Thermonuclear Fusion

- **In Stars:** High pressure + very high temperature → fusion is sustained naturally.
- **On Earth:** We try to reproduce this in reactors.
- **Requirements:**
 - Temperature of $\approx 10^8$ K to **overcome Coulomb repulsion**.
 - At this temperature, fuel becomes **plasma (ionised gas)**.
 - Plasma must be **confined** so it does not touch container walls, since **no container can withstand such high temperature**.
- **Methods of Confinement:**
 - **Magnetic confinement** → in toroidal devices (**tokamaks**).
 - **Inertial confinement** → using powerful **lasers** to compress fuel.

Controlled Thermonuclear Fusion

- **In Stars:** High pressure + very high temperature → fusion is sustained naturally.
- **On Earth:** We try to reproduce this in reactors.
- **Requirements:**
 - Temperature of $\approx 10^8$ K to **overcome Coulomb repulsion**.
 - At this temperature, fuel becomes **plasma (ionised gas)**.
 - Plasma must be **confined** so it does not touch container walls, since **no container can withstand such high temperature**.
- **Methods of Confinement:**
 - **Magnetic confinement** → in toroidal devices (**tokamaks**).
 - **Inertial confinement** → using powerful **lasers** to compress fuel.
- Fusion is attractive because **fuel (deuterium from water) is abundant** and **waste is less hazardous** compared to fission.