

Lorentz Force

- Consider a **point charge** q **moving with a velocity** $\vec{v}$ in the **presence of both electric field** $\vec{E}$ and **magnetic field** $\vec{B}$
- The moving **charge experience two force**:

Lorentz Force

- Consider a **point charge** q **moving with a velocity** $\vec{v}$ in the **presence of both electric field** $\vec{E}$ and **magnetic field** $\vec{B}$
- The moving **charge experience two force**:
 - ① **Electric Force**: $\vec{F}_e = q\vec{E}$ and,

Lorentz Force

- Consider a **point charge** q **moving with a velocity** $\vec{v}$ in the **presence of both electric field** $\vec{E}$ and **magnetic field** $\vec{B}$
- The moving **charge experience two force**:
 - ① **Electric Force**: $\vec{F}_e = q\vec{E}$ and,
 - ② **Magnetic Force**: $\vec{F}_m = q\vec{V} \times \vec{B}$

Lorentz Force

- Consider a **point charge** q **moving with a velocity** $\vec{v}$ in the **presence of both electric field** $\vec{E}$ and **magnetic field** $\vec{B}$
- The moving **charge experience two force**:
 - ① **Electric Force**: $\vec{F}_e = q\vec{E}$ and,
 - ② **Magnetic Force**: $\vec{F}_m = q\vec{v} \times \vec{B}$
- The **net force** on the moving charge **is called Lorentz force** F_L and is given by

$$\vec{F}_L = \vec{F}_e + \vec{F}_m$$

Lorentz Force

- Consider a **point charge** q **moving with a velocity** $\vec{v}$ in the **presence of both electric field** $\vec{E}$ and **magnetic field** $\vec{B}$
- The moving **charge experience two force**:
 - ① **Electric Force**: $\vec{F}_e = q\vec{E}$ and,
 - ② **Magnetic Force**: $\vec{F}_m = q\vec{v} \times \vec{B}$
- The **net force** on the moving charge **is called Lorentz force** F_L and is given by

$$\begin{aligned}\vec{F}_L &= \vec{F}_e + \vec{F}_m \\ \vec{F}_L &= q\vec{E} + q\vec{v} \times \vec{B}\end{aligned}$$


Lorentz Force

- Consider a **point charge** q **moving with a velocity** $\vec{v}$ in the **presence of both electric field** $\vec{E}$ and **magnetic field** $\vec{B}$
- The moving **charge experience two force**:
 - ① **Electric Force**: $\vec{F}_e = q\vec{E}$ and,
 - ② **Magnetic Force**: $\vec{F}_m = q\vec{v} \times \vec{B}$
- The **net force** on the moving charge **is called Lorentz force** F_L and is given by

$$\vec{F}_L = \vec{F}_e + \vec{F}_m$$
$$\vec{F}_L = q\vec{E} + q\vec{v} \times \vec{B}$$


$$\boxed{\vec{F}_L = q \left(\vec{E} + \vec{v} \times \vec{B} \right)}$$

Lorentz Force




- Force due to magnetic field depends on q , $\vec{v}$ and $\vec{B}$.

Lorentz Force




- Force due to magnetic field depends on q , $\vec{v}$ and $\vec{B}$.
- Value of magnetic force will be zero if

Lorentz Force




- Force due to magnetic field depends on q , $\vec{v}$ and $\vec{B}$.
- Value of magnetic force will be zero if
 - 1 Charge is at rest $v = 0$

Lorentz Force




- Force due to magnetic field depends on q , $\vec{v}$ and $\vec{B}$.
- Value of magnetic force will be zero if
 - 1 Charge is at rest $v = 0$
 - 2 $\vec{v}$ is parallel to $\vec{B}$ (i.e., $\sin(\theta) = 0$)

Lorentz Force



- **Force due to magnetic field depends on q , $\vec{v}$ and $\vec{B}$.**
- **Value of magnetic force will be zero if**
 - 1 **Charge is at rest $v = 0$**
 - 2 **$\vec{v}$ is parallel to $\vec{B}$ (i.e., $\sin(\theta) = 0$)**
 - 3 **$q = 0$ Or $B = 0$**

Lorentz Force



- **Force due to magnetic field depends on q , $\vec{v}$ and $\vec{B}$.**
- **Value of magnetic force will be zero if**
 - 1 **Charge is at rest $v = 0$**
 - 2 **$\vec{v}$ is parallel to $\vec{B}$ (i.e., $\sin(\theta) = 0$)**
 - 3 **$q = 0$ Or $B = 0$**
- **Magnetic Lorentz force $\vec{F}_m$ will always perpendicular to both $\vec{v}$ and $\vec{B}$ and its direction is given by **Right hand thumb rule** for vector product.**

Lorentz Force

- **Magnetic Lorentz force help us to define the unit of magnetic field**

Lorentz Force

- **Magnetic Lorentz force help us to define the unit of magnetic field**

$$B = \frac{F}{qv \sin \theta}$$

Lorentz Force

- **Magnetic Lorentz force help us to define the unit of magnetic field**

$$B = \frac{F}{qv \sin \theta}$$

The S.I unit of magnetic field is Tesla (T)

Lorentz Force

- **Magnetic Lorentz force help us to define the unit of magnetic field**


$$B = \frac{F}{qv \sin \theta}$$

The S.I unit of magnetic field is Tesla (T)

$$1T = \frac{N \ s}{C \ m}$$


- **A smaller unit, called the gauss (G), where $1G=10^{-4} \ T$, is sometimes used.**

Motion of a Charged Particle in a Magnetic Field




- Consider the **motion of a charge particle** in **uniform magnetic field** $\vec{B}$

Motion of a Charged Particle in a Magnetic Field




- Consider the **motion of a charge particle** in **uniform magnetic field** $\vec{B}$
- The magnetic **force is always perpendicular to velocity**, so that it does **no work on the charged particle** ($dW = \vec{F} \cdot d\vec{r} = 0$).

Motion of a Charged Particle in a Magnetic Field




- Consider the **motion of a charge particle** in **uniform magnetic field** $\vec{B}$
- The magnetic **force is always perpendicular to velocity**, so that it does **no work on the charged particle** ($dW = \vec{F} \cdot d\vec{r} = 0$).
- The particle's **kinetic energy and speed thus remain constant**.

Motion of a Charged Particle in a Magnetic Field




- Consider the **motion of a charge particle** in **uniform magnetic field** $\vec{B}$
- The magnetic **force is always perpendicular to velocity**, so that it does **no work on the charged particle** ($dW = \vec{F} \cdot d\vec{r} = 0$).
- The particle's **kinetic energy and speed thus remain constant**.
- The **direction of motion is affected but not the speed**.

Motion of a Charged Particle in a Magnetic Field



Case 1: When $\vec{v}$ perpendicular to $\vec{B}$


Motion of a Charged Particle in a Magnetic Field



Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- **Force acting on charge particle (q) is:**

Motion of a Charged Particle in a Magnetic Field




Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- Force acting on charge particle (q) is:

$$\vec{F} = q(\vec{v} \times \vec{B})$$

Motion of a Charged Particle in a Magnetic Field




Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- **Force acting on charge particle (q) is:**

$$\vec{F} = q(\vec{v} \times \vec{B})$$

- This magnetic force qvB acts as **centripetal force** $\frac{mv^2}{r}$ and produces a circular motion perpendicular to the magnetic field.

Motion of a Charged Particle in a Magnetic Field



Case 1: When $\vec{v}$ perpendicular to $\vec{B}$


- **Force acting on charge particle (q) is:**

$$\vec{F} = q(\vec{v} \times \vec{B})$$

- This magnetic force qvB acts as **centripetal force** $\frac{mv^2}{r}$ and produces a circular motion perpendicular to the magnetic field.
- Here, r be the radius of the circle, then

$$\frac{mv^2}{r} = qvB$$

Motion of a Charged Particle in a Magnetic Field



Case 1: When $\vec{v}$ perpendicular to $\vec{B}$


- Force acting on charge particle (q) is:

$$\vec{F} = q(\vec{v} \times \vec{B})$$

- This magnetic force qvB acts as **centripetal force** $\frac{mv^2}{r}$ and produces a circular motion perpendicular to the magnetic field.
- Here, r be the radius of the circle, then

$$\frac{mv^2}{r} = qvB \text{ Or } \boxed{r = \frac{mv}{qB}}$$


Motion of a Charged Particle in a Magnetic Field



Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- The time taken for one revolution is

Motion of a Charged Particle in a Magnetic Field




Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- The time taken for one revolution is

$$T = \frac{2\pi r}{v} = \frac{2\pi m}{qB}$$

Motion of a Charged Particle in a Magnetic Field




Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- The time taken for one revolution is

$$T = \frac{2\pi r}{v} = \frac{2\pi m}{qB}$$

- If ω is the angular frequency, then

Motion of a Charged Particle in a Magnetic Field



Case 1: When $\vec{v}$ perpendicular to $\vec{B}$

- The time taken for one revolution is

$$T = \frac{2\pi r}{v} = \frac{2\pi m}{qB}$$

- If ω is the angular frequency, then

$$\omega = \frac{2\pi}{T} = \frac{qB}{m}$$

Motion of a Charged Particle in a Magnetic Field

Case 2: If velocity $\vec{v}$ has a component along $\vec{B}$




FIGURE 4.6 Helical motion

Motion of a Charged Particle in a Magnetic Field




FIGURE 4.6 Helical motion

Case 2: If velocity $\vec{v}$ has a component along $\vec{B}$

- In this case the **component of velocity along $\vec{B}$** (*i.e.*, $v_{\parallel}$) make the particle move along the field and the **path of the particle** would be a **helical**.

Motion of a Charged Particle in a Magnetic Field




FIGURE 4.6 Helical motion

Case 2: If velocity $\vec{v}$ has a component along $\vec{B}$

- In this case the **component of velocity along $\vec{B}$** (*i.e.*, $v_\parallel$) make the particle move along the field and the **path of the particle** would be a **helical**.
- The **distance moved** along the magnetic field in **one rotation** is called **pitch p**.

Motion of a Charged Particle in a Magnetic Field




FIGURE 4.6 Helical motion

Case 2: If velocity $\vec{v}$ has a component along $\vec{B}$

- In this case the **component of velocity along $\vec{B}$** (*i.e.*, $v_{\parallel}$) make the particle move along the field and the **path of the particle** would be a **helical**.
- The **distance moved** along the magnetic field in **one rotation** is called **pitch p**.

$$p = v_{\parallel} \times T = v \cos \theta \times \frac{2\pi m}{qB}$$

Motion of a Charged Particle in a Magnetic Field




FIGURE 4.6 Helical motion

Case 2: If velocity $\vec{v}$ has a component along $\vec{B}$

- In this case the **component of velocity along $\vec{B}$** (*i.e.*, $v_{\parallel}$) make the particle move along the field and the **path of the particle** would be a **helical**.
- The **distance moved** along the magnetic field in **one rotation** is called **pitch p**.

$$p = v_{\parallel} \times T = v \cos \theta \times \frac{2\pi m}{qB}$$

$$p = \frac{2\pi m v \cos \theta}{qB}$$

Example 4.3

What is the radius of the path of an electron (mass $9 \times 10^{-31} \text{ kg}$ and charge $1.6 \times 10^{-19} \text{ C}$) moving at a speed of $3 \times 10^7 \text{ m/s}$ in a magnetic field of $6 \times 10^{-4} \text{ T}$ perpendicular to it? What is its frequency? Calculate its energy in keV. ($1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$).

Exercise 4.11

In a chamber, a uniform magnetic field of 6.5 G ($1 \text{ G} = 10^{-4} \text{ T}$) is maintained. An electron is shot into the field with a speed of $4.8 \times 10^6 \text{ ms}^{-1}$ normal to the field. Explain why the path of the electron is a circle. Determine the radius of the circular orbit. ($e = 1.5 \times 10^{-19} \text{ C}$, $m_e = 9.1 \times 10^{-31} \text{ kg}$)

Exercise 4.12

In Exercise 4.11 obtain the frequency of revolution of the electron in its circular orbit. Does the answer depend on the speed of the electron? Explain.

Reference Example

A proton enters a uniform magnetic field of $1.0 \times 10^{-4} \text{ T}$ with a speed of $5 \times 10^5 \text{ m/s}$. At what angle must the magnetic field be from the velocity so that the pitch of the resulting helical motion is equal to the radius of the helix?