

Deep Learning

Vazgen Mikayelyan



1 What is Convolutional Neural Network?

What is convolution?

Definition 1

Convolution of the functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ is defined as the integral of the product of the two functions after one is reversed and shifted:

$$(f * g)(t) =: \int_{-\infty}^{+\infty} f(x) g(t - x) dx.$$

What is convolution?

Definition 1

Convolution of the functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ is defined as the integral of the product of the two functions after one is reversed and shifted:

$$(f * g)(t) =: \int_{-\infty}^{+\infty} f(x) g(t - x) dx.$$

It easy to see that $f * g = g * f$.

What is convolution?

Definition 1

Convolution of the functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ is defined as the integral of the product of the two functions after one is reversed and shifted:

$$(f * g)(t) =: \int_{-\infty}^{+\infty} f(x) g(t-x) dx.$$

It easy to see that $f * g = g * f$.

Definition 2

Convolution of the sequences of real numbers $\{f_n\}_{n=-\infty}^{+\infty}$, $\{g_n\}_{n=-\infty}^{+\infty}$ is the following sequence:

$$z_n =: \sum_{k=-\infty}^{+\infty} f_k g_{n-k}.$$

Definition 3

Convolution of the functions $f, g : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the following function:

$$(f * g)(t, \tau) =: \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(x, y) g(t - x, \tau - y) dx dy.$$

Definition 3

Convolution of the functions $f, g : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the following function:

$$(f * g)(t, \tau) =: \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(x, y) g(t - x, \tau - y) dx dy.$$

It easy to see that $f * g = g * f$.

2D Convolution

Definition 3

Convolution of the functions $f, g : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the following function:

$$(f * g)(t, \tau) =: \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} f(x, y) g(t - x, \tau - y) dx dy.$$


It easy to see that $f * g = g * f$.

Definition 4

Let $f(x, y)$ is an image and $w(s, t)$ is a kernel where $s \in [-a, a], t \in [-b, b], x, y, s, t, a, b \in \mathbb{Z}$. The convolution between kernel w and image f is the following function


$$(w * f)(x, y) = \sum_{s=-a}^a \sum_{t=-b}^b w(s, t) f(x - s, y - t)$$

2D Convolution




Convolution on RGB Images

Convolution on RGB Images




Convolution on RGB Images



- How many dimensions has convolution kernel in general?

Convolution on RGB Images



- How many dimensions has convolution kernel in general?
- What does 1×1 convolution do?