

Mathematics for Machine Learning

Vazgen Mikayelyan



WELCOME TO

Mathematics for Machine Learning!

- Course: **Mathematics for Machine Learning**
- Instructor: Vazgen Mikayelyan

Course Content

- Linear Algebra
- Analytical Geometry
- Mathematical Analysis
- Probability Theory

Find other interesting stuff in the bibliography!

Definition

With $n \in \mathbb{N}$ a real-valued n vector a is a "sequence" of n elements a_i , $i = 1, \dots, n$:

$$a = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}, a_i \in \mathbb{R}.$$

Definition

With $n \in \mathbb{N}$ a real-valued n vector a is a "sequence" of n elements a_i , $i = 1, \dots, n$:

$$a = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}, a_i \in \mathbb{R}.$$

$\mathbb{R}^n$ is the set of all real-valued n -vectors.

Sum of Two Vectors

If a and b are vectors with n elements, then their sum $a + b$ is the following vector:

$$a + b = \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \\ \vdots \\ a_n + b_n \end{bmatrix}$$

Multiplication by a Scalar

If a is a vector with n elements and λ is a real scalar, then the scalar multiple λa is the following vector:

$$\lambda a = \begin{bmatrix} \lambda a_1 \\ \lambda a_2 \\ \vdots \\ \lambda a_n \end{bmatrix}$$

- **Distributivity:**

$$(\lambda_1 + \lambda_2)a = \lambda_1 a + \lambda_2 a, \quad a \in \mathbb{R}^n, \lambda_1, \lambda_2 \in \mathbb{R},$$

$$\lambda(a + b) = \lambda a + \lambda b, \quad a, b \in \mathbb{R}^n, \lambda \in \mathbb{R}.$$

- **Distributivity:**

$$(\lambda_1 + \lambda_2)a = \lambda_1 a + \lambda_2 a, \quad a \in \mathbb{R}^n, \lambda_1, \lambda_2 \in \mathbb{R},$$

$$\lambda(a + b) = \lambda a + \lambda b, \quad a, b \in \mathbb{R}^n, \lambda \in \mathbb{R}.$$

- **Associativity:**

$$(a + b) + c = a + (b + c), \quad a, b, c \in \mathbb{R}^n.$$

Matrices

Definition

With $m, n \in \mathbb{N}$ a real-valued (m, n) matrix A is matrix an $m \cdot n$ -tuple of elements a_{ij} , $i = 1, \dots, m$, $j = 1, \dots, n$, which is ordered in m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, a_{ij} \in \mathbb{R}.$$

$(1, n)$ -matrices are called **rows**, $(m, 1)$ -matrices are called **columns**. These special matrices are also called **row/column vectors**.

Definition

With $m, n \in \mathbb{N}$ a real-valued (m, n) matrix A is matrix an $m \cdot n$ -tuple of elements a_{ij} , $i = 1, \dots, m$, $j = 1, \dots, n$, which is ordered in m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, a_{ij} \in \mathbb{R}.$$

$(1, n)$ -matrices are called **rows**, $(m, 1)$ -matrices are called **columns**. These special matrices are also called **row/column vectors**.

$\mathbb{R}^{m \times n}$ is the set of all real-valued (m, n) -matrices.

Definition

If A and B are $m \times n$ matrices, then the sum $C = A + B$ is an $m \times n$ matrix and the (i, j) entry of the sum is computed as follows:

$$c_{ij} = a_{ij} + b_{ij}.$$

Matrix Addition

Definition

If A and B are $m \times n$ matrices, then the sum $C = A + B$ is an $m \times n$ matrix and the (i, j) entry of the sum is computed as follows:

$$c_{ij} = a_{ij} + b_{ij}.$$

Remark

Shapes of the matrices A and B must be equal.

- **Commutativity:**

$$\forall A, B \in \mathbb{R}^{m \times n} : A + B = B + A$$

- **Commutativity:**

$$\forall A, B \in \mathbb{R}^{m \times n} : A + B = B + A$$

- **Associativity:**

$$\forall A, B \in \mathbb{R}^{m \times n} : (A + B) + C = A + (B + C)$$

Matrix Multiplication

Definition

If A is an $m \times n$ matrix and B is an $n \times r$ matrix, then the product $C = AB$ is an $m \times r$ matrix. The (i, j) entry of the product is computed as follows:

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj} = \sum_{k=1}^n a_{ik}b_{kj}.$$

Matrix Multiplication

Definition

If A is an $m \times n$ matrix and B is an $n \times r$ matrix, then the product $C = AB$ is an $m \times r$ matrix. The (i, j) entry of the product is computed as follows:

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj} = \sum_{k=1}^n a_{ik}b_{kj}.$$

Remark

The number of columns of A must be the same as the number of rows of B .

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$$

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{i1} & a_{i2} & \dots & a_{in} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & \dots & b_{1j} & \dots & b_{1r} \\ b_{21} & \dots & b_{2j} & \dots & b_{2r} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \dots & b_{nj} & \dots & b_{nr} \end{bmatrix}$$

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$$

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{i1} & a_{i2} & \dots & a_{in} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & \dots & b_{1j} & \dots & b_{1r} \\ b_{21} & \dots & b_{2j} & \dots & b_{2r} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \dots & b_{nj} & \dots & b_{nr} \end{bmatrix}$$

Example

Compute AB if

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 3 & -1 & 0 \end{bmatrix}, \quad \text{and} \quad B = \begin{bmatrix} 1 & -3 \\ 4 & 0 \\ 2 & -2 \end{bmatrix}$$

Check that $AB \neq BA$.

Definition

In $\mathbb{R}^{n \times n}$, we define the **identity matrix**

$$I_n = \begin{bmatrix} 1 & 0 & 0 \dots & 0 \\ 0 & 1 & 0 \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 \dots & 1 \end{bmatrix} \in \mathbb{R}^{n \times n}$$

as the $n \times n$ -matrix containing 1 on the diagonal and 0 everywhere else. With this, $A \cdot I_n = A = I_n \cdot A$ for all $A \in \mathbb{R}^{n \times n}$.

- **Associativity:**

$$\forall A \in \mathbb{R}^{m \times n}, B \in \mathbb{R}^{n \times p}, C \in \mathbb{R}^{p \times q} : (AB)C = A(BC)$$

- **Associativity:**

$$\forall A \in \mathbb{R}^{m \times n}, B \in \mathbb{R}^{n \times p}, C \in \mathbb{R}^{p \times q} : (AB)C = A(BC)$$

- **Distributivity:**

$$\begin{aligned} \forall A, B \in \mathbb{R}^{m \times n}, C, D \in \mathbb{R}^{n \times p} : \quad & (A + B)C = AC + BC \\ & A(C + D) = AC + AD \end{aligned}$$

- **Associativity:**

$$\forall A \in \mathbb{R}^{m \times n}, B \in \mathbb{R}^{n \times p}, C \in \mathbb{R}^{p \times q} : (AB)C = A(BC)$$

- **Distributivity:**

$$\begin{aligned} \forall A, B \in \mathbb{R}^{m \times n}, C, D \in \mathbb{R}^{n \times p} : \quad & (A + B)C = AC + BC \\ & A(C + D) = AC + AD \end{aligned}$$

- **Neutral element (identity element):**

$$\forall A \in \mathbb{R}^{m \times n} : \quad I_m A = A I_n = A.$$

Note that $I_m \neq I_n$ for $m \neq n$.

Multiplication by a Scalar

If A is an $m \times n$ matrix and λ is a scalar, then the scalar multiple λA is the $m \times n$ matrix

$$\lambda A = \lambda[a_{ij}] = [\lambda a_{ij}].$$

Multiplication by a Scalar

If A is an $m \times n$ matrix and λ is a scalar, then the scalar multiple λA is the $m \times n$ matrix

$$\lambda A = \lambda[a_{ij}] = [\lambda a_{ij}].$$

- **Distributivity:**

$$(\lambda_1 + \lambda_2)C = \lambda_1 C + \lambda_2 C, \quad C \in \mathbb{R}^{m \times n}, \lambda_1, \lambda_2 \in \mathbb{R},$$

$$\lambda(B + C) = \lambda B + \lambda C, \quad B, C \in \mathbb{R}^{m \times n}, \lambda \in \mathbb{R}.$$

Multiplication by a Scalar

If A is an $m \times n$ matrix and λ is a scalar, then the scalar multiple λA is the $m \times n$ matrix

$$\lambda A = \lambda[a_{ij}] = [\lambda a_{ij}].$$

- **Distributivity:**

$$(\lambda_1 + \lambda_2)C = \lambda_1 C + \lambda_2 C, \quad C \in \mathbb{R}^{m \times n}, \lambda_1, \lambda_2 \in \mathbb{R},$$

$$\lambda(B + C) = \lambda B + \lambda C, \quad B, C \in \mathbb{R}^{m \times n}, \lambda \in \mathbb{R}.$$

- **Associativity:**

$$(\lambda_1 \lambda_2)C = \lambda_1(\lambda_2 C), \quad C \in \mathbb{R}^{m \times n}, \lambda_1, \lambda_2 \in \mathbb{R},$$

$$\lambda(BC) = (\lambda B)C = B(\lambda C) = (BC)\lambda, \quad B \in \mathbb{R}^{m \times n}, C \in \mathbb{R}^{n \times k}.$$

Multiplication by a Scalar

Example

Check that

$$(\lambda_1 + \lambda_2)C = \lambda_1 C + \lambda_2 C$$

for any $\lambda_1, \lambda_2 \in R$ and $C := \begin{bmatrix} 1 & 3 \\ 5 & 7 \end{bmatrix}$.

Definition

For a square matrix $A \in \mathbb{R}^{n \times n}$ a matrix $B \in \mathbb{R}^{n \times n}$ with $AB = I_n = BA$ is called **inverse** and denoted by A^{-1} .

Definition

For a square matrix $A \in \mathbb{R}^{n \times n}$ a matrix $B \in \mathbb{R}^{n \times n}$ with $AB = I_n = BA$ is called **inverse** and denoted by A^{-1} .

Not every matrix A has an inverse A^{-1} . If this inverse does exist, A is called **regular/invertible/non-singular**, otherwise **singular/non-invertible**.

Existence of the Inverse of a 2×2 -Matrix

Consider a matrix

$$A := \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in \mathbb{R}^{2 \times 2}.$$

Existence of the Inverse of a 2×2 -Matrix

Consider a matrix

$$A := \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in \mathbb{R}^{2 \times 2}.$$

Then for $B = \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix}$ we get

$$AB = \begin{bmatrix} a_{11}a_{22} - a_{12}a_{21} & 0 \\ 0 & a_{11}a_{22} - a_{12}a_{21} \end{bmatrix} = (a_{11}a_{22} - a_{12}a_{21})I_2.$$

Existence of the Inverse of a 2×2 -Matrix

Consider a matrix

$$A := \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in \mathbb{R}^{2 \times 2}.$$

Then for $B = \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix}$ we get

$$AB = \begin{bmatrix} a_{11}a_{22} - a_{12}a_{21} & 0 \\ 0 & a_{11}a_{22} - a_{12}a_{21} \end{bmatrix} = (a_{11}a_{22} - a_{12}a_{21})I_2.$$

Therefore

$$A^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix},$$

if $a_{11}a_{22} - a_{12}a_{21} \neq 0$.

Example

The matrices

$$A = \begin{bmatrix} 4 & 5 & 4 \\ 7 & 7 & 6 \\ 2 & 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -1 & 2 \\ 5 & -4 & 4 \\ -7 & 6 & -7 \end{bmatrix}$$

are inverse to each other since $AB = I = BA$.

Definition

For $A \in \mathbb{R}^{m \times n}$ the matrix $B \in \mathbb{R}^{n \times m}$ with $b_{ij} = a_{ji}$ is called the transpose of A . We write $B = A^T$.

Definition

For $A \in \mathbb{R}^{m \times n}$ the matrix $B \in \mathbb{R}^{n \times m}$ with $b_{ij} = a_{ji}$ is called the transpose of A . We write $B = A^T$.

A^T is achieved by any one of the following equivalent actions:

Definition

For $A \in \mathbb{R}^{m \times n}$ the matrix $B \in \mathbb{R}^{n \times m}$ with $b_{ij} = a_{ji}$ is called the transpose of A . We write $B = A^T$.

A^T is achieved by any one of the following equivalent actions:

- reflect A over its main diagonal (which runs from top-left to bottom-right) to obtain A^T ,

Definition

For $A \in \mathbb{R}^{m \times n}$ the matrix $B \in \mathbb{R}^{n \times m}$ with $b_{ij} = a_{ji}$ is called the transpose of A . We write $B = A^T$.

A^T is achieved by any one of the following equivalent actions:

- reflect A over its main diagonal (which runs from top-left to bottom-right) to obtain A^T ,
- write the rows of A as the columns of A^T ,

Definition

For $A \in \mathbb{R}^{m \times n}$ the matrix $B \in \mathbb{R}^{n \times m}$ with $b_{ij} = a_{ji}$ is called the transpose of A . We write $B = A^T$.

A^T is achieved by any one of the following equivalent actions:

- reflect A over its main diagonal (which runs from top-left to bottom-right) to obtain A^T ,
- write the rows of A as the columns of A^T ,
- write the columns of A as the rows of A^T .

Properties of inverses and transposes

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$
- $(AB)^{-1} = B^{-1}A^{-1}$

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$
- $(AB)^{-1} = B^{-1}A^{-1}$
- $(A^T)^T = A$

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$
- $(AB)^{-1} = B^{-1}A^{-1}$
- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$
- $(AB)^{-1} = B^{-1}A^{-1}$
- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(AB)^T = B^T A^T$

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$
- $(AB)^{-1} = B^{-1}A^{-1}$
- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(AB)^T = B^T A^T$
- If A is invertible, then so is A^T and $(A^{-1})^T = (A^T)^{-1}$

Properties of inverses and transposes

- $AA^{-1} = I = A^{-1}A$
- $(AB)^{-1} = B^{-1}A^{-1}$
- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(AB)^T = B^T A^T$
- If A is invertible, then so is A^T and $(A^{-1})^T = (A^T)^{-1}$

Definition

A square matrix A is symmetric if $A = A^T$.

Properties of inverses and transposes

Example

Prove that for any square matrix A the matrices $A + A^T$ and $A \cdot A^T$ are symmetric.

Properties of inverses and transposes

Example

Prove that for any square matrix A the matrices $A + A^T$ and $A \cdot A^T$ are symmetric.

Example

Prove that for any symmetric matrices A, B the matrix $A + B$ is symmetric.

Properties of inverses and transposes

Example

Prove that for any square matrix A the matrices $A + A^T$ and $A \cdot A^T$ are symmetric.

Example

Prove that for any symmetric matrices A, B the matrix $A + B$ is symmetric.

Remark

The product of two symmetric matrices is not symmetric in general, e.g. the matrices $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, but the matrix $AB = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ is not symmetric.

Compact Representations of Systems of Linear Equations

The system of linear equations

$$\begin{cases} 2x_1 + 3x_2 - 4x_3 &= 5 \\ x_1 - x_2 + 3x_3 &= -2 \\ -x_1 + 2x_2 + 5x_3 &= 7 \end{cases}$$

Compact Representations of Systems of Linear Equations

The system of linear equations

$$\begin{cases} 2x_1 + 3x_2 - 4x_3 &= 5 \\ x_1 - x_2 + 3x_3 &= -2 \\ -x_1 + 2x_2 + 5x_3 &= 7 \end{cases}$$

can be written as vector equation

$$\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} x_1 + \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix} x_2 + \begin{bmatrix} -4 \\ 3 \\ 5 \end{bmatrix} x_3 = \begin{bmatrix} 5 \\ -2 \\ 7 \end{bmatrix},$$

Compact Representations of Systems of Linear Equations

The system of linear equations

$$\begin{cases} 2x_1 + 3x_2 - 4x_3 &= 5 \\ x_1 - x_2 + 3x_3 &= -2 \\ -x_1 + 2x_2 + 5x_3 &= 7 \end{cases}$$

can be written as vector equation

$$\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} x_1 + \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix} x_2 + \begin{bmatrix} -4 \\ 3 \\ 5 \end{bmatrix} x_3 = \begin{bmatrix} 5 \\ -2 \\ 7 \end{bmatrix},$$

and can be compactly written in matrix notation as follows

$$\begin{bmatrix} 2 & 3 & -4 \\ 1 & -1 & 3 \\ -1 & 2 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 5 \\ -2 \\ 7 \end{bmatrix}$$

Compact Representations of Systems of Linear Equations

Note that Ax is a linear combination of the columns of A :

$$\begin{bmatrix} 2 & 3 & -4 \\ 1 & -1 & 3 \\ -1 & 2 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} x_1 + \begin{bmatrix} 3 \\ -1 \\ 2 \end{bmatrix} x_2 + \begin{bmatrix} -4 \\ 3 \\ 5 \end{bmatrix} x_3$$

Systems of Linear Equations

Let us introduce the general form of a system of linear equations,

$$\begin{aligned}a_{11}x_1 + \dots + a_{1n}x_n &= b_1 \\ \vdots \\ a_{m1}x_1 + \dots + a_{mn}x_n &= b_m\end{aligned}$$

where $a_{ij} \in \mathbb{R}$ and $b_i \in \mathbb{R}$ are known constants and x_j are unknowns, $i = 1, \dots, m$, $j = 1, \dots, n$.

Systems of Linear Equations

Let us introduce the general form of a system of linear equations,

$$\begin{aligned}a_{11}x_1 + \dots + a_{1n}x_n &= b_1 \\&\vdots \\a_{m1}x_1 + \dots + a_{mn}x_n &= b_m\end{aligned}$$

where $a_{ij} \in \mathbb{R}$ and $b_i \in \mathbb{R}$ are known constants and x_j are unknowns, $i = 1, \dots, m$, $j = 1, \dots, n$. It can be written in matrix form $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \\ a_{m1} & \dots & a_{mn} \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$$

Particular and General Solution

Consider the following system of linear equations (SLE):

$$\begin{bmatrix} 1 & 0 & 7 & 2 \\ 0 & 1 & 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \end{bmatrix}$$

Particular and General Solution

Consider the following system of linear equations (SLE):

$$\begin{bmatrix} 1 & 0 & 7 & 2 \\ 0 & 1 & 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \end{bmatrix}$$

or, equivalently $\sum_{i=1}^4 x_i \mathbf{c}_i = \mathbf{b}$, where $\mathbf{c}_i$ is the i th column of the matrix and $\mathbf{b}$ is the right-hand-side.

Particular and General Solution

Consider the following system of linear equations (SLE):

$$\begin{bmatrix} 1 & 0 & 7 & 2 \\ 0 & 1 & 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \end{bmatrix}$$

or, equivalently $\sum_{i=1}^4 x_i \mathbf{c}_i = \mathbf{b}$, where $\mathbf{c}_i$ is the i th column of the matrix and $\mathbf{b}$ is the right-hand-side. $[5, 8, 0, 0]^T$ is a solution, since $\mathbf{b} = 5\mathbf{c}_1 + 8\mathbf{c}_2 + 0\mathbf{c}_3 + 0\mathbf{c}_4$.

Particular and General Solution

Consider the following system of linear equations (SLE):

$$\begin{bmatrix} 1 & 0 & 7 & 2 \\ 0 & 1 & 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \end{bmatrix}$$

or, equivalently $\sum_{i=1}^4 x_i \mathbf{c}_i = \mathbf{b}$, where $\mathbf{c}_i$ is the i th column of the matrix and $\mathbf{b}$ is the right-hand-side. $[5, 8, 0, 0]^T$ is a solution, since $\mathbf{b} = 5\mathbf{c}_1 + 8\mathbf{c}_2 + 0\mathbf{c}_3 + 0\mathbf{c}_4$. This solution is called a **particular solution** or **special solution**.

Particular and General Solution

All solutions of the SLE, which is called the **general solution**, is the set

$$\left\{ \mathbf{x} \in \mathbb{R}^4; \mathbf{x} = \begin{bmatrix} 5 \\ 8 \\ 0 \\ 0 \end{bmatrix} + \lambda_1 \begin{bmatrix} -7 \\ -3 \\ 1 \\ 0 \end{bmatrix} + \lambda_2 \begin{bmatrix} -2 \\ 4 \\ 0 \\ 1 \end{bmatrix}, \lambda_1, \lambda_2 \in \mathbb{R} \right\}.$$

The algorithm.

Particular and General Solution

The algorithm.

1. Find a particular solution to $A\mathbf{x} = \mathbf{b}$.

Particular and General Solution

The algorithm.

1. Find a particular solution to $A\mathbf{x} = \mathbf{b}$.
2. Find all solutions to $A\mathbf{x} = \mathbf{0}$.

The algorithm.

1. Find a particular solution to $A\mathbf{x} = \mathbf{b}$.
2. Find all solutions to $A\mathbf{x} = \mathbf{0}$.
3. Combine the solutions from 1 and 2 to the general solution.