

# Image Compression by Partial Differential Equations

(Séminaire Doctorants IRIMAS)

Thomas Jacumin

sous la direction de Zakaria Belhachmi

Goal : reconstruct missing parts of an image  $f : D \rightarrow [0, 1]$  from  $K \subset D$ .




Figure – Inpainting




Figure – Examples [2]

# JPEG

Each block  $8 \times 8$  of an image is described as a function

$$F : \{0, \dots, 7\}^2 \rightarrow \mathbb{R}.$$

# JPEG

Each block  $8 \times 8$  of an image is described as a function

$$F : \{0, \dots, 7\}^2 \rightarrow \mathbb{R}.$$

For  $0 \leq u, v \leq 7$ ,

$$F_{u,v} : (x, y) \mapsto C \cos \frac{(2x+1)u\pi}{16} \cos \frac{(2y+1)v\pi}{16}.$$

## JPEG

Each block  $8 \times 8$  of an image is described as a function

$$F : \{0, \dots, 7\}^2 \rightarrow \mathbb{R}.$$

For  $0 \leq u, v \leq 7$ ,

$$F_{u,v} : (x, y) \mapsto C \cos \frac{(2x+1)u\pi}{16} \cos \frac{(2y+1)v\pi}{16}.$$

Then, we can decompose  $F$  as :

$$F = \sum_{0 \leq u, v \leq 7} c_{u,v} F_{u,v}.$$

## JPEG

Each block  $8 \times 8$  of an image is described as a function

$$F : \{0, \dots, 7\}^2 \rightarrow \mathbb{R}.$$

For  $0 \leq u, v \leq 7$ ,

$$F_{u,v} : (x, y) \mapsto C \cos \frac{(2x+1)u\pi}{16} \cos \frac{(2y+1)v\pi}{16}.$$

Then, we can decompose  $F$  as :

$$F = \sum_{0 \leq u, v \leq 7} c_{u,v} F_{u,v}.$$

For compression purpose, we can neglect small  $c_{u,v}$ .

We do not do this : we remove parts of the image and reconstruct them by inpainting :




Figure – Compression by Inpainting.

## Question 1

How to choose pixels to keep ?

## Question 1

How to choose pixels to keep ?

## Answer 1

We should keep relevant pixels to obtain a good reconstruction.

## Question 1

How to choose pixels to keep ?

## Answer 1

We should keep relevant pixels to obtain a good reconstruction.

## Questions 2-3

What is a good reconstruction ? How to find these pixels ?

Let  $D \subset \mathbb{R}^2$  the support of an image  $f : D \rightarrow [0, 1]$ . Some errors examples

### Example 1

$$\|u - f\|_{L^2(D)} := \left( \int_D (u - f)^2 \, dx \right)^{1/2}.$$

### Example 2

$$|u - f|_{H^1(D)} := \left( \int_D |\nabla u - \nabla f|^2 \, dx \right)^{1/2}.$$

## Shape Optimisation Problem

$$\min_{K \subseteq D} \{ \mathcal{E}(u_K) \mid u_K \text{ solution of an inpainting problem} \},$$

## Shape Optimisation Problem

$$\min_{K \subseteq D} \{ \mathcal{E}(u_K) \mid u_K \text{ solution of an inpainting problem} \},$$

A trivial solution is  $K = D$ . We need to add constraint on the “size” of  $K$ .

## Shape Optimisation Problem

$$\min_{K \subseteq D} \{ \mathcal{E}(u_K) \mid u_K \text{ solution of an inpainting problem} \},$$

A trivial solution is  $K = D$ . We need to add constraint on the "size" of  $K$ .

## Compression problem

$$\min_{K \subseteq D, m(K) \leq c} \{ \mathcal{E}(u_K) \mid u_K \text{ solution of an inpainting problem} \},$$

## Shape Optimisation Problem

$$\min_{K \subseteq D} \{ \mathcal{E}(u_K) \mid u_K \text{ solution of an inpainting problem} \},$$

A trivial solution is  $K = D$ . We need to add constraint on the "size" of  $K$ .

## Compression problem

$$\min_{K \subseteq D, m(K) \leq c} \{ \mathcal{E}(u_K) \mid u_K \text{ solution of an inpainting problem} \},$$

The existence of a solution depends on the inpainting problem, on the error  $\mathcal{E}$  and on  $m$ .

## Problem (Homogeneous Diffusion Inpainting [1])

Find  $u$  in  $H^1(D)$  such that


$$\begin{cases} -\Delta u = 0, & \text{in } D \setminus K, \\ u = f, & \text{in } K, \\ \frac{\partial u}{\partial n} = 0, & \text{on } \partial D. \end{cases} \quad (1)$$

## Problem (Heat equation)

Find  $u$  such that

$$\begin{cases} \partial_t u - \Delta u = 0, & \text{in } [0, +\infty[ \times D, \\ u = 100, & \text{on } [0, +\infty[ \times \Gamma, \\ \partial_n u = 0, & \text{on } [0, +\infty[ \times \partial D \setminus \Gamma. \end{cases} \quad (2)$$

and  $u(0, \cdot) = 0$  in  $D$ .



## Problem (Heat equation)

Find  $u$  such that

$$\begin{cases} \partial_t u - \Delta u = 0, & \text{in } [0, +\infty[ \times D, \\ u = 100, & \text{on } [0, +\infty[ \times \Gamma, \\ \partial_n u = 0, & \text{on } [0, +\infty[ \times \partial D \setminus \Gamma. \end{cases} \quad (3)$$


and  $u(0, \cdot) = 0$  in  $D$ .

## Problem (Heat equation ( $t \rightarrow +\infty$ ))

Find  $u$  such that

$$\begin{cases} \partial_t u - \Delta u = 0, & \text{in } [0, +\infty[ \times D, \\ u = 100, & \text{on } [0, +\infty[ \times \Gamma, \\ \partial_n u = 0, & \text{on } [0, +\infty[ \times \partial D \setminus \Gamma. \end{cases} \quad (4)$$

and  $u(0, \cdot) = 0$  in  $D$ .



## Theorem

*If the inpainting is the homogeneous diffusion inpaiting*

$$\mathcal{E}(u) = |u - f|_{H^1(D)} := \left( \int_D |\nabla u - \nabla f|^2 \, dx \right)^{1/2},$$

*and if*

$$m(K) = \text{cap}(K) := \inf \left\{ \int_D |\nabla u|^2 \, dx + \int_D u^2 \, dx \mid u \in H_0^1(D), \, u \geq 1 \text{ p.p. dans } K \right\},$$

*Then, the compression problem admits at least one solution.*

## Question

In practice, how to construct  $K$ ?

# Topological Gradient

**Goal :** Determine the influence of make a hole in  $K$  on the error.

Let  $K_\varepsilon := K \setminus B(x_0, \varepsilon)$  such that  $B(x_0, \varepsilon) \subset K$ .

# Topological Gradient

**Goal :** Determine the influence of make a hole in  $K$  on the error.

Let  $K_\varepsilon := K \setminus B(x_0, \varepsilon)$  such that  $B(x_0, \varepsilon) \subset K$ .

# Topological Gradient

**Goal :** Determine the influence of make a hole in  $K$  on the error.

Let  $K_\varepsilon := K \setminus B(x_0, \varepsilon)$  such that  $B(x_0, \varepsilon) \subset K$ .

$$j : A \subset D \mapsto \min_{v \in H^1(D), v=f \text{ in } A} \int_D |\nabla v|^2 \, dx.$$

## Proposition

*The compression problem is equivalent to  $\max_{K \subset D, m(K) \leq c} j(K)$ .*

# Topological Gradient

$$\begin{aligned} \max_{K \subset D, m(K) \leq c} j(K) &= \max_{K \subset D, m(K) \leq c} \min_{v \in H^1(D), v=f \text{ in } K} \int_D |\nabla v|^2 dx \\ &= \max_{K \subset D, m(K) \leq c} \left( \underbrace{\left( \min_{v \in H^1(D), v=f \text{ in } K} \int_{D \setminus K} |\nabla v|^2 dx \right)}_{\geq 0} + \int_K |\nabla f|^2 dx \right). \end{aligned}$$

Then,

$$\max_{K \subset D, m(K) \leq c} \int_K |\nabla f|^2 dx \leq \max_{K \subset D, m(K) \leq c} j(K).$$

# Topological Gradient

**Goal :** Determine the influence of make a hole in  $K$  on the error.

Let  $K_\varepsilon := K \setminus B(x_0, \varepsilon)$  such that  $B(x_0, \varepsilon) \subset K$ .

$$j : A \subset D \mapsto \min_{v \in H^1(D), v=f \text{ in } A} \int_D |\nabla v|^2 \, dx.$$

## Proposition

The compression problem is equivalent to  $\max_{K \subset D, m(K) \leq c} j(K)$ .

## Proposition

When  $\varepsilon$  tends to 0,

$$j(K_\varepsilon) - j(K) = -|\Delta f(x_0)|^2 \frac{\pi}{2} \varepsilon^4 + o(\varepsilon^4).$$

# Topological Gradient




Figure –  $f$  et  $|\Delta f|$ .

# Topological Gradient

## Question

How to build  $K$  ?

# Topological Gradient

## Question

How to build  $K$  ?

## Answer

Keep points  $x_0$  where the quantity  $|\Delta f(x_0)|$  is maximal.

# Topological Gradient

## Question

How to build  $K$ ?

## Answer

Keep points  $x_0$  where the quantity  $|\Delta f(x_0)|$  is maximal.

# Fat Pixels

For  $m > 0$  and  $n \in \mathbb{N}^*$ , we set

$$\mathcal{A}_{m,n} := \left\{ \bigcup_{i=1}^n \overline{B(x_i, r)} \mid x_i \in D, r = mn^{-1/2} \right\}.$$

# Fat Pixels

## Question

How to build  $K$ ?

# Fat Pixels

## Question

How to build  $K$  ?

## Answer

Built  $K$  such that the pixels density increase with the quantity  $|\Delta f|$ .



Figure – Topological Gradient vs Fat Pixels.






Figure – B-Tree vs Random.





Figure – Image with Gaussian Noise.






Figure – Image with Salt and Pepper Noise.




Figure – Image with Gaussian Noise.




Figure – Image with Salt and Pepper Noise.

# Thanks for your attention !

-  Z. Belhachmi, D. Bucur, B. Burgeth, and J. Weickert, *How to choose interpolation data in images*, 70, pp. 333–352.
-  C. Schmaltz, P. Peter, M. Mainberger, F. Huth, J. Weickert, and A. Bruhn, *Understanding, optimising, and extending data compression with anisotropic diffusion*, 108.