

Spectraltheory

Prof. Dr. Tobias Lamm

Sommersemester 2017

Karlsruher Institut für Technologie

Contents

I	Unbounded, adjoint and self-adjoint operators	2
II	Representation Theorems	9

Chapter I

Unbounded, adjoint and self-adjoint operators

Let H be a separable Hilbert space, i.e. a real or complex inner product space that is also a complete metric space with respect to the distance function induced by the inner product $\langle \cdot, \cdot \rangle$ on H .

Recall: A mapping $\langle \cdot, \cdot \rangle: H \times H \rightarrow \mathbb{C}$ is called an **inner product**, if for all $x, y \in H$, $\lambda \in \mathbb{C}$ holds:

$$(S1) \quad \langle x_1 + x_2, y \rangle = \langle x_1, y \rangle + \langle x_2, y \rangle, \quad \langle x, y_1 + y_2 \rangle = \langle x, y_1 \rangle + \langle x, y_2 \rangle$$

$$(S2) \quad \langle \lambda x, y \rangle = \lambda \langle x, y \rangle, \quad \langle x, \lambda y \rangle = \bar{\lambda} \langle x, y \rangle$$

$$(S3) \quad \langle x, y \rangle = \overline{\langle y, x \rangle}$$

$$(S4) \quad \langle x, x \rangle \geq 0, \quad \langle x, x \rangle = 0 \iff x = 0$$

A linear operator T in H is a linear map

$$u \mapsto Tu$$

defined on a subspace $\mathcal{D}(T)$ of H , and we call $\mathcal{D}(T)$ the **domain** of T . For $T: \mathcal{D}(T) \rightarrow H$ we denote the **range** of T with

$$\mathcal{R}(T) := \text{Image}(T).$$

We say that T is **bounded** if it is continuous from $\mathcal{D}(T)$ into H , with respect to the topology induced by H . We recall for the definition of bounded operators that if $\mathcal{D}(T) = H$ holds, boundedness is equivalent to continuity in 0, boundedness of $T(U_{(X, \|\cdot\|)})$ in Y and there $\exists c < \infty$ such that $\|Tx\| \leq c\|x\|$, for proof see the [functional analysis](#) course.

From now on, if $\mathcal{D}(T) \neq H$ we will assume that $\mathcal{D}(T)$ is **dense** in H , i.e. $\overline{\mathcal{D}(T)} = H$. If in this case T would be bounded then T has a unique continuous extension to all of H . As this simplifies many considerations some of the following theorems would be trivial, and hence, we won't focus on bounded operators during this lecture.

Recall: An operator is called **closed** if the graph

$$G(T) := \{(x, y) \in H \times H \mid x \in \mathcal{D}(T), y = Tx\}$$

is closed in $H \times H$.

Definition I.1: Let $T: \mathcal{D}(T) \rightarrow H$ be a (linear) operator with $\mathcal{D}(T)$ dense in H . Then T is called **closed** if for all

$$u_n \in \mathcal{D}(T), \quad u_n \rightarrow u \in H \quad \text{and} \quad Tu_n \rightarrow v \in H$$

follows that

$$u \in \mathcal{D}(T), \quad v = Tu$$

holds.

Example:

a) Let $H = L^2(\mathbb{R}^n)$, then $\mathcal{D}(T_0) = C_c^\infty(\mathbb{R}^n)$ is dense in H . Define the operator

$$T_0 = -\Delta,$$

and take $u \in W^{2,2}(\mathbb{R}^n) \setminus C_c^\infty(\mathbb{R}^n)$, s.t $u \in L^2(\mathbb{R}^n)$. Due to the density:

$$\exists (u_n)_{n \in \mathbb{N}} \subseteq C_c^\infty(\mathbb{R}^n): \quad u_n \rightarrow u \text{ in } W^{2,2}(\mathbb{R}^n).$$

As a result, $(u_n, -\Delta u_n) \in G(T_0)$ converges in $L^2 \times L^2$ to $(u, -\Delta u) \notin G(T_0)$.

b) Let $H = L^2(\mathbb{R}^n)$, and set $\mathcal{D}(T_1) = W^{2,2}(\mathbb{R}^n) \subseteq H$. Define the operator

$$T_1 = -\Delta.$$

For $(u_n)_{n \in \mathbb{N}} \subseteq \mathcal{D}(T_1)$ with

$$u_n \rightarrow u \in H \text{ and } (-\Delta u_n) \rightarrow v \in L^2$$

follows that $-\Delta u = v \in L^2(\mathbb{R}^n)$ weakly, i.e. for all $\varphi \in C_c^\infty(\mathbb{R}^n)$:

$$\int_{\mathbb{R}^n} v \varphi \leftarrow \int_{\mathbb{R}^n} (-\Delta u_n) \varphi = \int_{\mathbb{R}^n} u_n (-\Delta \varphi) \longrightarrow \int_{\mathbb{R}^n} u (-\Delta \varphi).$$

$\xrightarrow{PDE} u \in W^{2,2}(\mathbb{R}^n) = \mathcal{D}(T_1) \Rightarrow T_1$ is closed.

Definition I.2: An operator T is called **closable** $\iff \overline{G(T)}$ is a graph.

Remark: We call \overline{T} the **closure** of T , and in such case we have

$$\mathcal{D}(\overline{T}) := \{x \in H \mid \exists y: (x, y) \in \overline{G(T)}\}$$

For any $x \in \mathcal{D}(\overline{T})$ the assumption that $\overline{G(T)}$ is a graph implies that y is unique and hence

$$\Rightarrow G(\overline{T}) = \overline{G(T)}, \quad \overline{T}x := y$$

Equivalently, $\mathcal{D}(\overline{T})$ is the set of all $x \in H$ such that there exists a sequence $x_n \in \mathcal{D}(T)$ with $x_n \rightarrow x$ in H and Tx_n is a Cauchy sequence. For such x we define

$$\overline{T}x := \lim_{n \rightarrow \infty} Tx_n$$

Example: Let $T_0 := -\Delta$, $\mathcal{D}(T_0) = C_c^\infty(\mathbb{R}^n)$ is closable with $\overline{T_0} = T_1$.

Proof: Let $u \in L^2(\mathbb{R}^n)$ such that there exists $(u_n)_{n \in \mathbb{N}} \subseteq C_c^\infty(\mathbb{R}^n)$ with $u_n \rightarrow u$ in L^2 and $-\Delta u_n \rightarrow v$ in L^2 , as above:

$$-\Delta u = v \in L^2$$

For a given u the function v is unique, and hence, T_0 is closable. Let $\overline{T_0}$ be the closure with domain $\mathcal{D}(\overline{T_0})$ and $u \in \mathcal{D}(T_0)$

$$\Rightarrow \Delta u \in L^2 \Rightarrow u \in W^{2,2}(\mathbb{R}^n) = \mathcal{D}(T_1) \Rightarrow \mathcal{D}(\overline{T_0}) \subseteq \mathcal{D}(T_1)$$

but $C_c^\infty(\mathbb{R}^n)$ is dense in $\mathcal{D}(T_1) = W_{2,2}(\mathbb{R}^n)$

$$\Rightarrow W^{2,2}(\mathbb{R}^n) \subseteq \mathcal{D}(\overline{T_0}) \Rightarrow \mathcal{D}(\overline{T_0}) = \mathcal{D}(T_1)$$

$$\Rightarrow T_1 = \overline{T_0}.$$

□

Remark: Assume for a second in the example above that $W^{2,2} \not\subseteq \mathcal{D}(T_0)$ holds:

$$\Rightarrow \exists u \in W^{2,2} \setminus \mathcal{D}(\overline{T_0}), \exists (u_n)_{n \in \mathbb{N}} \in C_c^\infty(\mathbb{R}^n) : u_n \rightarrow u \text{ in } W^{2,2},$$

using the same arguments as in the example above $\Rightarrow \overline{T_0}$ not closed!

Recall: If $T: H \rightarrow H$ is bounded then T^* is defined through

$$\langle u, T^*v \rangle = \langle Tu, v \rangle, \forall u, v \in H$$

$u \mapsto \langle Tu, v \rangle$ defines a continuous linear map on H ($\in H'$). Riesz' representation theorem then ensures the existence of T^* .

Definition I.3: If T is an unbounded operator on H with dense domain we define

$$\mathcal{D}(T^*) := \left\{ v \in H : \mathcal{D}(T) \ni u \mapsto \langle Tu, v \rangle \text{ can be extended as a linear continuous form on } H \right\}$$

Using Riesz' representation theorem $\exists! f \in H$:

$$\langle u, f \rangle = \langle Tu, v \rangle, \forall u \in \mathcal{D}(T)$$

then define $T^*v = f$, where the uniqueness follows from the density of $\mathcal{D}(T)$ in H .

Remark: If $\mathcal{D}(T) = H$ and T is bounded then we recover the “old” adjoint.

Example: $T_0^* = T_1$,

$$\begin{aligned} \mathcal{D}(T_0^*) &= \left\{ v \in L^2(\mathbb{R}^n) : C_c^\infty(\mathbb{R}^n) \ni u \mapsto \langle -\Delta u, v \rangle \text{ extendable as a lin. continuous form on } L^2(\mathbb{R}^n) \right\} \\ &= \left\{ v \in L^2(\mathbb{R}^n) : -\Delta v \in L^2(\mathbb{R}^n) \right\} = W^{2,2}(\mathbb{R}^n) = \mathcal{D}(T_1) \end{aligned}$$

Damit ist

$$\langle T_1 u, v \rangle = \langle -\Delta u, v \rangle = \int v(-\Delta u) = - \int (-\Delta v) u = \langle u, T_1 v \rangle$$

Theorem I.1: T^* is a closed operator.

Proof: $v_n \in \mathcal{D}(T^*)$ such that $v_n \rightarrow v$ in H and $T^*v_n \rightarrow w^*$ in H for $(v, w^*) \in H \times H$. For all $u \in \mathcal{D}(T)$ we have

$$\langle Tu, v \rangle = \lim_{n \rightarrow \infty} \langle Tu, v_n \rangle = \lim_{n \rightarrow \infty} \langle u, T^*v_n \rangle = \langle u, w^* \rangle$$

$(H \ni u \mapsto \langle u, w^* \rangle \text{ is continuous}) \Rightarrow v \in \mathcal{D}(T^*)$ and $w^* = T^*v$ by definition. \square

Theorem I.2: Let T be an operator in H with domain $\mathcal{D}(T)$. Then

$$G(T^*) = \left(V \left(\overline{G(T)} \right) \right)^\perp$$

where $V: H \times H \rightarrow H \times H, V(x, y) = (y, -x)$ ($V^2 = -\mathbb{I}$).

Proof: Let $u \in \mathcal{D}(T), (v, w^*) \in H \times H$

$$\Rightarrow \langle V(u, Tu), (v, w^*) \rangle_{H \times H} = \langle Tu, v \rangle - \langle u, w^* \rangle$$

Considering the right-hand side it follows

$$\langle Tu, v \rangle - \langle u, w^* \rangle = 0 \quad \forall u \in \mathcal{D}(T) \iff v \in \mathcal{D}(T^*) \text{ and } w^* = T^*v \iff (v, w^*) \in G(T^*),$$

and considering the left-hand side:

$$\Rightarrow \langle V(u, Tu), (v, w^*) \rangle_{H \times H} = 0 \quad \forall u \in \mathcal{D}(T) \iff (v, w^*) \in V(G(T))^\perp$$

In general: $U^\perp = \overline{U}^\perp$, and hence

$$\Rightarrow V(G(T))^\perp = \left(\overline{V(G(T))} \right)^\perp = \left(V \left(\overline{G(T)} \right) \right)^\perp.$$

\square

Theorem I.3: Let T be a closable operator. Then:

a) $\mathcal{D}(T^*)$ is dense in H

b) $T^{**} := (T^*)^* = \overline{T}$

Proof:

a) Proof through contradiction: $D(T^*)$ not dense in $H \rightarrow \exists w \neq 0 : \langle w, v \rangle = 0 \ \forall v \in \overline{\mathcal{D}(T^*)}$

$$\implies \langle (0, w), (T^*v, -v) \rangle_{H \times H} = 0 \ \forall v \in \mathcal{D}(T^*)$$

$$\implies (0, w) \perp V(G(T^*))$$

$$\xrightarrow[\text{I.2}]{\text{Thm}} V(\overline{G(T)}) = G(T^*)^\perp$$

$$\implies V(G(T^*)^\perp) = \overline{G(T)}$$

For any $M \subseteq H \times H$ we have $V(M^\perp) = V(M)^\perp$ since for $(u, v) \in V(M)^\perp$, $(x, y) \in M$

$$\langle V(u, v), (x, y) \rangle_{H \times H} = -\langle (u, v), V(x, y) \rangle_{H \times H} \Rightarrow V(u, v) \in M^\perp \Rightarrow (u, v) \in V(M^\perp)$$

$$\implies V(G(T^*)^\perp) = \overline{G(T)} = G(\overline{T}) \implies (0, w) \in G(\overline{T}) \implies w = 0$$

b) $G(T^{**}) \xrightarrow[\text{I.2}]{\text{Thm}} V(\overline{G(T^*)})^\perp \xrightarrow[\text{I.1}]{\text{Thm}} V(G(T^*)^\perp)^\perp \stackrel{(\perp)}{=} G(\overline{T}) \implies \mathcal{D}(T^{**}) = \mathcal{D}(\overline{T}), T^{**} = \overline{T}$

□

Definition I.4: We say $T: \mathcal{D}(T) \rightarrow H$ is **symmetric** if and only if

$$\langle Tu, v \rangle = \langle u, Tv \rangle \quad \forall u, v \in \mathcal{D}(T)$$

Example: $T_0 = -\Delta$, $\mathcal{D}(T_0) = C_c^\infty(\mathbb{R}^n)$

$$\int_{\mathbb{R}^n} (-\Delta u) v = \int_{\mathbb{R}^n} u (-\Delta v)$$

Remark: If T is symmetric $\Rightarrow \mathcal{D}(T) \subseteq \mathcal{D}(T^*)$ and

$$Tu = T^*u \quad \forall u \in \mathcal{D}(T)$$

$\Rightarrow (T^*, \mathcal{D}(T^*))$ is an extension of $(T, \mathcal{D}(T))$.

Lemma I.1: A symmetric operator T is closable.

Proof: It suffice to show that for $u_n \in \mathcal{D}(T)$ with $u_n \rightarrow 0$ and $u_n \rightarrow x \in H$ we have $x = 0$

$$\langle x, v \rangle \leftarrow \langle Tu_n, v \rangle = \langle u_n, Tu \rangle \rightarrow \langle 0, Tu \rangle = 0 \quad \forall v \in \mathcal{D}(T)$$

$\Rightarrow x = 0$. □

Remark: The proof actually shows that if $\mathcal{D}(T^*)$ is dense in H , then T is closable.

Definition I.5: We call an operator T self-adjoint if

$$T = T^* \text{ and } Tu = T^*u \quad \forall u \in \mathcal{D}(T),$$

note that the first property implies that $\mathcal{D}(T) = \mathcal{D}(T^*)$.

Theorem I.4: Every self-adjoint operator is closable.

Proof: [Lemma I.1](#) □

Theorem I.5: Let T be an invertible self-adjoint operator, then T^{-1} is also self-adjoint.

Proof: For $T: \mathcal{D}(T) \rightarrow \mathcal{R}(T)$ consider

Step 1 $\mathcal{R}(T)$ is dense in H . We have to show that $\mathcal{R}(T)^\perp = \{0\}$.

Let $w \in H$ such that

$$\langle Tu, w \rangle = 0 \quad \forall u \in \mathcal{D}(T)$$

$$\implies w \in \mathcal{D}(T^*) \text{ and } T^*w = 0 \xrightarrow[\text{s.a.}]{\text{inj.}} w = 0.$$

Step 2 Let $w: H \times H \rightarrow H \times H$, $w(x, y) = (y, x)$

$$\implies G(T^{-1}) = \left\{ (x, T^{-1}x) : x \in \mathcal{D}(T) \right\} = w(G(T)) = \left\{ (Ty, y) : y \in \mathcal{D}(T) \right\}$$

$$\begin{aligned} G(T^{-1}) &= G((T^*)^{-1}) \xrightarrow[\text{Thm. I.2}]{\text{Proof}} w(V(G(T)^\perp)) \\ &= V(w(G(T))^\perp) = V(w(G(T)))^\perp \\ &= V(G(T^{-1}))^\perp \xrightarrow[\text{I.2}]{\text{Thm.}} G((T^{-1})^*) \end{aligned}$$

$$\Rightarrow T^{-1} = (T^{-1})^*$$

□

Chapter II

Representation Theorems

Theorem II.1 (Riesz): *Let $u \mapsto F(u)$ be a linear continuous function on H . Then $\exists! w \in H$:*

$$F(u) = \langle u, w \rangle \quad \forall u \in H$$

Lax-Milgram: V Hilbertspace, sesquilinear form is defined on $V \times V$, $(u, v) \mapsto \alpha(u, v)$ continuous with

$$|\alpha(u, v)| \leq c \|u\| \|v\| \quad \forall u, v \in V$$

Riesz: \exists linear map $A: V \rightarrow V$:

$$\alpha(u, v) = \langle Au, v \rangle$$

Definition II.1: A bilinear form $a: V \times V \rightarrow \mathbb{R}$ is ***V-coercive*** if there exists $\lambda > 0$ such that

$$a(u, u) \geq \lambda \|u\|^2 \quad \forall u \in V$$

Theorem II.2: *Let a be a continuous sesquilinear and V -coercive on $V \times V$ then A is an isomorphism.*

Proof:

Step 1: A is injective:

$$\|Au\| \|u\| \stackrel{C.S.}{\geq} |\langle Au, u \rangle| = |a(u, u)| \geq \lambda \|u\|^2$$

$$\Rightarrow \|Au\| \geq \lambda \|u\| \text{ for all } u \in V.$$

Step 2: $A(V)$ is dense in V . Let $u \in V$ such that

$$\langle Au, v \rangle = 0 \quad \forall v \in V$$

$$\text{take } v = u \Rightarrow a(u, u) = 0 \Rightarrow u = 0.$$

Step 3: $\mathbb{R}(A) = A(V)$ is closed. Let v_n be a sequence in $A(V)$ and let a_n be such that

$$Au_n = v_n$$

$$\stackrel{(+)}{\implies} u_n \text{ is a Cauchy sequence } \Rightarrow u_n \rightarrow u \in V \text{ und } Au_n \rightarrow Au \Rightarrow v_n \rightarrow Au \in A(V)$$

Step 4: $u = A^{-1}v \stackrel{(+)}{\implies} \|A^{-1}v\| \leq \lambda^{-1}\|v\| \quad \forall v \in V.$

□

Next we consider two Hilbert spaces V, H with $V \subset H$ (the inclusion is continuous), i.e.

$$\exists c < \infty: \quad \|u\|_H \leq c\|u\|_V \quad \forall u \in V$$

and we assume that V is dense in H .

Example: $V = W^{1,1}(\mathbb{R}^n)$, $H = L^2(\mathbb{R}^n)$

$$\|u\|_L^2 \leq \|u\|_{W^{1,2}}$$

There exists a natural injection from H into V' . Let $h \in H$ then $V \ni u \mapsto \langle u, h \rangle_H$ is continuous on $V \xrightarrow[\text{Thm. II.1}]{\implies} \exists l_h \in V'$:

$$l_h(u) = \langle u, h \rangle_H \quad \forall u \in V$$

injectivity follows from density of V in H . $V \subseteq H$

Stichwortverzeichnis

bounded, 3

closable, 4

closed, 3

closure, 4

coercive, 9

dense, 3

domain, 2

Graph, 3

inner product, 2

Operator

 bounded, 3

 closable, 4

 closed, 3

 closure of, 4

range, 2