

A van Galen 2019 AML (hdWGCNA networks): perturbation score, antagonist mode
Single-gene targets
Joint targets
log10 score
-3
-4
-5
-6
-7
B
Plasma
T
CTL
NK
earlyEry
lateEry
HSC
HSC-like
Prog
Prog-like
GMP
GMP-like
ProMono
ProMono-like
Mono
Mono-like
cDC
cDC-like
AURKA
AURKB
BCL2
CD33
CDK19
CDK8
CDK9
CHEK1
DHODH
DNMT1
EIF4A1
EIF4E
FLT3
GLS
HAVCR2
HDAC1
HMGCR
IDH1
IDH2
IL3RA
JAK2
KDM1A
KIT
MAP2K1
MDM2
MEN1
MTOR
NAE1
PLK1
PSMB5
SYK
TP53
WEE1
XPO1
BCL2;DNMT1
BCL2;EIF4A1
BCL2;FLT3
BCL2;IDH1
BCL2;IDH2
BCL2;MDM2
BCL2;MEN1
BCL2;XPO1
CDK19;BCL2
CDK8;BCL2
CDK8;CDK19
DNMT1;HAVCR2
DNMT1;HDAC1
DNMT1;IDH1
DNMT1;IDH2
DNMT1;MEN1
DNMT1;NAE1
DNMT1;TP53
DNMT1;XPO1
FLT3;DNMT1
FLT3;MEN1
B Malignant vs. normal counterpart
log10 (malignant / normal score)
2.5
0.0
-2.5
-5.0
HSC
Prog
GMP
ProMono
Mono
cDC
Malignant-like vs. matched normal identity
FLT3 / KIT / IL3RA / CD33
BCL2
C Identity-wide offset
Identity effect, log10 (vs. B)
0.5
0.0
B
Plasma
T
CTL
NK
earlyEry
lateEry
HSC
HSC-like
Prog
Prog-like
GMP
GMP-like
ProMono
ProMono-like
Mono
Mono-like
cDC
cDC-like
malignant
normal counterpart
other