

Linear Systems: Stability



1. Definitions in Lyapunov stability analysis
2. Stability of LTI systems: method of eigenvalue/pole locations
3. Lyapunov's approach to stability
 - Relevant tools
 - Lyapunov stability theorems
 - Instability theorem
 - Discrete-time case
4. Recap

Finite dimensional vector norms

Let $v \in \mathbb{R}^n$. A norm is:

Finite dimensional vector norms

Let $v \in \mathbb{R}^n$. A norm is:

- ▶ a metric in vector space: a function that assigns a real-valued length to each vector in a vector space

Finite dimensional vector norms

Let $v \in \mathbb{R}^n$. A norm is:

- ▶ a metric in vector space: a function that assigns a real-valued length to each vector in a vector space
- ▶ e.g., 2 (Euclidean) norm: $\|v\|_2 = \sqrt{v^T v} = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$

Finite dimensional vector norms

Let $v \in \mathbb{R}^n$. A norm is:

- ▶ a metric in vector space: a function that assigns a real-valued length to each vector in a vector space

- ▶ e.g., 2 (Euclidean) norm: $\|v\|_2 = \sqrt{v^T v} = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$

default in this set of notes: $\|\cdot\| = \|\cdot\|_2$

Equilibrium state

For an n -th order unforced system

$$\dot{x} = f(x, t), \quad x(t_0) = x_0$$

Equilibrium state

For an n -th order unforced system

$$\dot{x} = f(x, t), \quad x(t_0) = x_0$$

an equilibrium state/point x_e is one such that

$$f(x_e, t) = 0, \quad \forall t$$

Equilibrium state

For an n -th order unforced system

$$\dot{x} = f(x, t), \quad x(t_0) = x_0$$

an equilibrium state/point x_e is one such that

$$f(x_e, t) = 0, \quad \forall t$$

- the condition must be satisfied by all $t \geq 0$

Equilibrium state

For an n -th order unforced system

$$\dot{x} = f(x, t), \quad x(t_0) = x_0$$

an equilibrium state/point x_e is one such that

$$f(x_e, t) = 0, \quad \forall t$$

- ▶ the condition must be satisfied by all $t \geq 0$
- ▶ if a system starts at equilibrium state, it stays there

Equilibrium state of a linear system

For a linear system

$$\dot{x}(t) = A(t)x(t), \quad x(t_0) = x_0$$

Equilibrium state of a linear system

For a linear system

$$\dot{x}(t) = A(t)x(t), \quad x(t_0) = x_0$$

- ▶ origin $x_e = 0$ is always an equilibrium state

Equilibrium state of a linear system

For a linear system

$$\dot{x}(t) = A(t)x(t), \quad x(t_0) = x_0$$

- ▶ origin $x_e = 0$ is always an equilibrium state
- ▶ when $A(t)$ is singular, multiple equilibrium states exist

Lyapunov's definition of stability

- ▶ The equilibrium state 0 of $\dot{x} = f(x, t)$ is *stable in the sense of Lyapunov (s.i.L)* if

Lyapunov's definition of stability

- The equilibrium state 0 of $\dot{x} = f(x, t)$ is *stable in the sense of Lyapunov (s.i.L)* if for all $\epsilon > 0$, and t_0 , there exists $\delta(\epsilon, t_0) > 0$

Lyapunov's definition of stability

- The equilibrium state 0 of $\dot{x} = f(x, t)$ is *stable in the sense of Lyapunov (s.i.L)* if for all $\epsilon > 0$, and t_0 , there exists $\delta(\epsilon, t_0) > 0$ such that $\|x(t_0)\|_2 < \delta$ gives $\|x(t)\|_2 < \epsilon$ for all $t \geq t_0$

Lyapunov's definition of stability

- The equilibrium state 0 of $\dot{x} = f(x, t)$ is *stable in the sense of Lyapunov (s.i.L)* if for all $\epsilon > 0$, and t_0 , there exists $\delta(\epsilon, t_0) > 0$ such that $\|x(t_0)\|_2 < \delta$ gives $\|x(t)\|_2 < \epsilon$ for all $t \geq t_0$




Figure: Stable s.i.L: $\|x(t_0)\| < \delta \Rightarrow \|x(t)\| < \epsilon \forall t \geq t_0$.

Asymptotic stability

The equilibrium state 0 of $\dot{x} = f(x, t)$ is asymptotically stable if

Asymptotic stability

The equilibrium state 0 of $\dot{x} = f(x, t)$ is asymptotically stable if

- ▶ it is stable in the sense of Lyapunov, and

Asymptotic stability

The equilibrium state 0 of $\dot{x} = f(x, t)$ is asymptotically stable if

- ▶ it is stable in the sense of Lyapunov, and
- ▶ for all $\epsilon > 0$ and t_0 , there exists $\delta(\epsilon, t_0) > 0$ such that $\|x(t_0)\|_2 < \delta$ gives $x(t) \rightarrow 0$

Asymptotic stability

The equilibrium state 0 of $\dot{x} = f(x, t)$ is asymptotically stable if

- ▶ it is stable in the sense of Lyapunov, and
- ▶ for all $\epsilon > 0$ and t_0 , there exists $\delta(\epsilon, t_0) > 0$ such that $\|x(t_0)\|_2 < \delta$ gives $x(t) \rightarrow 0$




Figure: Asymptotically stable i.s.L: $\|x(t_0)\| < \delta \Rightarrow \|x(t)\| \rightarrow 0$.

1. Definitions in Lyapunov stability analysis
2. Stability of LTI systems: method of eigenvalue/pole locations
3. Lyapunov's approach to stability
 - Relevant tools
 - Lyapunov stability theorems
 - Instability theorem
 - Discrete-time case
4. Recap

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

- ▶ the response $e^{At}x(0)$ involves modes such as $e^{\lambda t}$, $te^{\lambda t}$, $e^{\sigma t} \cos \omega t$, $e^{\sigma t} \sin \omega t$

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

- ▶ the response $e^{At}x(0)$ involves modes such as $e^{\lambda t}$, $te^{\lambda t}$, $e^{\sigma t} \cos \omega t$, $e^{\sigma t} \sin \omega t$
- ▶ the response $A^k x(0)$ involves modes such as λ^k , $k\lambda^{k-1}$, $r^k \cos k\theta$, $r^k \sin k\theta$

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

- ▶ the response $e^{At}x(0)$ involves modes such as $e^{\lambda t}$, $te^{\lambda t}$, $e^{\sigma t} \cos \omega t$, $e^{\sigma t} \sin \omega t$
- ▶ the response $A^k x(0)$ involves modes such as λ^k , $k\lambda^{k-1}$, $r^k \cos k\theta$, $r^k \sin k\theta$
- ▶ $e^{\sigma t} \rightarrow 0$ if $\sigma < 0$; $e^{\lambda t} \rightarrow 0$ if $\lambda < 0$
- ▶ $\lambda^k \rightarrow 0$ if $|\lambda| < 1$; $r^k \rightarrow 0$ if $|r| = |\sqrt{\sigma^2 + \omega^2}| = |\lambda| < 1$

Stability of the origin for $\dot{x} = Ax$

stability at 0	$\lambda_i(A)$
unstable	$\text{Re}\{\lambda_i\} > 0$ for some λ_i or $\text{Re}\{\lambda_i\} \leq 0$ for all λ_i 's but for a repeated λ_m on the imaginary axis with multiplicity m , nullity $(A - \lambda_m I) < m$ (Jordan form)
stable i.s.L	$\text{Re}\{\lambda_i\} \leq 0$ for all λ_i 's and \forall repeated λ_m on the imaginary axis with multiplicity m , nullity $(A - \lambda_m I) = m$ (diagonal form)
asymptotically stable	$\text{Re}\{\lambda_i\} < 0 \forall \lambda_i$ (A is then called Hurwitz stable)

Example (Unstable moving mass)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Example (Unstable moving mass)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

► $\lambda_1 = \lambda_2 = 0$, $m = 2$,

$$\text{nullity}(A - \lambda_i I) = \text{nullity} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = 1 < m$$

Example (Unstable moving mass)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

- ▶ $\lambda_1 = \lambda_2 = 0, m = 2,$
 $\text{nullity}(A - \lambda_i I) = \text{nullity} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = 1 < m$
- ▶ i.e., two repeated eigenvalues but needs a generalized eigenvector \Rightarrow Jordan form after similarity transform

Example (Unstable moving mass)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

- ▶ $\lambda_1 = \lambda_2 = 0$, $m = 2$,
 $\text{nullity}(A - \lambda_i I) = \text{nullity} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = 1 < m$
- ▶ i.e., two repeated eigenvalues but needs a generalized eigenvector \Rightarrow Jordan form after similarity transform
- ▶ verify by checking $e^{At} = \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}$: t grows unbounded

Example (Stable in the sense of Lyapunov)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Example (Stable in the sense of Lyapunov)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

► $\lambda_1 = \lambda_2 = 0, \quad m = 2,$

$$\text{nullity}(A - \lambda_i I) = \text{nullity} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 2 = m$$

Example (Stable in the sense of Lyapunov)

$$\dot{x} = Ax, \quad A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

- ▶ $\lambda_1 = \lambda_2 = 0, \quad m = 2,$
 $\text{nullity}(A - \lambda_i I) = \text{nullity} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 2 = m$
- ▶ verify by checking $e^{At} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Routh-Hurwitz criterion

- ▶ the Routh Test (by E.J. Routh, in 1877): a simple algebraic procedure to determine how many roots a given polynomial

$$A(s) = a_n s^n + a_{n-1} s^{n-1} + \cdots + a_1 s + a_0$$

has in the closed right-half complex plane, without the need to explicitly solve for the roots

Routh-Hurwitz criterion

- ▶ the Routh Test (by E.J. Routh, in 1877): a simple algebraic procedure to determine how many roots a given polynomial

$$A(s) = a_n s^n + a_{n-1} s^{n-1} + \cdots + a_1 s + a_0$$

has in the closed right-half complex plane, without the need to explicitly solve for the roots

- ▶ German mathematician Adolf Hurwitz independently proposed in 1895 to approach the problem from a matrix perspective

Routh-Hurwitz criterion

- ▶ the Routh Test (by E.J. Routh, in 1877): a simple algebraic procedure to determine how many roots a given polynomial

$$A(s) = a_n s^n + a_{n-1} s^{n-1} + \cdots + a_1 s + a_0$$

has in the closed right-half complex plane, without the need to explicitly solve for the roots

- ▶ German mathematician Adolf Hurwitz independently proposed in 1895 to approach the problem from a matrix perspective
- ▶ popular if stability is the only concern and no details on eigenvalues (e.g., speed of response) are needed

Routh-Hurwitz criterion

- ▶ the asymptotic stability of the equilibrium point 0 for $\dot{x} = Ax$ can also be concluded based on the Routh-Hurwitz criterion

Routh-Hurwitz criterion

- ▶ the asymptotic stability of the equilibrium point 0 for $\dot{x} = Ax$ can also be concluded based on the Routh-Hurwitz criterion
- ▶ simply apply the Routh Test to $A(s) = \det(sI - A)$

Routh-Hurwitz criterion

- ▶ the asymptotic stability of the equilibrium point 0 for $\dot{x} = Ax$ can also be concluded based on the Routh-Hurwitz criterion
- ▶ simply apply the Routh Test to $A(s) = \det(sI - A)$
- ▶ recap: the poles of transfer function $G(s) = C(sI - A)^{-1}B + D$ come from $\det(sI - A)$ in computing the inverse $(sI - A)^{-1}$

The Routh Array

for $A(s) = a_n s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0$, construct

$$\begin{array}{c|cccccc} s^n & a_n & a_{n-2} & a_{n-4} & a_{n-6} & \cdots \\ s^{n-1} & a_{n-1} & a_{n-3} & a_{n-5} & a_{n-7} & \cdots \\ s^{n-2} & q_{n-2} & q_{n-4} & q_{n-6} & \cdots & \\ s^{n-3} & q_{n-3} & q_{n-5} & q_{n-7} & \cdots & \\ \vdots & \vdots & \vdots & \vdots & & \\ s^1 & x_2 & x_0 & & & \\ s^0 & x_0 & & & & \end{array}$$

The Routh Array

for $A(s) = a_n s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0$, construct

$$\begin{array}{c|cccc} s^n & a_n & a_{n-2} & a_{n-4} & a_{n-6} & \dots \\ s^{n-1} & a_{n-1} & a_{n-3} & a_{n-5} & a_{n-7} & \dots \\ s^{n-2} & q_{n-2} & q_{n-4} & q_{n-6} & \dots & \\ s^{n-3} & q_{n-3} & q_{n-5} & q_{n-7} & \dots & \\ \vdots & \vdots & \vdots & \vdots & & \\ s^1 & x_2 & x_0 & & & \\ s^0 & x_0 & & & & \end{array}$$

- first two rows contain the coefficients of $A(s)$

The Routh Array

for $A(s) = a_n s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0$, construct

$$\begin{array}{c|cccc}
 s^n & a_n & a_{n-2} & a_{n-4} & a_{n-6} & \cdots \\
 s^{n-1} & a_{n-1} & a_{n-3} & a_{n-5} & a_{n-7} & \cdots \\
 s^{n-2} & q_{n-2} & q_{n-4} & q_{n-6} & \cdots & \\
 s^{n-3} & q_{n-3} & q_{n-5} & q_{n-7} & \cdots & \\
 \vdots & \vdots & \vdots & \vdots & & \\
 s^1 & x_2 & x_0 & & & \\
 s^0 & x_0 & & & &
 \end{array}$$

- ▶ first two rows contain the coefficients of $A(s)$
- ▶ third row constructed from the previous two rows via

$$\begin{array}{c|ccc}
 \cdot & a & b & x & \cdot \\
 \cdot & c & d & y & \cdot \\
 \cdot & \frac{bc - ad}{c} & \frac{xc - ay}{c} & \cdot & \\
 \cdot & c & c & \cdot & \\
 \cdot & \cdot & \cdot & \cdot & \cdot
 \end{array}$$

The Routh Array

for $A(s) = a_n s^n + a_{n-1} s^{n-1} + \cdots + a_1 s + a_0$, construct

$$\begin{array}{c|ccccc} s^n & a_n & a_{n-2} & a_{n-4} & a_{n-6} & \cdots \\ s^{n-1} & a_{n-1} & a_{n-3} & a_{n-5} & a_{n-7} & \cdots \\ s^{n-2} & q_{n-2} & q_{n-4} & q_{n-6} & \cdots & \\ s^{n-3} & q_{n-3} & q_{n-5} & q_{n-7} & \cdots & \\ \vdots & \vdots & \vdots & \vdots & & \\ s^1 & x_2 & x_0 & & & \\ s^0 & x_0 & & & & \end{array}$$

The Routh Array

for $A(s) = a_n s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0$, construct

$$\begin{array}{c|cccc} s^n & a_n & a_{n-2} & a_{n-4} & a_{n-6} & \cdots \\ s^{n-1} & a_{n-1} & a_{n-3} & a_{n-5} & a_{n-7} & \cdots \\ s^{n-2} & q_{n-2} & q_{n-4} & q_{n-6} & \cdots & \\ s^{n-3} & q_{n-3} & q_{n-5} & q_{n-7} & \cdots & \\ \vdots & \vdots & \vdots & \vdots & & \\ s^1 & x_2 & x_0 & & & \\ s^0 & x_0 & & & & \end{array}$$

- All roots of $A(s)$ are on the left half s-plane if and only if all elements of the first column of the Routh array are positive.

The Routh Array

Example ($A(s) = 2s^4 + s^3 + 3s^2 + 5s + 10$)

$$\begin{array}{c|ccc} s^4 & 2 & 3 & 10 \\ s^3 & 1 & 5 & 0 \\ s^2 & 3 - \frac{2 \times 5}{1} = -7 & 10 & 0 \\ s^1 & 5 - \frac{1 \times 10}{-7} & 0 & 0 \\ s^0 & 10 & 0 & 0 \end{array}$$

The Routh Array

Example ($A(s) = 2s^4 + s^3 + 3s^2 + 5s + 10$)

$$\begin{array}{c|ccc} s^4 & 2 & 3 & 10 \\ s^3 & 1 & 5 & 0 \\ s^2 & 3 - \frac{2 \times 5}{1} = -7 & 10 & 0 \\ s^1 & 5 - \frac{1 \times 10}{-7} & 0 & 0 \\ s^0 & 10 & 0 & 0 \end{array}$$

- two sign changes in the first column

The Routh Array

Example ($A(s) = 2s^4 + s^3 + 3s^2 + 5s + 10$)

$$\begin{array}{c|ccc} s^4 & 2 & 3 & 10 \\ s^3 & 1 & 5 & 0 \\ s^2 & 3 - \frac{2 \times 5}{1} = -7 & 10 & 0 \\ s^1 & 5 - \frac{1 \times 10}{-7} & 0 & 0 \\ s^0 & 10 & 0 & 0 \end{array}$$

- ▶ two sign changes in the first column
- ▶ unstable and two roots in the right half side of s-plane

The Routh Array

special cases:

- If the 1st element in any one row of Routh's array is zero, one can replace the zero with a small number ϵ and proceed further.

The Routh Array

special cases:

- ▶ If the 1st element in any one row of Routh's array is zero, one can replace the zero with a small number ϵ and proceed further.
- ▶ There are other possible complications, which we will not pursue further. See, e.g. "Automatic Control Systems", by Kuo, 7th ed., pp. 339-340.

Stability of the origin for $x(k+1) = f(x(k), k)$

- stability analysis follows analogously for nonlinear time-varying discrete-time systems of the form

$$x(k+1) = f(x(k), k), \quad x(k_0) = x_0$$

Stability of the origin for $x(k+1) = f(x(k), k)$

- ▶ stability analysis follows analogously for nonlinear time-varying discrete-time systems of the form

$$x(k+1) = f(x(k), k), \quad x(k_0) = x_0$$

- ▶ equilibrium point x_e :

$$f(x_e, k) = x_e, \quad \forall k$$

- ▶ without loss of generality, 0 is assumed an equilibrium point

Stability of the origin for $x(k+1) = Ax(k)$

stability at 0	$\lambda_i(A)$
unstable	$ \lambda_i > 1$ for some λ_i or $ \lambda_i \leq 1$ for all λ_i 's but for a repeated λ_m on the unit circle with multiplicity m , $\text{nullity}(A - \lambda_m I) < m$ (Jordan form)
stable i.s.L	$ \lambda_i \leq 1$ for all λ_i 's but for any repeated λ_m on the unit circle with multiplicity m , $\text{nullity}(A - \lambda_m I) = m$ (diagonal form)
asymptotically stable	$ \lambda_i < 1 \ \forall \lambda_i$ (such a matrix is called Schur stable)

Routh-Hurwitz criterion for DT LTI systems


- ▶ the stability domain $|\lambda_i| < 1$ is a unit disk

Routh-Hurwitz criterion for DT LTI systems

- ▶ the stability domain $|\lambda_i| < 1$ is a unit disk
- ▶ Routh array validates stability in the left-half plane

Routh-Hurwitz criterion for DT LTI systems

- ▶ the stability domain $|\lambda_i| < 1$ is a unit disk
- ▶ Routh array validates stability in the left-half plane
- ▶ bilinear transformation maps the closed left half s -plane to the closed unit disk in z -plane



Routh-Hurwitz criterion for DT LTI systems

- ▶ Given $A(z) = z^n + a_1 z^{n-1} + \dots + a_n$, procedures of Routh-Hurwitz test:
 - ▶ apply bilinear transform
$$A(z)|_{z=\frac{1+s}{1-s}} = \left(\frac{1+s}{1-s}\right)^n + a_1 \left(\frac{1+s}{1-s}\right)^{n-1} + \dots + a_n =: \frac{A^*(s)}{(1-s)^n}$$
 - ▶ apply Routh test to
$$A^*(s) = a_n^* s^n + a_{n-1}^* s^{n-1} + \dots + a_0^* = A(z)|_{z=\frac{1+s}{1-s}} (1-s)^n$$

Routh-Hurwitz criterion for DT LTI systems

Example ($A(z) = z^3 + 0.8z^2 + 0.6z + 0.5$)

$$\blacktriangleright A^*(s) = A(z)|_{z=\frac{1+s}{1-s}} (1-s)^3 = (1+s)^3 + 0.8(1+s)^2(1-s) + 0.6(1+s)(1-s)^2 + 0.5(1-s)^3 = 0.3s^3 + 3.1s^2 + 1.7s + 2.9$$

Routh-Hurwitz criterion for DT LTI systems

Example ($A(z) = z^3 + 0.8z^2 + 0.6z + 0.5$)

► $A^*(s) = A(z)|_{z=\frac{1+s}{1-s}} (1-s)^3 = (1+s)^3 + 0.8(1+s)^2(1-s) + 0.6(1+s)(1-s)^2 + 0.5(1-s)^3 = 0.3s^3 + 3.1s^2 + 1.7s + 2.9$

► Routh array

$$\begin{array}{c|cc} s^3 & 0.3 & 1.7 \\ s^2 & 3.1 & 2.9 \\ s & 1.7 - \frac{0.3 \times 2.9}{3.1} = 1.42 & 0 \\ s^0 & 2.9 & 0 \end{array}$$

Routh-Hurwitz criterion for DT LTI systems

Example ($A(z) = z^3 + 0.8z^2 + 0.6z + 0.5$)

► $A^*(s) = A(z)|_{z=\frac{1+s}{1-s}} (1-s)^3 = (1+s)^3 + 0.8(1+s)^2(1-s) + 0.6(1+s)(1-s)^2 + 0.5(1-s)^3 = 0.3s^3 + 3.1s^2 + 1.7s + 2.9$

► Routh array

$$\begin{array}{c|cc} s^3 & 0.3 & 1.7 \\ s^2 & 3.1 & 2.9 \\ s & 1.7 - \frac{0.3 \times 2.9}{3.1} = 1.42 & 0 \\ s^0 & 2.9 & 0 \end{array}$$

► all elements in first column are positive \Rightarrow roots of $A(z)$ are all in the unit circle

1. Definitions in Lyapunov stability analysis
2. Stability of LTI systems: method of eigenvalue/pole locations
3. Lyapunov's approach to stability
 - Relevant tools
 - Lyapunov stability theorems
 - Instability theorem
 - Discrete-time case
4. Recap

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

- ▶ the response $e^{At}x(0)$ involves modes such as $e^{\lambda t}$, $te^{\lambda t}$, $e^{\sigma t} \cos \omega t$, $e^{\sigma t} \sin \omega t$

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

- ▶ the response $e^{At}x(0)$ involves modes such as $e^{\lambda t}$, $te^{\lambda t}$, $e^{\sigma t} \cos \omega t$, $e^{\sigma t} \sin \omega t$
- ▶ the response $A^k x(0)$ involves modes such as λ^k , $k\lambda^{k-1}$, $r^k \cos k\theta$, $r^k \sin k\theta$

Stability of LTI systems: method of eigenvalue/pole locations

the stability of the equilibrium point 0 for $\dot{x} = Ax$ or $x(k+1) = Ax(k)$ can be concluded immediately based on $\lambda(A)$:

- ▶ the response $e^{At}x(0)$ involves modes such as $e^{\lambda t}$, $te^{\lambda t}$, $e^{\sigma t} \cos \omega t$, $e^{\sigma t} \sin \omega t$
- ▶ the response $A^k x(0)$ involves modes such as λ^k , $k\lambda^{k-1}$, $r^k \cos k\theta$, $r^k \sin k\theta$
- ▶ $e^{\sigma t} \rightarrow 0$ if $\sigma < 0$; $e^{\lambda t} \rightarrow 0$ if $\lambda < 0$
- ▶ $\lambda^k \rightarrow 0$ if $|\lambda| < 1$; $r^k \rightarrow 0$ if $|r| = |\sqrt{\sigma^2 + \omega^2}| = |\lambda| < 1$

Lyapunov's approach to stability

The direct method of Lyapunov to stability problems:

Lyapunov's approach to stability

The direct method of Lyapunov to stability problems:

- ▶ no need for explicit solutions to system responses

Lyapunov's approach to stability

The direct method of Lyapunov to stability problems:

- ▶ no need for explicit solutions to system responses
- ▶ an “energy” perspective

Lyapunov's approach to stability

The direct method of Lyapunov to stability problems:

- ▶ no need for explicit solutions to system responses
- ▶ an “energy” perspective
- ▶ fit for general dynamic systems (linear/nonlinear, time-invariant/time-varying)

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2 : \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2 : \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

► $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2: \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

- ▶ $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable
- ▶ total energy

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2: \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

- ▶ $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable
- ▶ total energy

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

- ▶ energy dissipates / is dissipative:

$$\dot{\mathcal{E}}(t) = kx_1\dot{x}_1 + mx_2\dot{x}_2 = -bx_2^2 \leq 0$$

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2: \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

- ▶ $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable
- ▶ total energy

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

- ▶ energy dissipates / is dissipative:

$$\dot{\mathcal{E}}(t) = kx_1\dot{x}_1 + mx_2\dot{x}_2 = -bx_2^2 \leq 0$$

- ▶ $\dot{\mathcal{E}} = 0$ only when $x_2 = 0$.

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2: \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

- ▶ $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable
- ▶ total energy

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

- ▶ energy dissipates / is dissipative:

$$\dot{\mathcal{E}}(t) = kx_1\dot{x}_1 + mx_2\dot{x}_2 = -bx_2^2 \leq 0$$

- ▶ $\dot{\mathcal{E}} = 0$ only when $x_2 = 0$. As $[x_1, x_2]^T = 0$ is the only equilibrium, the motion will not stop at $x_2 = 0, x_1 \neq 0$.

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2: \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

- ▶ $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable
- ▶ total energy

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

- ▶ energy dissipates / is dissipative:

$$\dot{\mathcal{E}}(t) = kx_1\dot{x}_1 + mx_2\dot{x}_2 = -bx_2^2 \leq 0$$

- ▶ $\dot{\mathcal{E}} = 0$ only when $x_2 = 0$. As $[x_1, x_2]^T = 0$ is the only equilibrium, the motion will not stop at $x_2 = 0$, $x_1 \neq 0$. Thus energy will keep decreasing toward 0

Stability from an energy viewpoint: Example

Consider spring-mass-damper systems:

$$\dot{x}_1 = x_2 \quad (x_1: \text{position}; x_2: \text{velocity})$$

$$\dot{x}_2 = -\frac{k}{m}x_1 - \frac{b}{m}x_2, \quad b > 0 \quad (\text{Newton's law})$$

- ▶ $\lambda(A)$'s are in the left-half s -plane \Rightarrow asymptotically stable
- ▶ total energy

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

- ▶ energy dissipates / is dissipative:

$$\dot{\mathcal{E}}(t) = kx_1\dot{x}_1 + mx_2\dot{x}_2 = -bx_2^2 \leq 0$$

- ▶ $\dot{\mathcal{E}} = 0$ only when $x_2 = 0$. As $[x_1, x_2]^T = 0$ is the only equilibrium, the motion will not stop at $x_2 = 0$, $x_1 \neq 0$. Thus energy will keep decreasing toward 0 which is achieved at the origin.

Stability from an energy viewpoint: Generalization

Consider unforced, time-varying, nonlinear systems

$$\begin{aligned}\dot{x}(t) &= f(x(t), t), \quad x(t_0) = x_0 \\ x(k+1) &= f(x(k), k), \quad x(k_0) = x_0\end{aligned}$$

Stability from an energy viewpoint: Generalization

Consider unforced, time-varying, nonlinear systems

$$\begin{aligned}\dot{x}(t) &= f(x(t), t), \quad x(t_0) = x_0 \\ x(k+1) &= f(x(k), k), \quad x(k_0) = x_0\end{aligned}$$

- assume the origin is an equilibrium state

Stability from an energy viewpoint: Generalization

Consider unforced, time-varying, nonlinear systems

$$\begin{aligned}\dot{x}(t) &= f(x(t), t), \quad x(t_0) = x_0 \\ x(k+1) &= f(x(k), k), \quad x(k_0) = x_0\end{aligned}$$

- ▶ assume the origin is an equilibrium state
- ▶ energy function \Rightarrow Lyapunov function: a scalar function of x and t

Stability from an energy viewpoint: Generalization

Consider unforced, time-varying, nonlinear systems

$$\begin{aligned}\dot{x}(t) &= f(x(t), t), \quad x(t_0) = x_0 \\ x(k+1) &= f(x(k), k), \quad x(k_0) = x_0\end{aligned}$$

- ▶ assume the origin is an equilibrium state
- ▶ energy function \Rightarrow Lyapunov function: a scalar function of x and t (or x and k)

Stability from an energy viewpoint: Generalization

Consider unforced, time-varying, nonlinear systems

$$\begin{aligned}\dot{x}(t) &= f(x(t), t), \quad x(t_0) = x_0 \\ x(k+1) &= f(x(k), k), \quad x(k_0) = x_0\end{aligned}$$

- ▶ assume the origin is an equilibrium state
- ▶ energy function \Rightarrow Lyapunov function: a scalar function of x and t (or x and k)
- ▶ goal is to relate properties of the state through the Lyapunov function

Stability from an energy viewpoint: Generalization

Consider unforced, time-varying, nonlinear systems

$$\begin{aligned}\dot{x}(t) &= f(x(t), t), \quad x(t_0) = x_0 \\ x(k+1) &= f(x(k), k), \quad x(k_0) = x_0\end{aligned}$$

- ▶ assume the origin is an equilibrium state
- ▶ energy function \Rightarrow Lyapunov function: a scalar function of x and t (or x and k)
- ▶ goal is to relate properties of the state through the Lyapunov function
- ▶ main tool: matrix formulation, linear algebra, positive definite functions

Relevant tools

Quadratic functions

- intrinsic in energy-like analysis, e.g.

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 = \frac{1}{2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} k & 0 \\ 0 & m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Relevant tools

Quadratic functions

- intrinsic in energy-like analysis, e.g.

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 = \frac{1}{2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} k & 0 \\ 0 & m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- convenience of matrix formulation:

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 + x_1x_2 = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} \frac{k}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{m}{2} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Relevant tools

Quadratic functions

- intrinsic in energy-like analysis, e.g.

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 = \frac{1}{2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} k & 0 \\ 0 & m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- convenience of matrix formulation:

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 + x_1x_2 = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} \frac{k}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{m}{2} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 + x_1x_2 + c = \begin{bmatrix} x_1 \\ x_2 \\ 1 \end{bmatrix}^T \begin{bmatrix} \frac{k}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & \frac{m}{2} & 0 \\ 0 & 0 & c \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ 1 \end{bmatrix}$$

Relevant tools

Quadratic functions

- intrinsic in energy-like analysis, e.g.

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 = \frac{1}{2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} k & 0 \\ 0 & m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- convenience of matrix formulation:

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 + x_1x_2 = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}^T \begin{bmatrix} \frac{k}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{m}{2} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2 + x_1x_2 + c = \begin{bmatrix} x_1 \\ x_2 \\ 1 \end{bmatrix}^T \begin{bmatrix} \frac{k}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & \frac{m}{2} & 0 \\ 0 & 0 & c \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ 1 \end{bmatrix}$$

- general quadratic functions in matrix form

$$Q(x) = x^T P x, \quad P^T = P$$

Relevant tools

Symmetric matrices

- ▶ recall: a real square matrix A is
 - ▶ *symmetric* if $A = A^T$
 - ▶ *skew-symmetric* if $A = -A^T$

Relevant tools

Symmetric matrices

- ▶ recall: a real square matrix A is
 - ▶ *symmetric* if $A = A^T$
 - ▶ *skew-symmetric* if $A = -A^T$
- ▶ examples:

$$\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

Relevant tools

Symmetric matrices

- ▶ recall: a real square matrix A is
 - ▶ *symmetric* if $A = A^T$
 - ▶ *skew-symmetric* if $A = -A^T$
- ▶ examples:

$$\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

- ▶ Any real square matrix can be decomposed as the sum of a *symmetric* matrix and a *skew-symmetric* matrix:

Relevant tools

Symmetric matrices

- ▶ recall: a real square matrix A is
 - ▶ *symmetric* if $A = A^T$
 - ▶ *skew-symmetric* if $A = -A^T$
- ▶ examples:

$$\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

- ▶ Any real square matrix can be decomposed as the sum of a *symmetric* matrix and a *skew-symmetric* matrix:

$$\text{e.g. } \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 2.5 \\ 2.5 & 4 \end{bmatrix} + \begin{bmatrix} 0 & -0.5 \\ 0.5 & 0 \end{bmatrix}$$

Relevant tools

Symmetric matrices

- ▶ recall: a real square matrix A is
 - ▶ *symmetric* if $A = A^T$
 - ▶ *skew-symmetric* if $A = -A^T$
- ▶ examples:

$$\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

- ▶ Any real square matrix can be decomposed as the sum of a *symmetric* matrix and a *skew-symmetric* matrix:

$$\text{e.g. } \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 2.5 \\ 2.5 & 4 \end{bmatrix} + \begin{bmatrix} 0 & -0.5 \\ 0.5 & 0 \end{bmatrix}$$

$$\text{general case: } P = \frac{P + P^T}{2} + \frac{P - P^T}{2}$$

Relevant tools

Symmetric matrices

- ▶ a real square matrix $A \in \mathbb{R}^{n \times n}$ is *orthogonal* if $A^T A = A A^T = I$

Relevant tools

Symmetric matrices

- ▶ a real square matrix $A \in \mathbb{R}^{n \times n}$ is *orthogonal* if $A^T A = A A^T = I$
- ▶ meaning that the columns of A form an orthonormal basis of \mathbb{R}^n

$$A = \begin{bmatrix} | & | & | & | \\ a_1 & a_2 & \dots & a_n \\ | & | & | & | \end{bmatrix}$$

Relevant tools

Symmetric matrices

- ▶ a real square matrix $A \in \mathbb{R}^{n \times n}$ is *orthogonal* if $A^T A = A A^T = I$
- ▶ meaning that the columns of A form an orthonormal basis of \mathbb{R}^n

$$A = \begin{bmatrix} | & | & | & | \\ a_1 & a_2 & \dots & a_n \\ | & | & | & | \end{bmatrix}$$

$$A^T A = \begin{bmatrix} a_1^T a_1 & a_1^T a_2 & \dots & a_1^T a_n \\ a_2^T a_1 & a_2^T a_2 & \dots & a_2^T a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_n^T a_1 & a_n^T a_2 & \dots & a_n^T a_n \end{bmatrix} = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 \end{bmatrix}$$

Relevant tools

Symmetric matrices

- ▶ a real square matrix $A \in \mathbb{R}^{n \times n}$ is *orthogonal* if $A^T A = A A^T = I$
- ▶ meaning that the columns of A form an orthonormal basis of \mathbb{R}^n

$$A = \begin{bmatrix} | & | & | & | \\ a_1 & a_2 & \dots & a_n \\ | & | & | & | \end{bmatrix}$$

$$A^T A = \begin{bmatrix} a_1^T a_1 & a_1^T a_2 & \dots & a_1^T a_n \\ a_2^T a_1 & a_2^T a_2 & \dots & a_2^T a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_n^T a_1 & a_n^T a_2 & \dots & a_n^T a_n \end{bmatrix} = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 \end{bmatrix}$$

namely, $a_j^T a_j = 1$ and $a_j^T a_m = 0 \ \forall j \neq m$.

Theorem

The eigenvalues of symmetric matrices are all real.

Theorem

The eigenvalues of symmetric matrices are all real.

Proof: $\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$.

Eigenvalue-eigenvector pair: $Au = \lambda u$

Theorem

The eigenvalues of symmetric matrices are all real.

Proof: $\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$.

Eigenvalue-eigenvector pair: $Au = \lambda u \Rightarrow \bar{u}^T Au = \lambda \bar{u}^T u$, where \bar{u} is the complex conjugate of u .

Theorem

The eigenvalues of symmetric matrices are all real.

Proof: $\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$.

Eigenvalue-eigenvector pair: $Au = \lambda u \Rightarrow \bar{u}^T Au = \lambda \bar{u}^T u$, where \bar{u} is the complex conjugate of u . $\bar{u}^T Au$ is a real number, as

$$\begin{aligned}\overline{\bar{u}^T Au} &= u^T \overline{A\bar{u}} \\ &= u^T A\bar{u} \quad \because A \in \mathbb{R}^{n \times n} \\ &= u^T A^T \bar{u} \quad \because A = A^T \\ &= \lambda u^T \bar{u} \quad \because (Au)^T = (\lambda u)^T \\ &= \lambda \bar{u}^T u \quad \because u^T \bar{u} \in \mathbb{R} \\ &= \bar{u}^T Au \quad \because Au = \lambda u\end{aligned}$$

Theorem

The eigenvalues of symmetric matrices are all real.

Proof: $\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$.

Eigenvalue-eigenvector pair: $Au = \lambda u \Rightarrow \bar{u}^T Au = \lambda \bar{u}^T u$, where \bar{u} is the complex conjugate of u . $\bar{u}^T Au$ is a real number, as

$$\begin{aligned}\overline{\bar{u}^T Au} &= u^T \overline{A\bar{u}} \\ &= u^T A\bar{u} \quad \because A \in \mathbb{R}^{n \times n} \\ &= u^T A^T \bar{u} \quad \because A = A^T \\ &= \lambda u^T \bar{u} \quad \because (Au)^T = (\lambda u)^T \\ &= \lambda \bar{u}^T u \quad \because u^T \bar{u} \in \mathbb{R} \\ &= \bar{u}^T Au \quad \because Au = \lambda u\end{aligned}$$

Also, $\bar{u}^T u \in \mathbb{R}$.

Theorem

The eigenvalues of symmetric matrices are all real.

Proof: $\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$.

Eigenvalue-eigenvector pair: $Au = \lambda u \Rightarrow \bar{u}^T Au = \lambda \bar{u}^T u$, where \bar{u} is the complex conjugate of u . $\bar{u}^T Au$ is a real number, as

$$\begin{aligned}\overline{\bar{u}^T Au} &= u^T \overline{A\bar{u}} \\ &= u^T A\bar{u} \quad \because A \in \mathbb{R}^{n \times n} \\ &= u^T A^T \bar{u} \quad \because A = A^T \\ &= \lambda u^T \bar{u} \quad \because (Au)^T = (\lambda u)^T \\ &= \lambda \bar{u}^T u \quad \because u^T \bar{u} \in \mathbb{R} \\ &= \bar{u}^T Au \quad \because Au = \lambda u\end{aligned}$$

Also, $\bar{u}^T u \in \mathbb{R}$. Thus $\lambda = \frac{\bar{u}^T Au}{\bar{u}^T u}$ must also be a real number. □

Example

► $\begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}: \lambda = \pm 2$

Example

► $\begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix} : \lambda = \pm 2$

► $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix} : \lambda = 1 \pm 2$

Example

► $\begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix} : \lambda = \pm 2$

► $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix} : \lambda = 1 \pm 2$

Theorem

The eigenvalues of skew-symmetric matrices are all imaginary or zero.

Theorem

The eigenvalues of skew-symmetric matrices are all imaginary or zero.

► $\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$: $\lambda = \pm 2j$

Theorem

The eigenvalues of skew-symmetric matrices are all imaginary or zero.

► $\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$: $\lambda = \pm 2j$

Theorem

All eigenvalues of an orthogonal matrix have a magnitude of 1.

Theorem

All eigenvalues of an orthogonal matrix have a magnitude of 1.

$$\blacktriangleright \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} : \lambda = 1 \pm 2j$$

Theorem

All eigenvalues of an orthogonal matrix have a magnitude of 1.

$$\blacktriangleright \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} : \lambda = 1 \pm 2j$$

Important properties of symmetric matrices

Theorem

The eigenvalues of symmetric matrices are all real.

Theorem

The eigenvalues of skew-symmetric matrices are all imaginary or zero.

Theorem

All eigenvalues of an orthogonal matrix have a magnitude of 1.

Important properties of symmetric matrices

Theorem

The eigenvalues of symmetric matrices are all real.

Theorem

The eigenvalues of skew-symmetric matrices are all imaginary or zero.

Theorem

All eigenvalues of an orthogonal matrix have a magnitude of 1.

matrix structure	analogy in complex plane
symmetric	real line
skew-symmetric	imaginary line
orthogonal	unit circle

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization
 $A = U\Lambda U^{-1}$.

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization $A = U\Lambda U^{-1}$. When A is symmetric, things are even better:

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization $A = U\Lambda U^{-1}$. When A is symmetric, things are even better:

Theorem (Symmetric eigenvalue decomposition (SED))

$\forall : A \in \mathbb{R}^{n \times n}, A^T = A$, there always exist $\lambda_i \in \mathbb{R}$ and $u_i \in \mathbb{R}^n$, s.t.

$$A = \sum_{i=1}^n \lambda_i u_i u_i^T = U\Lambda U^T \quad (1)$$

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization $A = U\Lambda U^{-1}$. When A is symmetric, things are even better:

Theorem (Symmetric eigenvalue decomposition (SED))

$\forall : A \in \mathbb{R}^{n \times n}$, $A^T = A$, there always exist $\lambda_i \in \mathbb{R}$ and $u_i \in \mathbb{R}^n$, s.t.

$$A = \sum_{i=1}^n \lambda_i u_i u_i^T = U\Lambda U^T \quad (1)$$

► λ_i 's: eigenvalues of A

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization $A = U \Lambda U^{-1}$. When A is symmetric, things are even better:

Theorem (Symmetric eigenvalue decomposition (SED))

$\forall : A \in \mathbb{R}^{n \times n}, A^T = A$, there always exist $\lambda_i \in \mathbb{R}$ and $u_i \in \mathbb{R}^n$, s.t.

$$A = \sum_{i=1}^n \lambda_i u_i u_i^T = U \Lambda U^T \quad (1)$$

- ▶ λ_i 's: eigenvalues of A
- ▶ u_i : eigenvector associated to λ_i , normalized to have unity norms

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization $A = U \Lambda U^{-1}$. When A is symmetric, things are even better:

Theorem (Symmetric eigenvalue decomposition (SED))

$\forall : A \in \mathbb{R}^{n \times n}, A^T = A$, there always exist $\lambda_i \in \mathbb{R}$ and $u_i \in \mathbb{R}^n$, s.t.

$$A = \sum_{i=1}^n \lambda_i u_i u_i^T = U \Lambda U^T \quad (1)$$

- ▶ λ_i 's: eigenvalues of A
- ▶ u_i : eigenvector associated to λ_i , normalized to have unity norms
- ▶ $U = [u_1, u_2, \dots, u_n]$ is orthogonal: $U^T U = U U^T = I$

The spectral theorem for symmetric matrices

When $A \in \mathbb{R}^{n \times n}$ has n distinct eigenvalues, we can do diagonalization $A = U\Lambda U^{-1}$. When A is symmetric, things are even better:

Theorem (Symmetric eigenvalue decomposition (SED))

$\forall : A \in \mathbb{R}^{n \times n}$, $A^T = A$, there always exist $\lambda_i \in \mathbb{R}$ and $u_i \in \mathbb{R}^n$, s.t.

$$A = \sum_{i=1}^n \lambda_i u_i u_i^T = U\Lambda U^T \quad (1)$$

- ▶ λ_i 's: eigenvalues of A
- ▶ u_i : eigenvector associated to λ_i , normalized to have unity norms
- ▶ $U = [u_1, u_2, \dots, u_n]$ is orthogonal: $U^T U = U U^T = I$
- ▶ $\Lambda = \text{diagonal}(\lambda_1, \lambda_2, \dots, \lambda_n)$

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$,

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$.

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$.

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$.

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$. So $\lambda_i u_i^T u_j = \lambda_j u_i^T u_j$.

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$. So $\lambda_i u_i^T u_j = \lambda_j u_i^T u_j$. But $\lambda_i \neq \lambda_j$. It must be that $u_i^T u_j = 0$. \square

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$. So $\lambda_i u_i^T u_j = \lambda_j u_i^T u_j$. But $\lambda_i \neq \lambda_j$. It must be that $u_i^T u_j = 0$. \square

SED now follows:

- If A has distinct eigenvalues,

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$. So $\lambda_i u_i^T u_j = \lambda_j u_i^T u_j$. But $\lambda_i \neq \lambda_j$. It must be that $u_i^T u_j = 0$. \square

SED now follows:

- If A has distinct eigenvalues, then $U = [u_1, u_2, \dots, u_n]$ is orthogonal after normalizing all the eigenvectors to unity norm.

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are **orthogonal**.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$. So $\lambda_i u_i^T u_j = \lambda_j u_i^T u_j$. But $\lambda_i \neq \lambda_j$. It must be that $u_i^T u_j = 0$. \square

SED now follows:

- ▶ If A has distinct eigenvalues, then $U = [u_1, u_2, \dots, u_n]$ is orthogonal after normalizing all the eigenvectors to unity norm.
- ▶ If A has $r(< n)$ distinct eigenvalues,

Elements of proof for SED

Theorem

$\forall : A \in \mathbb{R}^{n \times n}$ with $A^T = A$, then eigenvectors of A , associated with different eigenvalues, are *orthogonal*.

Proof.

Let $Au_i = \lambda_i u_i$ and $Au_j = \lambda_j u_j$. Then $u_i^T Au_j = u_i^T \lambda_j u_j = \lambda_j u_i^T u_j$. Also, $u_i^T Au_j = u_i^T A^T u_j = (Au_i)^T u_j = \lambda_i u_i^T u_j$. So $\lambda_i u_i^T u_j = \lambda_j u_i^T u_j$. But $\lambda_i \neq \lambda_j$. It must be that $u_i^T u_j = 0$. \square

SED now follows:

- ▶ If A has distinct eigenvalues, then $U = [u_1, u_2, \dots, u_n]$ is orthogonal after normalizing all the eigenvectors to unity norm.
- ▶ If A has $r(< n)$ distinct eigenvalues, we can *choose* multiple orthogonal eigenvectors for the eigenvalues with none-unity multiplicities.

Rethinking symmetric matrices

With the spectral theorem, next time we see a symmetric matrix A , we immediately know that

Rethinking symmetric matrices

With the spectral theorem, next time we see a symmetric matrix A , we immediately know that

- ▶ λ_i is real for all i

Rethinking symmetric matrices

With the spectral theorem, next time we see a symmetric matrix A , we immediately know that

- ▶ λ_i is real for all i
- ▶ associated with λ_i , we can always find a real eigenvector

Rethinking symmetric matrices

With the spectral theorem, next time we see a symmetric matrix A , we immediately know that

- ▶ λ_i is real for all i
- ▶ associated with λ_i , we can always find a real eigenvector
- ▶ \exists an orthonormal basis $\{u_i\}_{i=1}^n$, which consists of the eigenvectors

Rethinking symmetric matrices

With the spectral theorem, next time we see a symmetric matrix A , we immediately know that

- ▶ λ_i is real for all i
- ▶ associated with λ_i , we can always find a real eigenvector
- ▶ \exists an orthonormal basis $\{u_i\}_{i=1}^n$, which consists of the eigenvectors
- ▶ if $A \in \mathbb{R}^{2 \times 2}$, then if you compute first λ_1 , λ_2 and u_1 , you won't need to go through the regular math to get u_2 ,

Rethinking symmetric matrices

With the spectral theorem, next time we see a symmetric matrix A , we immediately know that

- ▶ λ_i is real for all i
- ▶ associated with λ_i , we can always find a real eigenvector
- ▶ \exists an orthonormal basis $\{u_i\}_{i=1}^n$, which consists of the eigenvectors
- ▶ if $A \in \mathbb{R}^{2 \times 2}$, then if you compute first λ_1 , λ_2 and u_1 , you won't need to go through the regular math to get u_2 , but can simply solve for a u_2 that is orthogonal to u_1 with $\|u_2\| = 1$.

Example: $A = \begin{bmatrix} 5 & \sqrt{3} \\ \sqrt{3} & 7 \end{bmatrix}$

Computing the eigenvalues gives

$$\det \begin{bmatrix} 5 - \lambda & \sqrt{3} \\ \sqrt{3} & 7 - \lambda \end{bmatrix} = 35 - 12\lambda + \lambda^2 - 3 = (\lambda - 4)(\lambda - 8) = 0$$
$$\Rightarrow \lambda_1 = 4, \lambda_2 = 8$$

Example: $A = \begin{bmatrix} 5 & \sqrt{3} \\ \sqrt{3} & 7 \end{bmatrix}$

Computing the eigenvalues gives

$$\det \begin{bmatrix} 5 - \lambda & \sqrt{3} \\ \sqrt{3} & 7 - \lambda \end{bmatrix} = 35 - 12\lambda + \lambda^2 - 3 = (\lambda - 4)(\lambda - 8) = 0$$
$$\Rightarrow \lambda_1 = 4, \lambda_2 = 8$$

► first normalized eigenvector:

$$(A - \lambda_1 I) t_1 = 0 \Rightarrow \begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & 3 \end{bmatrix} t_1 = 0 \Rightarrow t_1 = \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix}$$

Example: $A = \begin{bmatrix} 5 & \sqrt{3} \\ \sqrt{3} & 7 \end{bmatrix}$

Computing the eigenvalues gives

$$\det \begin{bmatrix} 5 - \lambda & \sqrt{3} \\ \sqrt{3} & 7 - \lambda \end{bmatrix} = 35 - 12\lambda + \lambda^2 - 3 = (\lambda - 4)(\lambda - 8) = 0$$
$$\Rightarrow \lambda_1 = 4, \lambda_2 = 8$$

► first normalized eigenvector:

$$(A - \lambda_1 I) t_1 = 0 \Rightarrow \begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & 3 \end{bmatrix} t_1 = 0 \Rightarrow t_1 = \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix}$$

► A is symmetric \Rightarrow eigenvectors are orthogonal to each other:

Example: $A = \begin{bmatrix} 5 & \sqrt{3} \\ \sqrt{3} & 7 \end{bmatrix}$

Computing the eigenvalues gives

$$\det \begin{bmatrix} 5 - \lambda & \sqrt{3} \\ \sqrt{3} & 7 - \lambda \end{bmatrix} = 35 - 12\lambda + \lambda^2 - 3 = (\lambda - 4)(\lambda - 8) = 0$$
$$\Rightarrow \lambda_1 = 4, \lambda_2 = 8$$

► first normalized eigenvector:

$$(A - \lambda_1 I) t_1 = 0 \Rightarrow \begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & 3 \end{bmatrix} t_1 = 0 \Rightarrow t_1 = \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix}$$

► A is symmetric \Rightarrow eigenvectors are orthogonal to each other:

choose $t_2 = \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix}$.

Example: $A = \begin{bmatrix} 5 & \sqrt{3} \\ \sqrt{3} & 7 \end{bmatrix}$

Computing the eigenvalues gives

$$\det \begin{bmatrix} 5 - \lambda & \sqrt{3} \\ \sqrt{3} & 7 - \lambda \end{bmatrix} = 35 - 12\lambda + \lambda^2 - 3 = (\lambda - 4)(\lambda - 8) = 0$$
$$\Rightarrow \lambda_1 = 4, \lambda_2 = 8$$

► first normalized eigenvector:

$$(A - \lambda_1 I) t_1 = 0 \Rightarrow \begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & 3 \end{bmatrix} t_1 = 0 \Rightarrow t_1 = \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix}$$

► A is symmetric \Rightarrow eigenvectors are orthogonal to each other:

choose $t_2 = \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix}$. No need to solve $(A - \lambda_2 I) t_2 = 0$!

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Proof.

Perform SED to get $A = \sum_{i=1}^n \lambda_i u_i u_i^T$

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Proof.

Perform SED to get $A = \sum_{i=1}^n \lambda_i u_i u_i^T$ where $\{u_i\}_{i=1}^n$ spans \mathbb{R}^n .

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Proof.

Perform SED to get $A = \sum_{i=1}^n \lambda_i u_i u_i^T$ where $\{u_i\}_{i=1}^n$ spans \mathbb{R}^n . Then any vector $x \in \mathbb{R}^n$ can be decomposed as $x = \sum_{i=1}^n \alpha_i u_i$.

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Proof.

Perform SED to get $A = \sum_{i=1}^n \lambda_i u_i u_i^T$ where $\{u_i\}_{i=1}^n$ spans \mathbb{R}^n . Then any vector $x \in \mathbb{R}^n$ can be decomposed as $x = \sum_{i=1}^n \alpha_i u_i$. Thus

$$\max_{x \neq 0} \frac{x^T A x}{\|x\|_2^2} = \max_{\alpha_i} \frac{(\sum_i \alpha_i u_i)^T \sum_i \lambda_i \alpha_i u_i}{\sum_i \alpha_i^2}$$

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Proof.

Perform SED to get $A = \sum_{i=1}^n \lambda_i u_i u_i^T$ where $\{u_i\}_{i=1}^n$ spans \mathbb{R}^n . Then any vector $x \in \mathbb{R}^n$ can be decomposed as $x = \sum_{i=1}^n \alpha_i u_i$. Thus

$$\max_{x \neq 0} \frac{x^T A x}{\|x\|_2^2} = \max_{\alpha_i} \frac{(\sum_i \alpha_i u_i)^T \sum_i \lambda_i \alpha_i u_i}{\sum_i \alpha_i^2} = \max_{\alpha_i} \frac{\sum_i \lambda_i \alpha_i^2}{\sum_i \alpha_i^2}$$

Theorem (Eigenvalues of symmetric matrices)

If $A = A^T \in \mathbb{R}^{n \times n}$, then the eigenvalues of A satisfy

$$\lambda_{\max} = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (2)$$

$$\lambda_{\min} = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (3)$$

Proof.

Perform SED to get $A = \sum_{i=1}^n \lambda_i u_i u_i^T$ where $\{u_i\}_{i=1}^n$ spans \mathbb{R}^n . Then any vector $x \in \mathbb{R}^n$ can be decomposed as $x = \sum_{i=1}^n \alpha_i u_i$. Thus

$$\max_{x \neq 0} \frac{x^T A x}{\|x\|_2^2} = \max_{\alpha_i} \frac{(\sum_i \alpha_i u_i)^T \sum_i \lambda_i \alpha_i u_i}{\sum_i \alpha_i^2} = \max_{\alpha_i} \frac{\sum_i \lambda_i \alpha_i^2}{\sum_i \alpha_i^2} = \lambda_{\max}$$

□

Positive definite matrices

- ▶ eigenvalues of symmetric matrices are real \Rightarrow we can order the eigenvalues

Positive definite matrices

- ▶ eigenvalues of symmetric matrices are real \Rightarrow we can order the eigenvalues
- ▶ a symmetric matrix P is called positive-definite if all its eigenvalues are positive

Positive definite matrices

- ▶ eigenvalues of symmetric matrices are real \Rightarrow we can order the eigenvalues
- ▶ a symmetric matrix P is called positive-definite if all its eigenvalues are positive
- ▶ equivalently:

Definition (Positive Definite Matrices)

A symmetric matrix $P \in \mathbb{R}^{n \times n}$ is called **positive-definite**, written $P \succ 0$,

Positive definite matrices

- ▶ eigenvalues of symmetric matrices are real \Rightarrow we can order the eigenvalues
- ▶ a symmetric matrix P is called positive-definite if all its eigenvalues are positive
- ▶ equivalently:

Definition (Positive Definite Matrices)

A symmetric matrix $P \in \mathbb{R}^{n \times n}$ is called **positive-definite**, written $P \succ 0$, if $x^T P x > 0$ for all $x (\neq 0) \in \mathbb{R}^n$.

Positive definite matrices

- ▶ eigenvalues of symmetric matrices are real \Rightarrow we can order the eigenvalues
- ▶ a symmetric matrix P is called positive-definite if all its eigenvalues are positive
- ▶ equivalently:

Definition (Positive Definite Matrices)

A symmetric matrix $P \in \mathbb{R}^{n \times n}$ is called **positive-definite**, written $P \succ 0$, if $x^T P x > 0$ for all $x (\neq 0) \in \mathbb{R}^n$.

P is called **positive-semidefinite**, written $P \succeq 0$, if $x^T P x \geq 0$ for all $x \in \mathbb{R}^n$

Positive definite matrices

- ▶ eigenvalues of symmetric matrices are real \Rightarrow we can order the eigenvalues
- ▶ a symmetric matrix P is called **positive-definite** if all its eigenvalues are positive
- ▶ equivalently:

Definition (Positive Definite Matrices)

A symmetric matrix $P \in \mathbb{R}^{n \times n}$ is called **positive-definite**, written $P \succ 0$, if $x^T P x > 0$ for all $x (\neq 0) \in \mathbb{R}^n$.

P is called **positive-semidefinite**, written $P \succeq 0$, if $x^T P x \geq 0$ for all $x \in \mathbb{R}^n$

- ▶ $P \succ 0$ ($P \succeq 0$) $\Leftrightarrow P$ can be decomposed as $P = N^T N$ where N is nonsingular (singular)

Negative definite matrices

Definition

A symmetric matrix $Q \in \mathbb{R}^{n \times n}$ is called **negative-definite**, written $Q \prec 0$, if $-Q \succ 0$, i.e., $x^T Q x < 0$ for all $x (\neq 0) \in \mathbb{R}^n$.

Negative definite matrices

Definition

A symmetric matrix $Q \in \mathbb{R}^{n \times n}$ is called **negative-definite**, written $Q \prec 0$, if $-Q \succ 0$, i.e., $x^T Q x < 0$ for all $x (\neq 0) \in \mathbb{R}^n$.

Q is called **negative-semidefinite**, written $Q \preceq 0$, if $x^T Q x \leq 0$ for all $x \in \mathbb{R}^n$

Updated matrix analogies

matrix structure	eigenvalues	analogy in complex plane
symmetric	real	real axis
skew-symmetric	on imaginary axis	imaginary axis
orthogonal	magnitude 1	unit circle
positive definite	positive	\mathbb{R}_+ axis
negative definite	negative	\mathbb{R}_- axis

Caution

- ▶ positive-definite matrices can have negative entries:

Caution

- positive-definite matrices can have negative entries:

Example

$P = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ is positive-definite, as $P = P^T$ and take any $v = [x, y]^T$, we have

$$\begin{aligned} v^T P v &= \begin{bmatrix} x \\ y \end{bmatrix}^T \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 2x^2 + 2y^2 - 2xy \\ &= x^2 + y^2 + (x - y)^2 \geq 0 \end{aligned}$$

Caution

- positive-definite matrices can have negative entries:

Example

$P = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ is positive-definite, as $P = P^T$ and take any $v = [x, y]^T$, we have

$$\begin{aligned} v^T P v &= \begin{bmatrix} x \\ y \end{bmatrix}^T \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 2x^2 + 2y^2 - 2xy \\ &= x^2 + y^2 + (x - y)^2 \geq 0 \end{aligned}$$

and the equality sign holds only when $x = y = 0$.

Caution

- conversely, matrices whose entries are all positive are not necessarily positive-definite:

Example

$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ is not positive-definite:

Caution

- conversely, matrices whose entries are all positive are not necessarily positive-definite:

Example

$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ is not positive-definite:

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = -2 < 0$$

Positive definite matrices

Theorem

For a symmetric matrix P , $P \succ 0$ if and only if all the eigenvalues of P are positive.

Positive definite matrices

Theorem

For a symmetric matrix P , $P \succ 0$ if and only if all the eigenvalues of P are positive.

Proof.

Since P is symmetric, we have

$$\lambda_{\max}(P) = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (4)$$

$$\lambda_{\min}(P) = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (5)$$

Positive definite matrices

Theorem

For a symmetric matrix P , $P \succ 0$ if and only if all the eigenvalues of P are positive.

Proof.

Since P is symmetric, we have

$$\lambda_{\max}(P) = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (4)$$

$$\lambda_{\min}(P) = \min_{x \in \mathbb{R}^n, x \neq 0} \frac{x^T A x}{\|x\|_2^2} \quad (5)$$

which gives $x^T A x \in [\lambda_{\min} \|x\|_2^2, \lambda_{\max} \|x\|_2^2]$. Thus
 $x^T A x > 0, x \neq 0 \Leftrightarrow \lambda_{\min} > 0$. □

Relevant tools

Checking positive definiteness of a matrix.

We often use the following necessary and sufficient conditions to check positive (semi-)definiteness:

Relevant tools

Checking positive definiteness of a matrix.

We often use the following necessary and sufficient conditions to check positive (semi-)definiteness:

- ▶ $P \succ 0$ ($P \succeq 0$) \Leftrightarrow the leading principle minors defined below are positive (nonnegative)

Relevant tools

Checking positive definiteness of a matrix.

We often use the following necessary and sufficient conditions to check positive (semi-)definiteness:

- ▶ $P \succ 0$ ($P \succeq 0$) \Leftrightarrow the leading principle minors defined below are positive (nonnegative)

Definition

The leading principle minors of $P = \begin{bmatrix} p_{11} & p_{12} & p_{13} \\ p_{21} & p_{22} & p_{23} \\ p_{31} & p_{32} & p_{33} \end{bmatrix}$ are defined as

$$p_{11}, \det \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}, \det P.$$

Relevant tools


Checking positive definiteness of a matrix.

Example


None of the following matrices are positive definite:

$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 1 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$


Recap



Recap



Recap



Relevant tools

Definition (Positive Definite Functions)

A continuous time function $W : \mathbb{R}^n \rightarrow \mathbb{R}_+$, called to be PD, satisfying

- ▶ $W(x) > 0$ for all $x \neq 0$
- ▶ $W(0) = 0$
- ▶ $W(x) \rightarrow \infty$ as $|x| \rightarrow \infty$ uniformly in x

Relevant tools

Definition (Positive Definite Functions)

A continuous time function $W : \mathbb{R}^n \rightarrow \mathbb{R}_+$, called to be PD, satisfying

- ▶ $W(x) > 0$ for all $x \neq 0$
- ▶ $W(0) = 0$
- ▶ $W(x) \rightarrow \infty$ as $|x| \rightarrow \infty$ uniformly in x

In the 3D space, positive definite functions are “bowl-shaped”


Relevant tools

Definition (Positive Definite Functions)

A continuous time function $W : \mathbb{R}^n \rightarrow \mathbb{R}_+$, called to be PD, satisfying

- ▶ $W(x) > 0$ for all $x \neq 0$
- ▶ $W(0) = 0$
- ▶ $W(x) \rightarrow \infty$ as $|x| \rightarrow \infty$ uniformly in x

In the 3D space, positive definite functions are “bowl-shaped”, e.g., $W(x_1, x_2) = x_1^2 + x_2^2$.



Relevant tools

Definition (Locally Positive Definite Functions)

A continuous time function $W : \mathbb{R}^n \rightarrow \mathbb{R}_+$, called to be LPD, satisfying

- ▶ $W(x) > 0$ for all $x \neq 0$ and $|x| < r$
- ▶ $W(0) = 0$

Relevant tools

Definition (Locally Positive Definite Functions)

A continuous time function $W : \mathbb{R}^n \rightarrow \mathbb{R}_+$, called to be LPD, satisfying

- ▶ $W(x) > 0$ for all $x \neq 0$ and $|x| < r$
- ▶ $W(0) = 0$

In the 3D space, locally positive definite functions are “bowl-shaped” locally


Relevant tools

Definition (Locally Positive Definite Functions)

A continuous time function $W : \mathbb{R}^n \rightarrow \mathbb{R}_+$, called to be LPD, satisfying

- ▶ $W(x) > 0$ for all $x \neq 0$ and $|x| < r$
- ▶ $W(0) = 0$

In the 3D space, locally positive definite functions are “bowl-shaped” locally, e.g., $W(x_1, x_2) = x_1^2 + \sin^2 x_2$ for $x_1 \in \mathbb{R}$ and $|x_2| < \pi$



Relevant tools

Exercise

Let $x = [x_1, x_2, x_3]^T$. Check the positive definiteness of the following functions

1. $V(x) = x_1^4 + x_2^2 + x_3^4$

Exercise

Let $x = [x_1, x_2, x_3]^T$. Check the positive definiteness of the following functions

1. $V(x) = x_1^4 + x_2^2 + x_3^4$ (PD)
2. $V(x) = x_1^2 + x_2^2 + 3x_3^2 - x_3^4$

Relevant tools

Exercise

Let $x = [x_1, x_2, x_3]^T$. Check the positive definiteness of the following functions

1. $V(x) = x_1^4 + x_2^2 + x_3^4$ (PD)
2. $V(x) = x_1^2 + x_2^2 + 3x_3^2 - x_3^4$ (LPD for $|x_3| < \sqrt{3}$)

1. Definitions in Lyapunov stability analysis
2. Stability of LTI systems: method of eigenvalue/pole locations
3. Lyapunov's approach to stability
 - Relevant tools
 - Lyapunov stability theorems
 - Instability theorem
 - Discrete-time case
4. Recap

Lyapunov stability theorems

- recall the spring mass damper example in matrix form

$$\frac{d}{dt} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{b}{m} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Lyapunov stability theorems

- recall the spring mass damper example in matrix form

$$\frac{d}{dt} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{b}{m} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- energy function is PD:

$$\mathcal{E}(t) = \text{potential energy} + \text{kinetic energy} = \frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$$

Lyapunov stability theorems

- recall the spring mass damper example in matrix form

$$\frac{d}{dt} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{b}{m} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- energy function is PD:

$\mathcal{E}(t)$ = potential energy + kinetic energy = $\frac{1}{2}kx_1^2 + \frac{1}{2}mx_2^2$
and its derivative is NSD:

$$\dot{\mathcal{E}}(t) = \left[\frac{\partial \mathcal{E}}{\partial x_1}, \frac{\partial \mathcal{E}}{\partial x_2} \right] \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = k_1 x_1 \dot{x}_1 + m x_2 \dot{x}_2 \quad (6)$$

$$\begin{aligned} &= k_1 x_1 x_2 + m x_2 \left(-\frac{k}{m} x_1 - \frac{b}{m} x_2 \right) = \left[\frac{\partial \mathcal{E}}{\partial x_1}, \frac{\partial \mathcal{E}}{\partial x_2} \right] A x \quad (7) \\ &= -b x_2^2 \end{aligned}$$

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$ such that $\dot{V}(x, t) \leq 0$ for all $t \geq t_0$

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$ such that $\dot{V}(x, t) \leq 0$ for all $t \geq t_0$ and all x in a local region $x : |x| < r$ for some $r > 0$.

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$ such that $\dot{V}(x, t) \leq 0$ for all $t \geq t_0$ and all x in a local region $x : |x| < r$ for some $r > 0$.

- such a $V(x, t)$ is called a Lyapunov function

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$ such that $\dot{V}(x, t) \leq 0$ for all $t \geq t_0$ and all x in a local region $x : |x| < r$ for some $r > 0$.

- ▶ such a $V(x, t)$ is called a Lyapunov function
- ▶ i.e., $V(x)$ is PD and $\dot{V}(x)$ is negative semidefinite in a local region $|x| < r$

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$ such that $\dot{V}(x, t) \leq 0$ for all $t \geq t_0$ and all x in a local region $x : |x| < r$ for some $r > 0$.

- ▶ such a $V(x, t)$ is called a Lyapunov function
- ▶ i.e., $V(x)$ is PD and $\dot{V}(x)$ is negative semidefinite in a local region $|x| < r$

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is locally asymptotically stable if

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is stable in the sense of Lyapunov if there exists a locally positive definite function $V(x, t)$ such that $\dot{V}(x, t) \leq 0$ for all $t \geq t_0$ and all x in a local region $x : |x| < r$ for some $r > 0$.

- ▶ such a $V(x, t)$ is called a Lyapunov function
- ▶ i.e., $V(x)$ is PD and $\dot{V}(x)$ is negative semidefinite in a local region $|x| < r$

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is locally asymptotically stable if there exists a Lyapunov function $V(x)$ such that $\dot{V}(x)$ is locally negative definite.

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is globally asymptotically stable if

Theorem

The equilibrium point 0 of $\dot{x}(t) = f(x(t), t)$, $x(t_0) = x_0$ is globally asymptotically stable if there exists a Lyapunov function $V(x)$ such that $V(x)$ is positive definite and $\dot{V}(x)$ is negative definite.

Lyapunov stability concept for linear systems

- ▶ for linear system $\dot{x} = Ax$, a good Lyapunov candidate is the quadratic function $V(x) = x^T P x$ where $P = P^T$ and $P \succ 0$

Lyapunov stability concept for linear systems

- ▶ for linear system $\dot{x} = Ax$, a good Lyapunov candidate is the quadratic function $V(x) = x^T P x$ where $P = P^T$ and $P \succ 0$
- ▶ the derivative along the state trajectory is then

$$\begin{aligned}\dot{V}(x) &= \dot{x}^T P x + x^T P \dot{x} \\ &= (Ax)^T P x + x^T P A x \\ &= x^T (A^T P + P A) x\end{aligned}$$

Lyapunov stability concept for linear systems

- ▶ for linear system $\dot{x} = Ax$, a good Lyapunov candidate is the quadratic function $V(x) = x^T P x$ where $P = P^T$ and $P \succ 0$
- ▶ the derivative along the state trajectory is then

$$\begin{aligned}\dot{V}(x) &= \dot{x}^T P x + x^T P \dot{x} \\ &= (Ax)^T P x + x^T P A x \\ &= x^T (A^T P + P A) x\end{aligned}$$

- ▶ such a $V(x) = x^T P x$ is a Lyapunov function for $\dot{x} = Ax$ when $A^T P + P A \preceq 0$

Lyapunov stability concept for linear systems

- ▶ for linear system $\dot{x} = Ax$, a good Lyapunov candidate is the quadratic function $V(x) = x^T P x$ where $P = P^T$ and $P \succ 0$
- ▶ the derivative along the state trajectory is then

$$\begin{aligned}\dot{V}(x) &= \dot{x}^T P x + x^T P \dot{x} \\ &= (Ax)^T P x + x^T P Ax \\ &= x^T (A^T P + PA) x\end{aligned}$$

- ▶ such a $V(x) = x^T P x$ is a Lyapunov function for $\dot{x} = Ax$ when $A^T P + PA \preceq 0$
- ▶ and the origin is stable in the sense of Lyapunov

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$,

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Essense of the Lyapunov Eq.

Observations:

- ▶ $A^T P + PA$ is a linear operation on P

Essense of the Lyapunov Eq.

Observations:

► $A^T P + P A$ is a linear operation on P : e.g.,

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad Q = \begin{bmatrix} | & | \\ q_1 & q_2 \\ | & | \end{bmatrix}, \quad P = \begin{bmatrix} | & | \\ p_1 & p_2 \\ | & | \end{bmatrix}$$

$$A^T \begin{bmatrix} | & | \\ p_1 & p_2 \\ | & | \end{bmatrix} + \begin{bmatrix} | & | \\ p_1 & p_2 \\ | & | \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = - \begin{bmatrix} | & | \\ q_1 & q_2 \\ | & | \end{bmatrix}$$

Essense of the Lyapunov Eq.

Observations:

► $A^T P + PA$ is a linear operation on P : e.g.,

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad Q = \begin{bmatrix} | & | \\ q_1 & q_2 \\ | & | \end{bmatrix}, \quad P = \begin{bmatrix} | & | \\ p_1 & p_2 \\ | & | \end{bmatrix}$$

$$A^T \begin{bmatrix} | & | \\ p_1 & p_2 \\ | & | \end{bmatrix} + \begin{bmatrix} | & | \\ p_1 & p_2 \\ | & | \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = - \begin{bmatrix} | & | \\ q_1 & q_2 \\ | & | \end{bmatrix}$$

$$A^T p_1 + a_{11}p_1 + a_{21}p_2 = -q_1$$

$$A^T p_2 + a_{12}p_1 + a_{22}p_2 = -q_2$$

Essense of the Lyapunov Eq.

Observations: with now

$$A^T P + PA = Q \Leftrightarrow \begin{cases} A^T p_1 + a_{11}p_1 + a_{21}p_2 &= -q_1 \\ A^T p_2 + a_{12}p_1 + a_{22}p_2 &= -q_2 \end{cases}$$

Essense of the Lyapunov Eq.

Observations: with now

$$A^T P + PA = Q \Leftrightarrow \begin{cases} A^T p_1 + a_{11}p_1 + a_{21}p_2 = -q_1 \\ A^T p_2 + a_{12}p_1 + a_{22}p_2 = -q_2 \end{cases}$$

► can stack the columns of $A^T P + PA$ and Q to yield

$$\begin{aligned} \begin{bmatrix} A^T & 0 \\ 0 & A^T \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} + \begin{bmatrix} a_{11}/ & a_{21}/ \\ a_{12}/ & a_{22}/ \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} &= - \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} \\ \underbrace{\left\{ \begin{bmatrix} A^T & 0 \\ 0 & A^T \end{bmatrix} + \begin{bmatrix} a_{11}/ & a_{21}/ \\ a_{12}/ & a_{22}/ \end{bmatrix} \right\}}_{L_A} \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} &= - \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} \end{aligned}$$

The Lyapunov Eq.: Existence of solution

$$L_A(P) = A^T P + PA$$

- L_A is invertible if and only if $\lambda_i + \lambda_j \neq 0$ for all eigenvalues of A

The Lyapunov Eq.: Existence of solution

$$L_A(P) = A^T P + PA$$

- ▶ L_A is invertible if and only if $\lambda_i + \lambda_j \neq 0$ for all eigenvalues of A :
 - ▶ let $A^T u_i = \lambda_i u_i$ and $A^T u_j = \lambda_j u_j$

The Lyapunov Eq.: Existence of solution

$$L_A(P) = A^T P + PA$$

- ▶ L_A is invertible if and only if $\lambda_i + \lambda_j \neq 0$ for all eigenvalues of A :
 - ▶ let $A^T u_i = \lambda_i u_i$ and $A^T u_j = \lambda_j u_j$
 - ▶ $L_A(u_i u_j^T) = u_i u_j^T A + A^T u_i u_j^T = u_i (\lambda_j u_j)^T + \lambda_i u_i u_j^T = (\lambda_i + \lambda_j) u_i u_j^T$

The Lyapunov Eq.: Existence of solution

$$L_A(P) = A^T P + PA$$

- ▶ L_A is invertible if and only if $\lambda_i + \lambda_j \neq 0$ for all eigenvalues of A :
 - ▶ let $A^T u_i = \lambda_i u_i$ and $A^T u_j = \lambda_j u_j$
 - ▶ $L_A(u_i u_j^T) = u_i u_j^T A + A^T u_i u_j^T = u_i (\lambda_j u_j)^T + \lambda_i u_i u_j^T = (\lambda_i + \lambda_j) u_i u_j^T$
 - ▶ so $\lambda_i + \lambda_j$ is an eigenvalue of the operator $L_A(\cdot)$

The Lyapunov Eq.: Existence of solution

$$L_A(P) = A^T P + PA$$

- ▶ L_A is invertible if and only if $\lambda_i + \lambda_j \neq 0$ for all eigenvalues of A :
 - ▶ let $A^T u_i = \lambda_i u_i$ and $A^T u_j = \lambda_j u_j$
 - ▶ $L_A(u_i u_j^T) = u_i u_j^T A + A^T u_i u_j^T = u_i (\lambda_j u_j)^T + \lambda_i u_i u_j^T = (\lambda_i + \lambda_j) u_i u_j^T$
 - ▶ so $\lambda_i + \lambda_j$ is an eigenvalue of the operator $L_A(\cdot)$
 - ▶ if $\lambda_i + \lambda_j \neq 0$, the operator is invertible

The Lyapunov operator: eigenvalues

$$L_A = \begin{bmatrix} A^T & 0 \\ 0 & A^T \end{bmatrix} + \begin{bmatrix} a_{11}I & a_{21}I \\ a_{12}I & a_{22}I \end{bmatrix}$$

- can simply write $L_A = \underbrace{I \otimes A^T + A^T \otimes I}_{\text{mirror symmetric}}$ using the Kronecker

product notation $B \otimes C = \begin{bmatrix} b_{11}C & b_{12}C & \dots & b_{1n}C \\ b_{21}C & b_{22}C & \dots & b_{2n}C \\ \vdots & \vdots & \dots & \vdots \\ b_{m1}C & b_{m2}C & \dots & b_{mn}C \end{bmatrix}$

The Lyapunov operator: eigenvalues

$$L_A = \begin{bmatrix} A^T & 0 \\ 0 & A^T \end{bmatrix} + \begin{bmatrix} a_{11}I & a_{21}I \\ a_{12}I & a_{22}I \end{bmatrix}$$

► e.g., $A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}$

$$\begin{aligned} L_A &= I \otimes A^T + A^T \otimes I = \begin{bmatrix} A^T + a_{11}I & a_{21}I \\ a_{12}I & A^T + a_{22}I \end{bmatrix} \\ &= \left[\begin{array}{cc|cc} -1 & -1 & -1 & 0 \\ 1 & 0 & 0 & -1 \\ \hline 1 & 0 & -1 & -1 \\ 0 & 1 & 1 & 0 \end{array} \right] = \left[\begin{array}{cc|cc} -2 & -1 & -1 & 0 \\ 1 & -1 & 0 & -1 \\ \hline 1 & 0 & -1 & -1 \\ 0 & 1 & 1 & 0 \end{array} \right] \end{aligned}$$

Example: $A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}$, $\lambda_{1,2} = -0.5 \pm i\sqrt{3}/2$

$$L_A = I \otimes A^T + A^T \otimes I = \left[\begin{array}{cc|cc} -2 & -1 & -1 & 0 \\ 1 & -1 & 0 & -1 \\ \hline 1 & 0 & -1 & -1 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

Example: $A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}$, $\lambda_{1,2} = -0.5 \pm i\sqrt{3}/2$

$$L_A = I \otimes A^T + A^T \otimes I = \left[\begin{array}{cc|cc} -2 & -1 & -1 & 0 \\ 1 & -1 & 0 & -1 \\ \hline 1 & 0 & -1 & -1 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

The eigenvalues of L_A are $-1, -1, -1 - \sqrt{3}, -1 + \sqrt{3}$, which are precisely $\lambda_1 + \lambda_1, \lambda_1 + \lambda_2, \lambda_2 + \lambda_1, \lambda_2 + \lambda_2$.

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$,

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"}\Rightarrow\text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x}$$

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"}\Rightarrow\text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha}$$

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"}\Rightarrow\text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha} \implies V(t) \leq e^{-\alpha t} V(0).$$

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"} \Rightarrow \text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha} \implies V(t) \leq e^{-\alpha t} V(0). \quad Q \succ 0 \text{ and}$$

$$P \succ 0 \implies$$

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\begin{aligned} \text{"}\Rightarrow\text{"}: \quad \frac{\dot{V}}{V} &= -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha} \implies V(t) \leq e^{-\alpha t} V(0). \quad Q \succ 0 \text{ and} \\ P \succ 0 &\implies (\lambda_Q)_{\min} > 0 \text{ and } (\lambda_P)_{\max} > 0. \end{aligned}$$

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"}\Rightarrow\text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha} \Rightarrow V(t) \leq e^{-\alpha t} V(0). \quad Q \succ 0 \text{ and}$$

$P \succ 0 \Rightarrow (\lambda_Q)_{\min} > 0$ and $(\lambda_P)_{\max} > 0$. Thus $\alpha > 0$; $V(t)$ decays exponentially to zero.

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"}\Rightarrow\text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha} \Rightarrow V(t) \leq e^{-\alpha t} V(0). \quad Q \succ 0 \text{ and}$$

$P \succ 0 \Rightarrow (\lambda_Q)_{\min} > 0$ and $(\lambda_P)_{\max} > 0$. Thus $\alpha > 0$; $V(t)$ decays exponentially to zero. $V(x) \succ 0 \Rightarrow V(x) = 0$ only at $x = 0$.

Theorem (Lyapunov stability theorem for linear systems)

For $\dot{x} = Ax$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if for any symmetric positive definite matrix $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

Proof.

$$\text{"}\Rightarrow\text{"}: \frac{\dot{V}}{V} = -\frac{x^T Q x}{x^T P x} \leq -\underbrace{\frac{(\lambda_Q)_{\min}}{(\lambda_P)_{\max}}}_{\triangleq \alpha} \implies V(t) \leq e^{-\alpha t} V(0). \quad Q \succ 0 \text{ and}$$

$P \succ 0 \implies (\lambda_Q)_{\min} > 0$ and $(\lambda_P)_{\max} > 0$. Thus $\alpha > 0$; $V(t)$ decays exponentially to zero. $V(x) \succ 0 \implies V(x) = 0$ only at $x = 0$.

Therefore, $x \rightarrow 0$ as $t \rightarrow \infty$, regardless of the initial condition. \square

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable,

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable, then all eigenvalues of A have negative real parts.

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable, then all eigenvalues of A have negative real parts. For any Q , the Lyapunov equation has a unique solution P .

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable, then all eigenvalues of A have negative real parts. For any Q , the Lyapunov equation has a unique solution P . Note $x(t) = e^{At}x_0 \rightarrow 0$ as $t \rightarrow \infty$. We have

$$\begin{aligned} \cancel{x^T(\infty) P x(\infty)} - x^T(0) P x(0) &= \int_0^\infty \frac{d}{dt} x^T(t) P x(t) dt = \int_0^\infty x^T(t) (A^T P + P A) x(t) dt \\ &\Rightarrow x^T(0) P x(0) = \int_0^\infty x^T(t) Q x(t) dt = \int_0^\infty x^T(0) e^{A^T t} Q e^{At} x(0) dt \end{aligned}$$

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable, then all eigenvalues of A have negative real parts. For any Q , the Lyapunov equation has a unique solution P . Note $x(t) = e^{At}x_0 \rightarrow 0$ as $t \rightarrow \infty$. We have

$$\begin{aligned} \cancel{x^T(\infty)Px(\infty)} - x^T(0)Px(0) &= \int_0^\infty \frac{d}{dt} x^T(t)Px(t) dt = \int_0^\infty x^T(t) (A^T P + PA) x(t) dt \\ &\Rightarrow x^T(0)Px(0) = \int_0^\infty x^T(t) Q x(t) dt = \int_0^\infty x^T(0) e^{A^T t} Q e^{At} x(0) dt \end{aligned}$$

If $Q \succ 0$, there exists a nonsingular N matrix: $Q = N^T N$. Thus

$$x^T(0)Px(0) = \int_0^\infty \|Ne^{At}x(0)\|^2 dt \geq 0$$

$$x^T(0)Px(0) = 0 \text{ only if } x_0 = 0$$

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable, then all eigenvalues of A have negative real parts. For any Q , the Lyapunov equation has a unique solution P . Note $x(t) = e^{At}x_0 \rightarrow 0$ as $t \rightarrow \infty$. We have

$$\begin{aligned} \cancel{x^T(\infty)Px(\infty)} - x^T(0)Px(0) &= \int_0^\infty \frac{d}{dt} x^T(t)Px(t) dt = \int_0^\infty x^T(t) (A^T P + PA) x(t) dt \\ &\Rightarrow x^T(0)Px(0) = \int_0^\infty x^T(t) Q x(t) dt = \int_0^\infty x^T(0) e^{A^T t} Q e^{At} x(0) dt \end{aligned}$$

If $Q \succ 0$, there exists a nonsingular N matrix: $Q = N^T N$. Thus

$$x^T(0)Px(0) = \int_0^\infty \|Ne^{At}x(0)\|^2 dt \geq 0$$

$$x^T(0)Px(0) = 0 \text{ only if } x_0 = 0$$

Thus $P \succ 0$.

Proof.

“ \Leftarrow ”: if 0 of $\dot{x} = Ax$ is asymptotically stable, then all eigenvalues of A have negative real parts. For any Q , the Lyapunov equation has a unique solution P . Note $x(t) = e^{At}x_0 \rightarrow 0$ as $t \rightarrow \infty$. We have

$$\begin{aligned} \cancel{x^T(\infty)Px(\infty)} - x^T(0)Px(0) &= \int_0^\infty \frac{d}{dt} x^T(t)Px(t) dt = \int_0^\infty x^T(t) (A^T P + PA) x(t) dt \\ &\Rightarrow x^T(0)Px(0) = \int_0^\infty x^T(t) Q x(t) dt = \int_0^\infty x^T(0) e^{A^T t} Q e^{At} x(0) dt \end{aligned}$$

If $Q \succ 0$, there exists a nonsingular N matrix: $Q = N^T N$. Thus

$$x^T(0)Px(0) = \int_0^\infty \|Ne^{At}x(0)\|^2 dt \geq 0$$

$$x^T(0)Px(0) = 0 \text{ only if } x_0 = 0$$

Thus $P \succ 0$. Furthermore

$$P = \int_0^\infty e^{A^T t} Q e^{At} dt$$



Procedures of Lyapunov's direct method

1. Given A , select an arbitrary positive-definite symmetric matrix Q (e.g., I).
2. Find the solution matrix P to the Lyapunov equation
$$A^T P + PA = -Q.$$

Procedures of Lyapunov's direct method

1. Given A , select an arbitrary positive-definite symmetric matrix Q (e.g., I).
2. Find the solution matrix P to the Lyapunov equation
$$A^T P + PA = -Q.$$
3. If a solution P cannot be found, the origin is not asymptotically stable.

Procedures of Lyapunov's direct method

1. Given A , select an arbitrary positive-definite symmetric matrix Q (e.g., I).
2. Find the solution matrix P to the Lyapunov equation
$$A^T P + PA = -Q.$$
3. If a solution P cannot be found, the origin is not asymptotically stable.
4. If a solution is found:
 - ▶ if P is positive-definite, then A is Hurwitz stable and the origin is asymptotically stable;

Procedures of Lyapunov's direct method

1. Given A , select an arbitrary positive-definite symmetric matrix Q (e.g., I).
2. Find the solution matrix P to the Lyapunov equation
$$A^T P + PA = -Q.$$
3. If a solution P cannot be found, the origin is not asymptotically stable.
4. If a solution is found:
 - ▶ if P is positive-definite, then A is Hurwitz stable and the origin is asymptotically stable;
 - ▶ if P is not positive-definite, then A has at least one eigenvalue with a positive real part and the origin is an unstable equilibrium.

Lyapunov stability theorems

Example

$\dot{x} = Ax$, $A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}$. The Lyapunov equation is

$$\begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}^T \underbrace{\begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix}}_P + \begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix} \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix} = - \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}}_Q$$

Lyapunov stability theorems

Example

$\dot{x} = Ax$, $A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}$. The Lyapunov equation is

$$\begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}^T \underbrace{\begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix}}_P + \underbrace{\begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix}}_P \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix} = - \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}}_Q$$

We need

$$\begin{cases} -2p_{11} - 2p_{12} = -1 \\ -p_{12} - p_{22} + p_{11} = 0 \\ 2p_{12} = -1 \end{cases} \Rightarrow \begin{cases} p_{11} = 1 \\ p_{22} = 3/2 \\ p_{12} = -1/2 \end{cases}$$

Lyapunov stability theorems

Example

$\dot{x} = Ax$, $A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}$. The Lyapunov equation is


$$\begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}^T \underbrace{\begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix}}_P + \underbrace{\begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix}}_P \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix} = - \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}}_Q$$

We need

$$\begin{cases} -2p_{11} - 2p_{12} = -1 \\ -p_{12} - p_{22} + p_{11} = 0 \\ 2p_{12} = -1 \end{cases} \Rightarrow \begin{cases} p_{11} = 1 \\ p_{22} = 3/2 \\ p_{12} = -1/2 \end{cases}$$


Leading principle minors: $p_{11} > 0$, $p_{11}p_{22} - p_{12}^2 > 0$
 $\Rightarrow P \succ 0 \Rightarrow$ asymptotically stable

Lyapunov analysis with Matlab

$$\dot{x} = Ax, A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}.$$


Lyapunov analysis with Python

$$\dot{x} = Ax, A = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}.$$



It suffices to select $Q = I$

For linear systems we can let $Q = I$ and check whether the resulting P is positive definite. If it is, then we can assert the asymptotic stability:

It suffices to select $Q = I$

For linear systems we can let $Q = I$ and check whether the resulting P is positive definite. If it is, then we can assert the asymptotic stability:

- take any $Q \succ 0$. there exists $Q = N^T N$, where N is invertible, yielding

$$\begin{aligned} A^T P + P A &= -I \\ \Downarrow \\ \underbrace{N^T A^T N^{-T}}_{\tilde{A}^T} \underbrace{N^T P N}_{\tilde{P}} + \underbrace{N^T P N}_{\tilde{P}} \underbrace{N^{-1} A N}_{\tilde{A}} &= -N^T N \end{aligned}$$

It suffices to select $Q = I$

For linear systems we can let $Q = I$ and check whether the resulting P is positive definite. If it is, then we can assert the asymptotic stability:

- take any $Q \succ 0$. there exists $Q = N^T N$, where N is invertible, yielding

$$\begin{aligned} A^T P + P A &= -I \\ \Updownarrow \\ \underbrace{N^T A^T N^{-T}}_{\tilde{A}^T} \underbrace{N^T P N}_{\tilde{P}} + \underbrace{N^T P N}_{\tilde{P}} \underbrace{N^{-1} A N}_{\tilde{A}} &= -N^T N \end{aligned}$$

- $\tilde{A} = N^{-1} A N$ and A are similar matrices and have the same eigenvalues.

It suffices to select $Q = I$

For linear systems we can let $Q = I$ and check whether the resulting P is positive definite. If it is, then we can assert the asymptotic stability:

- take any $Q \succ 0$. there exists $Q = N^T N$, where N is invertible, yielding

$$\begin{aligned} A^T P + P A &= -I \\ \Updownarrow \\ \underbrace{N^T A^T N^{-T}}_{\tilde{A}^T} \underbrace{N^T P N}_{\tilde{P}} + \underbrace{N^T P N}_{\tilde{P}} \underbrace{N^{-1} A N}_{\tilde{A}} &= -N^T N \end{aligned}$$

- $\tilde{A} = N^{-1} A N$ and A are similar matrices and have the same eigenvalues.
- $\tilde{P} = N^T P N$ and P have the same definiteness. If we can find a positive definite solution P then the \tilde{P} will also be positive definite. Vise versa.

Instability theorem

- ▶ for nonlinear systems, Lyapunov function can be nontrivial to find

Instability theorem

- ▶ for nonlinear systems, Lyapunov function can be nontrivial to find
- ▶ failure to find a Lyapunov function does not imply instability

Instability theorem

- ▶ for nonlinear systems, Lyapunov function can be nontrivial to find
- ▶ failure to find a Lyapunov function does not imply instability

Theorem

The equilibrium state 0 of $\dot{x} = f(x)$ is unstable if there exists a function $W(x)$ such that

- ▶ $\dot{W}(x)$ is PD locally: $\dot{W}(x) > 0 \quad \forall |x| < r$ for some r and $\dot{W}(0) = 0$
- ▶ $W(0) = 0$
- ▶ *there exist states x arbitrarily close to the origin such that $W(x) > 0$*

Discrete-time case: key concept of Lyapunov

For the discrete-time system

$$x(k+1) = Ax(k)$$

we consider a quadratic Lyapunov function candidate

$$V(x) = x^T P x, \quad P = P^T \succ 0$$

Discrete-time case: key concept of Lyapunov

For the discrete-time system

$$x(k+1) = Ax(k)$$

we consider a quadratic Lyapunov function candidate

$$V(x) = x^T P x, \quad P = P^T \succ 0$$

and compute $\Delta V(x)$ along the trajectory of the state

$$V(x(k+1)) - V(x(k)) = x^T(k) \underbrace{(A^T P A - P)}_{\triangleq -Q} x(k)$$

Discrete-time case: key concept of Lyapunov

For the discrete-time system

$$x(k+1) = Ax(k)$$

we consider a quadratic Lyapunov function candidate

$$V(x) = x^T P x, \quad P = P^T \succ 0$$

and compute $\Delta V(x)$ along the trajectory of the state

$$V(x(k+1)) - V(x(k)) = x^T(k) \underbrace{(A^T P A - P)}_{\triangleq -Q} x(k)$$

Asymptotic stability desires $\Delta V(x)$ to be negative.

DT Lyapunov stability theorem for linear systems

Theorem

For system $x(k+1) = Ax(k)$ with $A \in \mathbb{R}^{n \times n}$, the origin is asymptotically stable if and only if $\exists Q \succ 0$, such that the discrete-time Lyapunov equation

$$A^T P A - P = -Q$$

has a unique positive definite solution $P \succ 0$, $P^T = P$.

The DT Lyapunov Eq.

$$\boxed{A^T P A - P = -Q}$$

- Solution to the DT Lyapunov equation, when asymptotic stability holds (A is Schur stable), comes from:

$$\begin{aligned} \cancel{V(x(\infty))} - V(x(0)) &= \sum_{k=0}^{\infty} x^T(k) [A^T P A - P] x(k) \\ &= - \sum_{k=0}^{\infty} x^T(0) (A^T)^k Q A^k x(0) \\ \Rightarrow P &= \sum_{k=0}^{\infty} (A^T)^k Q A^k \end{aligned}$$

The DT Lyapunov Eq.

$$\boxed{A^T P A - P = -Q}$$

- Solution to the DT Lyapunov equation, when asymptotic stability holds (A is Schur stable), comes from:

$$\begin{aligned} \cancel{V(x(\infty))} - V(x(0)) &= \sum_{k=0}^{\infty} x^T(k) [A^T P A - P] x(k) \\ &= - \sum_{k=0}^{\infty} x^T(0) (A^T)^k Q A^k x(0) \\ \Rightarrow P &= \sum_{k=0}^{\infty} (A^T)^k Q A^k \end{aligned}$$

- can show that the DT Lyapunov operator $L_A = A^T P A - P$ is invertible if and only if $\forall i, j \ (\lambda_A)_i (\lambda_A)_j \neq 1$

DT Lyapunov analysis with MATLAB

Example

$$x(k+1) = Ax(k), \quad A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0.275 & -0.225 & -0.1 \end{bmatrix}$$

DT Lyapunov analysis with Python

Example

$$x(k+1) = Ax(k), \quad A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0.275 & -0.225 & -0.1 \end{bmatrix}$$

Recap

- ▶ Internal stability
 - ▶ Stability in the sense of Lyapunov: ϵ, δ conditions
 - ▶ Asymptotic stability
- ▶ Stability analysis of linear time invariant systems ($\dot{x} = Ax$ or $x(k+1) = Ax(k)$)
 - ▶ Based on the eigenvalues of A
 - ▶ Time response modes
 - ▶ Repeated eigenvalues on the imaginary axis
 - ▶ Routh's criterion
 - ▶ No need to solve the characteristic equation
 - ▶ Discrete time case: bilinear transform ($z = \frac{1+s}{1-s}$)

Recap

► Lyapunov equations

Theorem: All eigenvalues of A have negative real parts iff for any given $Q \succ 0$, the Lyapunov equation

$$A^T P + PA = -Q$$

has a unique solution P and $P \succ 0$.

Given Q , the Lyapunov equation $A^T P + PA = -Q$ has a unique solution when $\lambda_{A,i} + \lambda_{A,j} \neq 0$ for all i and j .

Theorem: All eigenvalues of A are inside the unit circle iff for any given $Q \succ 0$, the Lyapunov equation

$$A^T P A - P = -Q$$

has a unique solution P and $P \succ 0$.

Given Q , the Lyapunov equation $A^T P A - P = -Q$ has a unique solution when $\lambda_{A,i} \lambda_{A,j} \neq 1$ for all i and j .

Recap

- ▶ P is positive definite if and only if any one of the following conditions holds:
 1. All the eigenvalues of P are positive.
 2. All the leading principle minors of P are positive.
 3. There exists a nonsingular matrix N such that $P = N^T N$.