

# MS 120 In-class Problems

September 24, 2024

# Table of Contents

## Ch 1

Ch R1.2

Ch R1.3

Ch R1.6

Ch 1.5

Ch R1.6 again

Ch 3.3

Ch 3.3 Numerical

## Chapter R Section R1.2

**1.2.001** Use the values in the following table.

$x$	-6	-1	0	3	4.2	9	12	14	15	22
$y$	0	0	1	5	9	12	38	22	22	70

- Explain why the table defines  $y$  as a function of  $x$ .
  - ☐ For each value of  $y$  there are multiple values for  $x$ .
  - ☐ For each value of  $y$  there is only one  $x$ .
  - ☐ For each value of  $x$  there are multiple values for  $y$ .
  - ☐ For each value of  $x$  there is only one  $y$ .
  - ☐ For some values of  $y$  there are multiple values for  $x$ .
- State the domain and range of this function.

domain:


range:

- If the table expresses  $y = f(x)$ , find  $f(0)$  and  $f(12)$ . (If the table does not express  $y = f(x)$ , enter DNE.)

$$f(0) =$$


$$f(12) =$$

**1.2.005a** Determine whether the graph represents  $y$  as a function of  $x$ . Explain your answer.



- ▶ Yes, the vertical line test shows that the graph represents  $y$  as a function of  $x$ .
- ▶ Yes, the horizontal line test shows that the graph represents  $y$  as a function of  $x$ .
- ▶ No, the vertical line test shows that the graph does not represent  $y$  as a function of  $x$ .
- ▶ No, the horizontal line test shows that the graph does not represent  $y$  as a function of  $x$ .
- ▶ There is no way to determine this using the graph.

**1.2.005b** Determine whether the graph represents  $y$  as a function of  $x$ . Explain your answer.



- ▶ Yes, the vertical line test shows that the graph represents  $y$  as a function of  $x$ .
- ▶ Yes, the horizontal line test shows that the graph represents  $y$  as a function of  $x$ .
- ▶ No, the vertical line test shows that the graph does not represent  $y$  as a function of  $x$ .
- ▶ No, the horizontal line test shows that the graph does not represent  $y$  as a function of  $x$ .
- ▶ There is no way to determine this using the graph.

**1.2.009** If  $R(x) = 8x - 11$ , find the following. (Give exact answers. Do not round.)

1.  $R(0) =$


2.  $R(2) =$

3.  $R(-3) =$

4.  $R(1.6) =$

**1.2.029** A function and its graph are given. Find the domain. (Enter your answer using interval notation.)

$$f(x) = \frac{\sqrt{x-3}}{x-8}$$





## Chapter R Section R1.3

**1.3.001** Find the intercepts and graph.

$$5x + 8y = 40$$

**1.3.005** Find the slope  $m$  of the line passing through the given pair of points. (If an answer is undefined, enter UNDEFINED.)

$(20, 21)$  and  $(14, -3)$


**1.3.011** If a line is horizontal, then its slope is \_\_\_\_\_.

**1.3.013** What is the rate of change of the function whose graph is a line passing through  $(3, 4)$  and  $(-1, 4)$ ?

**1.3.015a** For the given graph, determine whether the line has a slope that is positive, negative, 0, or undefined.



**1.3.015b** For the given graph, determine whether the line has a slope that is positive, negative, 0, or undefined.



**1.3.017** Find the slope  $m$  and  $y$ -intercept  $b$ . (Give exact answers. Do not round. If an answer is undefined, enter UNDEFINED. If an answer does not exist, enter DNE.)

$$y = \frac{7}{3}x - \frac{1}{2}.$$



**1.3.023** Find the slope  $m$  and  $y$ -intercept  $b$ . (Give exact answers. Do not round. If an answer is undefined, enter UNDEFINED. If an answer does not exist, enter DNE.)

$$2x + 7y = 14.$$

**1.3.025** Write the slope-intercept form of the equation of the line that has the given slope and  $y$ -intercept.

Slope  $\frac{1}{3}$  and  $y$ -intercept  $-3$

**1.3.033** Write the equation of the line that passes through the given point and has the given slope.

$(-2, 2)$  with undefined slope

**1.3.035** Write the equation of the line described.

Through  $(4, 5)$  and  $(-1, -5)$

**1.3.041** Determine whether the following pair of equations represents parallel lines, perpendicular lines, or neither of these.

$$3x + 8y = 24; \quad 8x - 3y = 24$$


**1.3.045** Write the equation of the line passing through  $(-2, -1)$  that is parallel to  $3x + 5y = 11$ .



In engineering and science, dimensional analysis is the analysis of the relationships between different physical quantities by identifying their base quantities (such as length, mass, time, and electric current) and units of measurement (such as metres and grams) and tracking these dimensions as calculations or comparisons are performed.

The term dimensional analysis is also used to refer to conversion of units from one dimensional unit to another, which can be used to evaluate scientific formulae.<sup>1</sup>

---

<sup>1</sup>[https://en.wikipedia.org/wiki/Dimensional\\_analysis](https://en.wikipedia.org/wiki/Dimensional_analysis) 



**1.6.005a** A linear cost function is  $C(x) = 3x + 750$ . (Assume  $C$  is measured in dollars.)

1. What are the slope and the  $C$ -intercept?
2. What is the marginal cost  $C'$  ( $\overline{MC}$ )?
3. What does the marginal cost mean?
  - a) Each additional unit produced costs this much (in dollars).
  - b) If production is increased by this many units, the cost decreases by \$1.
  - c) If production is increased by this many units, the cost increases by \$1.
  - d) Each additional unit produced reduces the cost by this much (in dollars).
4. What are the fixed costs?
5. How are your answers to parts (1), (2), and (3) related?
  - a)  $\frac{C\text{-intercept}}{\text{slope}} = \text{marginal cost}$
  - b)  $\text{slope} = \text{fixed costs, and } C\text{-intercept} = \text{marginal cost}$
  - c)  $\text{slope} = \text{marginal cost, and } C\text{-intercept} = \text{fixed costs}$
  - d)  $\frac{\text{slope}}{C\text{-intercept}} = \text{marginal cost}$
6. What is the cost of producing one more item if 50 are currently being produced?  
What is the cost of producing one more item if 100 are currently being produced?

**1.6.007** A linear revenue function is  $R = 26x$ . (Assume  $R$  is measured in dollars.)

1. What is the slope  $m$ ?
2. What is the marginal revenue  $R'$ ?  
What does the marginal revenue mean?

- |  |   |
|--|---|
| a) Each additional unit sold decreases the revenue by this many dollars. | b) If the number of units sold is increased by this amount, the revenue increases by \$1. |
| c) Each additional unit sold yields this many dollars in revenue.        | d) If the number of units sold is increased by this amount, the revenue decreases by \$1. |

3. What is the revenue received from selling one more item if 50 are currently being sold?  
What is the revenue received from selling one more item if 100 are being sold?

**1.6.001** Suppose a calculator manufacturer has the total cost function  $C(x) = 22x + 6600$  and the total revenue function  $R(x) = 56x$ .

1. What is the equation of the profit function  $P(x)$  for the calculator?

2. What is the profit on 2800 units?

**1.6.003** Suppose a ceiling fan manufacturer has the total cost function  $C(x) = 34x + 560$  and the total revenue function  $R(x) = 48x$ .

1. What is the equation of the profit function  $P(x)$  for this commodity?

$$P(x) =$$

2. What is the profit on 20 units?

Interpret your result.

- ▶ The total costs are less than the revenue.
  - ▶ The total costs are more than the revenue.
  - ▶ The total costs are exactly the same as the revenue.
3. How many fans must be sold to avoid losing money?

**1.6.009** Let  $C(x) = 3x + 750$  and  $R(x) = 21x$ .

1. Write the profit function  $P(x)$ .
2. What is the slope  $m$  of the profit function?
3. What is the marginal profit  $P'$ ?
4. Interpret the marginal profit.
  - a) Each additional unit sold decreases the profit by this much.
  - b) Each additional unit sold increases the profit by this much.
  - c) This is the smallest number of units that can be sold in order to make a profit.
  - d) The profit is maximized when this many units are sold.

**1.6.013 1-3** Extreme Protection, Inc. manufactures helmets for skiing and snowboarding. The fixed costs for one model of helmet are \$4700 per month. Materials and labor for each helmet of this model are \$50, and the company sells this helmet to dealers for \$70 each. (Let  $x$  represent the number of helmets sold. Let  $C$ ,  $R$ , and  $P$  be measured in dollars.)

1. For this helmet, write the function for monthly total costs  $C(x)$ .

$$C(x) =$$

2. Write the function for total revenue  $R(x)$ .

$$R(x) =$$

3. Write the function for profit  $P(x)$ .

$$P(x) =$$

**1.6.013 4** Extreme Protection, Inc. manufactures helmets for skiing and snowboarding. The fixed costs for one model of helmet are \$4700 per month. Materials and labor for each helmet of this model are \$50, and the company sells this helmet to dealers for \$70 each. (Let  $x$  represent the number of helmets sold. Let  $C$ ,  $R$ , and  $P$  be measured in dollars.)

4. Find  $C(200)$ .

$$C(200) =$$

Interpret  $C(200)$ .

- ▶ For each \$1 increase in cost this many more helmets can be produced.
- ▶ This is the cost (in dollars) of producing 200 helmets.
- ▶ For every additional helmet produced the cost increases by this much.
- ▶ When this many helmets are produced the cost is \$200.

**1.6.013 4** Extreme Protection, Inc. manufactures helmets for skiing and snowboarding. The fixed costs for one model of helmet are \$4700 per month. Materials and labor for each helmet of this model are \$50, and the company sells this helmet to dealers for \$70 each. (Let  $x$  represent the number of helmets sold. Let  $C$ ,  $R$ , and  $P$  be measured in dollars.)

4. Find  $R(200)$ .

$$R(200) =$$

Interpret  $R(200)$ .

- ▶ When this many helmets are produced the revenue generated is \$200.
- ▶ For each \$1 increase in revenue this many more helmets can be produced.
- ▶ For every additional helmet produced the revenue generated increases by this much.
- ▶ This is the revenue (in dollars) generated from the sale of 200 helmets.



**1.6.013 4** Extreme Protection, Inc. manufactures helmets for skiing and snowboarding. The fixed costs for one model of helmet are \$4700 per month. Materials and labor for each helmet of this model are \$50, and the company sells this helmet to dealers for \$70 each. (Let  $x$  represent the number of helmets sold. Let  $C$ ,  $R$ , and  $P$  be measured in dollars.)

4. Find  $P(200)$ .

$$P(200) =$$

Interpret  $P(200)$ .

- ▶ This is the profit (in dollars) when 200 helmets are sold, but since it is negative it means that the company loses money when 200 helmets are sold.
- ▶ For each additional helmet sold the profit (in dollars) increases by this much, but since it is positive it means that the company is producing too many helmets.
- ▶ For each additional helmet sold the profit (in dollars) increases by this much, but since it is negative it means that the company needs to decrease the number of helmets sold in order to make a profit.
- ▶ This is the profit (in dollars) when 200 helmets are sold, and since it is positive it means that the company makes money when 200 helmets are sold.

**1.6.013 5,6** Extreme Protection, Inc. manufactures helmets for skiing and snowboarding. The fixed costs for one model of helmet are \$4700 per month. Materials and labor for each helmet of this model are \$50, and the company sells this helmet to dealers for \$70 each. (Let  $x$  represent the number of helmets sold. Let  $C$ ,  $R$ , and  $P$  be measured in dollars.)

5. Same as the last part but with \$300 instead of \$200.

6. Find the marginal profit  $P'$ .

$$P' =$$

Write a sentence that explains its meaning.

- ▶ When revenue is increased by this much the profit is increased by \$1.
- ▶ For each \$1 increase in profit this many more helmets can be produced.
- ▶ When costs are decreased by this much the profit is increased by \$1.
- ▶ Each additional helmet sold increases the profit by this many dollars.

**1.6.015** The figure shows graphs of the total cost function and the total revenue function for a commodity. (Assume cost and revenue are measured in dollars.)



1. Label each function correctly.  
Choose from *total revenue function*, *total cost function*

a) function  
A

b) function  
B

2. Determine the fixed costs.
3. Locate the break-even point.  
 $(x, y) =$   
Determine the number of units sold to break even.
4. Estimate the marginal cost  $C'$  and marginal revenue  $R'$ .

## Definition 1 (Market equilibrium)


*occurs when the quantity of a commodity demanded is equal to the quantity supplied.*

## Law 1 (Law of demand)



*states that the quantity demanded will increase as price decreases and that the quantity demanded will decrease as price increases.*

## Law 2 (Law of supply)

*states that the quantity supplied for sale will increase as the price of a product increases.*



**1.6.031** The graphs of the demand function and supply function for a certain product, are given below. Use these graphs to answer the questions.



1. How many units  $q$  are demanded when the price  $p$  is \$50?
2. How many units  $q$  are supplied when the price  $p$  is \$50?
3. Will there be a market surplus (more supplied) or shortage (more demanded) when  $p = \$50$ ?

**1.6.033** If the demand for a pair of shoes is given by  $2p + 5q = 200$  and the supply function for it is  $p - 2q = 10$ , compare the quantity demanded and the quantity supplied when the price is \$90.

quantity demanded \_\_\_\_\_ pairs of shoes

quantity supplied \_\_\_\_\_ pairs of shoes

Will there be a surplus or shortfall at this price?





**Ex 1** Graphical method

$$\begin{cases} x + y = 2 \\ x - 2y = -1 \end{cases}$$

**Ex 2** Graphical method

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

**Ex 3** Graphical method

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

**Ex 4** Graphical method

$$\begin{cases} y = x^2 + 1 \\ y = x + 1 \end{cases}$$

**Ex 5** Graphical method

$$\begin{cases} x + y = 2 \\ x - 2y = -1 \end{cases}$$

**Ex 6** Substitution method

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

**Ex 7** Substitution method

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

**Ex 8** Substitution method

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$



**Ex 9** Substitution method

$$\begin{cases} x - 3y + z = 0 \\ y - z = 3 \\ z = -2 \end{cases}$$

**Ex 10** Elimination method

$$\begin{cases} x + y = 1 \\ x - y = 2 \end{cases}$$

**Ex 11** Elimination method

$$\begin{cases} x + y = 2 \\ x - 2y = -1 \end{cases}$$

**Ex 12** Elimination method

$$\begin{cases} 2x + 3y = 9 \\ x - y = 2 \end{cases}$$

**Ex 13** Elimination method

$$\begin{cases} 8x - 3y = -11 \\ 5x - 2y = -6 \end{cases}$$

**Ex 14** Elimination method

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

**Ex 15** Elimination method

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

**Ex 16** Elimination method

$$\begin{cases} x - 3y + z = 0 \\ y - z = 3 \\ z = -2 \end{cases}$$



**Ex 17** Elimination method

$$\begin{cases} x + y + z = 4 \\ x + 3y + 3z = 10 \\ 2x + y - z = 3 \end{cases}$$

**Ex 1.5.039** A freight company has shipping orders for two products. The first product has a unit volume of 10 cu ft and weighs 50 lb. The second product's unit volume is 3 cu ft, and it weighs 40 lb. If the company's trucks have 2,290 cu ft of space and can carry 20,700 lb, how many units of each product can be transported in a single shipment with one truck using the entire volume and weight capacity?

**Ex 1.5.044** A biologist has a 40% solution and a 10% solution of the same plant nutrient. How many cubic centimeters of each solution should be mixed to obtain 25 cc of a 22% solution?

## Chapter R Section R1.6 Part Deux

**1.6.033** If the demand for a pair of shoes is given by  $2p + 5q = 200$  and the supply function for it is  $p - 2q = 10$ , compare the quantity demanded and the quantity supplied when the price is \$90.

quantity demanded \_\_\_\_\_ pairs of shoes

quantity supplied \_\_\_\_\_ pairs of shoes

Will there be a surplus or shortfall at this price?

**1.6.044** Find the market equilibrium point for the following demand and supply functions.

**Demand:**  $p = -2q + 318$

**Supply:**  $p = 8q + 1$

$(q, p) =$

**1.6.049.EP** A group of retailers will buy 104 televisions from a wholesaler if the price is \$325 and 144 if the price is \$275. The wholesaler is willing to supply 84 if the price is \$255 and 164 if the price is \$345. Assume that the resulting supply and demand functions are linear. Let  $p$  represent price (in dollars) and  $q$  represent quantity.

State the two ordered pairs for the demand function in the form  $(q, p)$ .

$$(q, p) =$$

$$(q, p) =$$

Write the demand function in terms of  $q$ .

$$p =$$

State the two ordered pairs for the supply function in the form  $(q, p)$ .

$$(q, p) =$$

$$(q, p) =$$

Write the supply function in terms of  $q$ .

$$p =$$

Find the equilibrium point for the market in the form  $(q, p)$ .

$$(q, p) =$$

## Chapter 3 Section 3



$$\begin{cases} x + y + z = 4 \\ x + 3y + 3z = 10 \\ 2x + y - z = 3 \end{cases}$$

**Ex 01** Make the augmented matrix

$$\begin{cases} x & & - & 4z & = & 1 \\ 2x & - & y & - & 6z & = & 4 \\ 2x & + & 3y & - & 2z & = & 8 \end{cases}$$

**Ex 02** Make the augmented matrix and solve.

$$\begin{cases} x - 3y + z = 0 \\ y - z = 3 \\ z = -2 \end{cases}$$

**Ex HarMathAp12 3.3.017** A system of linear equations and a reduced matrix for the system are given.

$$\begin{cases} x + 3y + 2z = 4 \\ 3x - y = 2 \\ x + 3y + 2z = 5 \end{cases} \quad \left[ \begin{array}{ccc|c} 1 & 0 & \frac{1}{5} & 0 \\ 0 & 1 & \frac{3}{5} & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

1. Use the reduced matrix to find the general solution of the system, if one exists. (If there is no solution, enter NO SOLUTION. If there are infinitely many solutions, express your answers in terms of  $z$  as in Example 3.)
2. If multiple solutions exist, find two specific solutions. (Enter your answers as a comma-separated list of ordered triples. If there is no solution, enter NO SOLUTION.)

**Ex HarMathAp12 3.3.020** A system of linear equations and a reduced matrix for the system are given.

$$\begin{cases} x - y + z = 5 \\ 3x \quad \quad + 2z = 13 \\ x - 4y + 2z = 7 \end{cases} \quad \left[ \begin{array}{ccc|c} 1 & 0 & \frac{2}{3} & \frac{13}{3} \\ 0 & 1 & -\frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 0 & 0 \end{array} \right]$$

1. Use the reduced matrix to find the general solution of the system, if one exists. (If there is no solution, enter NO SOLUTION. If there are infinitely many solutions, express your answers in terms of  $z$  as in Example 3.)
2. If multiple solutions exist, find two specific solutions. (Enter your answers as a comma-separated list of ordered triples. If there is no solution, enter NO SOLUTION.)

**Ex 03** Use Gauss-Jordan Elimination to solve:

$$\begin{cases} x + y = 2 \\ x - 2y = -1 \end{cases}$$

**Ex 04** Use Gauss-Jordan Elimination to solve:

$$\begin{cases} 2x + 3y = 9 \\ x - y = 2 \end{cases}$$

**Ex 05** Use Gauss-Jordan Elimination to solve:

$$\begin{cases} 8x - 3y = -11 \\ 5x - 2y = -6 \end{cases}$$



**Ex 06** Use Gauss-Jordan Elimination to solve:

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

**Ex 07** Use Gauss-Jordan Elimination to solve:

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

**Ex 08** Use Gauss-Jordan Elimination to solve:

$$\begin{cases} x + y + z = 4 \\ x + 3y + 3z = 10 \\ 2x + y - z = 3 \end{cases}$$

**Ex KAUFACS10 11.1.061** The sum of two numbers is 57, and their difference is 17. Find the numbers.

**Ex KAUFACS10 11.1.068** A video store rents new release movies for \$6 and favorites for \$3.25. One day the number of new release movies rented was twice the number of favorites. If the total income from those rentals was \$838.75, how many movies of each kind were rented?

## Chapter 3 Section 3 Numerical

Let's solve this system of equations numerically using the programming language Julia.

$$\begin{cases} 2x - 6y - 12z = -20 \\ 3x - 10y - 20z = -38 \\ 2x - 17z = -40 \end{cases}$$

# Matrix Multiplication

Let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} e & f \\ g & h \end{bmatrix}$$



# Matrix Multiplication

Let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} e & f \\ g & h \end{bmatrix}$$

To find their product

$$AB$$

# Matrix Multiplication

Let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} e & f \\ g & h \end{bmatrix}$$

To find their product

$$AB$$

We do

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{bmatrix}$$

# Matrix Multiplication

Let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} e & f \\ g & h \end{bmatrix}$$

To find their product

$$AB$$

We do

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{bmatrix}$$

This is *messy*!

Let

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \quad B = \begin{bmatrix} 6 & 7 \\ 8 & 9 \end{bmatrix}$$

Have Julia multiply these for you.

## Identity matrix

An identity matrix is a square matrix with 1s along the diagonal and zeros everywhere else.

## Identity matrix

An identity matrix is a square matrix with 1s along the diagonal and zeros everywhere else.

There is an identity matrix for every different size matrix:

$$2 \times 2, 3 \times 3, 4 \times 4, \dots, n \times n, \dots$$

## Identity matrix

An identity matrix is a square matrix with 1s along the diagonal and zeros everywhere else.

There is an identity matrix for every different size matrix:

$$2 \times 2, 3 \times 3, 4 \times 4, \dots, n \times n, \dots$$

$$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \dots$$

## Inverse matrix

If you have a square matrix  $A$  sometimes you can find an inverse for that matrix  $A^{-1}$ .



## Inverse matrix

If you have a square matrix  $A$  sometimes you can find an inverse for that matrix  $A^{-1}$ .

If  $A$  and  $A^{-1}$  are inverse matrices then

$$AA^{-1} = A^{-1}A = I$$

```
julia> A = [2 -6 -12; 3 -10 -20; 2 0 -17]
```

```
3×3 Matrix{Int64}:
```

```
2  -6  -12
```

```
3  -10 -20
```

```
2   0  -17
```

```
julia> A = [2 -6 -12; 3 -10 -20; 2 0 -17]
```

```
3×3 Matrix{Int64}:
```

```
2  -6  -12
```

```
3  -10 -20
```

```
2   0  -17
```

```
julia> Ainv = inv(A)
```

```
3×3 Matrix{Float64}:
```

```
5.0      -3.0      2.22045e-16
```

```
0.323529 -0.294118  0.117647
```

```
0.588235 -0.352941 -0.0588235
```

```
julia> A = [2 -6 -12; 3 -10 -20; 2 0 -17]
```

```
3×3 Matrix{Int64}:
```

```
2  -6  -12
```

```
3  -10 -20
```

```
2   0  -17
```

```
julia> Ainv = inv(A)
```

```
3×3 Matrix{Float64}:
```

```
5.0      -3.0      2.22045e-16
```

```
0.323529 -0.294118  0.117647
```

```
0.588235 -0.352941 -0.0588235
```

```
julia> Ainv*A
```

```
3×3 Matrix{Float64}:
```

```
1.0      -3.55271e-15 -1.08802e-14
```

```
1.11022e-16  1.0      0.0
```

```
-1.52656e-16  0.0      1.0
```

```
julia> A = [2 -6 -12; 3 -10 -20; 2 0 -17]
```

```
3×3 Matrix{Int64}:
```

```
2  -6  -12
```

```
3  -10 -20
```

```
2   0  -17
```

```
julia> Ainv = inv(A)
```

```
3×3 Matrix{Float64}:
```

```
5.0          -3.0          2.22045e-16
```

```
0.323529    -0.294118     0.117647
```

```
0.588235    -0.352941    -0.0588235
```

```
julia> Ainv*A
```

```
3×3 Matrix{Float64}:
```

```
1.0          -3.55271e-15  -1.08802e-14
```

```
1.11022e-16   1.0          0.0
```

```
-1.52656e-16  0.0          1.0
```

The reason we aren't getting 0s in all the off diagonals is round off errors which is always a major concern when working with mathematics on computers.

Resolving  $ax = b$  using the above idea of a multiplicative inverse we get:

$$ax = b$$

$$a^{-1}ax = a^{-1}b$$

$$1x = a^{-1}b$$

$$x = a^{-1}b$$

Resolving  $ax = b$  using the above idea of a multiplicative inverse we get:

$$ax = b$$

$$a^{-1}ax = a^{-1}b$$

$$1x = a^{-1}b$$

$$x = a^{-1}b$$

So back to

$$Ax = b$$

$$A^{-1}Ax = A^{-1}b$$

$$Ix = A^{-1}b$$

$$x = A^{-1}b$$

To solve this system

$$\begin{cases} 2x - 6y - 12z = -20 \\ 3x - 10y - 20z = -38 \\ 2x - 17z = -40 \end{cases}$$

we first need to make the coefficient matrix

```
julia> A = [2 -6 -12; 3 -10 -20; 2 0 -17]
```

```
3×3 Matrix{Int64}:
```

```
2 -6 -12
```

```
3 -10 -20
```

```
2 0 -17
```

and the constant vector

```
julia> b = [-20, -38, -40]
```

```
3-element Vector{Int64}:
```

```
-20
```

```
-38
```

```
-40
```



Now we can solve for  $x$  using Julia

```
julia> x = Ainv*b  
3-element Vector{Float64}:  
13.999999999999999  
0.0  
4.0000000000000002
```

Since this is such an important and often used sequence of operations, Julia provides an operator to do this:

```
julia> x = A\b  
3-element Vector{Float64}:  
14.000000000000001  
1.0658141036401504e-15  
4.0000000000000001
```

To solve

$$\begin{cases} x + 2y + 3z = 1 \\ 2x - y = 2 \\ x + 2y + 3z = 2 \end{cases}$$

```
julia> A = [1 2 3; 2 -1 0; 1 2 3]
```

```
3×3 Matrix{Int64}:
```

```
1 2 3
2 -1 0
1 2 3
```

```
julia> b = [1;2;2]
```

```
3-element Vector{Int64}:
```

```
1
2
2
```

```
julia> x = A\b  
ERROR: LinearAlgebra.SingularException(3)  
Stacktrace:  
 [1] checknonsingular  
 @ \.julia\...\src\factorization.jl:68 [inlined]  
 ...
```

To avoid this issue we use the determinant function

`det`

from the package `LinearAlgebra`. To add a package you can follow *doggo dot jl*'s video instructions at the clickable link:  
[10x27] [How to use External Packages in Julia](#)

Load the LinearAlgebra using:

```
using LinearAlgebra # Needed for the determinant function: det(A)
```

Looking at three examples:

```
A = [1 -3 1; 0 1 -1; 0 0 1]  
det(A)
```

This gives the result

```
3×3 Matrix{Int64}:  
 1 -3 1  
 0 1 -1  
 0 0 1  
1.0
```

Since the determinant  $\det(A) \neq 0$  we can solve using the  $\backslash$  operator.

But for the following two examples:

$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$   
 $\det(A)$

This gives the result

2×2 MatrixInt64:  
 1 1  
 1 1  
 0.0

$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$   
 $\det(A)$

This gives the result

2×2 MatrixInt64:  
 1 1  
 2 2  
 0.0

In both of these cases the determinant  $\det(A) = 0$  we cannot solve using the  $\backslash$  operator.

For now we will just say that there is no unique solution and will come back later to dissect what's happening.

It is now later...

Recall that

$$\begin{cases} x + y = 2 \\ x - 2y = -1 \end{cases}$$

has one solution.

```
using LinearAlgebra
A = [1 1; 1 2]
b = [2, -1]
det(A)
> 1.0
x, y = A\b
> 2-element Vector{Float64}:
 5.0
-3.0
```

And that

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

has infinitely many solutions since they are actually the same lines.

```
A = [1 1; 2 2]
```

```
b = [1, 2]
```

```
det(A)
```

```
> 0.0
```

And that

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

has no solutions because the lines are parallel.

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$b = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

$$\det(A)$$

$$> 0.0$$



There is a difference between the second and third systems, but we cannot tell that from the determinant. How can we tell the difference?

**Ex 1** Make augmented matrix and solve

$$\begin{cases} x + y = 2 \\ x - 2y = -1 \end{cases}$$

**Ex 2** Make augmented matrix and solve

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

**Ex 3** Make augmented matrix and solve

$$\begin{cases} x + y = 2 \\ x + y = -1 \end{cases}$$

**Ex 4** Make augmented matrix and solve

$$\begin{cases} x + 2y + z = 2 \\ 2x + y + 2z = 1 \\ 3x + 3y + 3z = 3 \end{cases}$$

**Ex 5** Make augmented matrix and solve

$$\begin{cases} x + 2y + z = 2 \\ 2x + y + 2z = 1 \\ 3x + 3y + 3z = 0 \end{cases}$$

Give me a  $3 \times 3$  augmented matrix that is

1. Independent

2. Dependent

3. Inconsistent