

Notes for ddCOSMO/ddPCM

December 1, 2016

1 Intro

Let $\{\mathbf{r}_j\}_{1 \leq j \leq M}$ be the centers of domains, and $\{\rho_j\}_{1 \leq j \leq M}$ be the radii. Let us set

$$U_j^n = U_j(\mathbf{r}_j + \rho_j \mathbf{s}_n)$$

and define:

$$[A_{jj}]_{\ell\ell'}^{mm'} = 2\pi \frac{\varepsilon + 1}{\varepsilon - 1} \delta_{\ell\ell'} \delta_{mm'} - \frac{2\pi}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) U_j^n Y_{\ell'}^{m'}(\mathbf{s}_n)$$

$$[A_{jk}]_{\ell\ell'}^{mm'} = -\frac{4\pi\ell'}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) U_j^n \left(\frac{\rho_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right)^{\ell'+1} Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right)$$

Let us set $\partial_{i,\alpha} = \partial/\partial r_{i,\alpha}$. The action of A is obtained as:

$$[A\phi_j]_\ell^m = \sum_k \sum_{\ell', m'} [A_{jk}]_{\ell\ell'}^{mm'} [\phi_k]_{\ell'}^{m'}$$

and its partial derivative is:

$$\partial_{i,\alpha} [A\phi_j]_l^m = [(\partial_{i,\alpha} A)\phi_j]_l^m + [A\partial_{i,\alpha} \phi]_j]_l^m$$

Let us focus on the first term in the sum. Since the entries of $[A_{jj}]_{\ell\ell'}^{mm'}$ are functions of \mathbf{r}_j only, we obtain:

$$[(\partial_{i,\alpha} A)\phi_j]_\ell^m = \sum_{k \neq j} \sum_{\ell', m'} \partial_{i,\alpha} [A_{jk}]_{\ell\ell'}^{mm'} [\phi_k]_{\ell'}^{m'} \quad i \neq j$$

$$[(\partial_{j,\alpha} A)\phi_j]_\ell^m = \sum_{k \neq j} \sum_{\ell', m'} \partial_{j,\alpha} [A_{jk}]_{\ell\ell'}^{mm'} [\phi_k]_{\ell'}^{m'} + \sum_{\ell', m'} \partial_{j,\alpha} [A_{jj}]_{\ell\ell'}^{mm'} [\phi_j]_{\ell'}^{m'}$$

Then, in the case $i \neq j$, we have:

$$\begin{aligned}
[(\partial_{i,\alpha} A)\phi_j]_\ell^m &= - \sum_{k \neq j} \sum_{\ell', m'} \frac{4\pi\ell'}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) U_j^n \times \\
&\quad \partial_{i,\alpha} \left[\left(\frac{\rho_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right)^{\ell'+1} Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right) \right] [\phi_k]_{\ell'}^{m'} \\
&= - \sum_{\ell', m'} \frac{4\pi\ell'}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) U_j^n \times \\
&\quad \sum_{k \neq j} \partial_{i,\alpha} \left[\left(\frac{\rho_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right)^{\ell'+1} Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right) \right] [\phi_k]_{\ell'}^{m'} \\
&= - \sum_{\ell', m'} \frac{4\pi\ell'}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) U_j^n \times \\
&\quad \partial_{i,\alpha} \left[\left(\frac{\rho_i}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|} \right)^{\ell'+1} Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|} \right) \right] [\phi_i]_{\ell'}^{m'}
\end{aligned}$$

Similarly, when $i = j$, we obtain:

$$\begin{aligned}
[(\partial_{j,\alpha} A)\phi_j]_\ell^m &= - \sum_{k \neq j} \sum_{\ell', m'} \frac{4\pi\ell'}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) \times \\
&\quad \partial_{j,\alpha} \left[U_j^n \left(\frac{\rho_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right)^{\ell'+1} Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right) \right] [\phi_k]_{\ell'}^{m'} + \\
&\quad - \sum_{\ell', m'} \frac{2\pi}{2\ell' + 1} \sum_n w_n Y_\ell^m(\mathbf{s}_n) \partial_{j,\alpha} U_j^n Y_{\ell'}^{m'}(\mathbf{s}_n) [\phi_j]_{\ell'}^{m'}
\end{aligned}$$

Notice that:

$$\begin{aligned}
\partial_{i,\alpha} \left(\frac{\rho_i}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|} \right)^{\ell'+1} &= (\ell' + 1) (\cdot)^{\ell'} \rho_i \frac{r_{j,\alpha} + \rho_j s_{n,\alpha} - r_{i,\alpha}}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|^3} \\
\partial_{j,\alpha} \left(\frac{\rho_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right)^{\ell'+1} &= -(\ell' + 1) (\cdot)^{\ell'} \rho_k \frac{r_{j,\alpha} + \rho_j s_{n,\alpha} - r_{k,\alpha}}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|^3}
\end{aligned}$$

Let $\mathbf{s} = \mathbf{r}/|\mathbf{r}| \in \mathbb{S}^2$. Using index notation we have that:

$$\partial_{i,\alpha} Y = \frac{\partial Y}{\partial s_\beta} \frac{\partial s_\beta}{\partial r_\gamma} \frac{\partial r_\gamma}{\partial r_{i,\alpha}}$$

and:

$$\frac{\partial s_\beta}{\partial r_\gamma} = \frac{\delta_{\beta\gamma}}{|\mathbf{r}|} - \frac{r_\beta r_\gamma}{|\mathbf{r}|^3}$$

Finally, we have that:

$$\begin{aligned} \partial_{i,\alpha} \left[Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|} \right) \right] &= \frac{\partial Y_{\ell'}^{m'}}{\partial s_\beta} \left(\frac{\delta_{\beta\gamma}}{|\mathbf{r}|} - \frac{r_\beta r_\gamma}{|\mathbf{r}|^3} \right) (-\delta_{\gamma\alpha}) = \frac{\partial Y_{\ell'}^{m'}}{\partial s_\beta} \left(\frac{r_\beta r_\alpha}{|\mathbf{r}|^3} - \frac{\delta_{\beta\alpha}}{|\mathbf{r}|} \right) = \\ &= \frac{\partial Y_{\ell'}^{m'}}{\partial s_\beta} \left(\frac{(r_{j,\beta} + \rho_j s_{n,\beta} - r_{i,\beta})(r_{j,\alpha} + \rho_j s_{n,\alpha} - r_{i,\alpha})}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|^3} - \frac{\delta_{\alpha\beta}}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_i|} \right) \end{aligned}$$

and

$$\begin{aligned} \partial_{j,\alpha} \left[Y_{\ell'}^{m'} \left(\frac{\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} \right) \right] &= \frac{\partial Y_{\ell'}^{m'}}{\partial s_\beta} \left(\frac{\delta_{\beta\gamma}}{|\mathbf{r}|} - \frac{r_\beta r_\gamma}{|\mathbf{r}|^3} \right) \delta_{\gamma\alpha} = \frac{\partial Y_{\ell'}^{m'}}{\partial s_\beta} \left(\frac{\delta_{\beta\alpha}}{|\mathbf{r}|} - \frac{r_\beta r_\alpha}{|\mathbf{r}|^3} \right) = \\ &= \frac{\partial Y_{\ell'}^{m'}}{\partial s_\beta} \left(\frac{\delta_{\alpha\beta}}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|} - \frac{(r_{j,\beta} + \rho_j s_{n,\beta} - r_{k,\beta})(r_{j,\alpha} + \rho_j s_{n,\alpha} - r_{k,\alpha})}{|\mathbf{r}_j + \rho_j \mathbf{s}_n - \mathbf{r}_k|^3} \right) \end{aligned}$$

A Appendix