

Kinetic scheme of the non-equilibrium discharge in nitrogen-oxygen mixtures

To cite this article: I A Kossyi *et al* 1992 *Plasma Sources Sci. Technol.* **1** 207

View the [article online](#) for updates and enhancements.

Related content

- [Kinetic processes in a highly-ionized non-equilibrium hydrogen plasma](#)
A A Matveyev and V P Silakov
- [A freely localized microwave discharge for removal of chlorofluorocarbon contamination from the atmosphere](#)
G A Askaryan, G M Batanov, A E Barkhudarov *et al.*
- [Theoretical study of the role of ultraviolet radiation of the non-equilibrium plasma in the dynamics of the microwave discharge in molecular nitrogen](#)
A A Matveyev and V P Silakov

Recent citations

- [Ignition by Electrical Discharges](#)
Detlev Markus *et al*
- [Numerical modeling of repetitive nanosecond discharge in air](#)
S. Rassou *et al*
- [Ionic composition of a humid air plasma under ionizing radiation](#)
A. V. Filippov *et al*

Kinetic scheme of the non-equilibrium discharge in nitrogen–oxygen mixtures

I A Kossyi, A Yu Kostinsky, A A Matveyev and V P Silakov

General Physics Institute, Vavilova St. 38, Moscow, Russia 117924

Received 17 February 1992, in final form 23 July 1992

Abstract. A kinetic scheme for non-equilibrium discharge in nitrogen–oxygen mixtures is developed, which almost wholly describes chemical transformations of particles in the cold ($200\text{ K} \leq T \leq 500\text{ K}$) vibrationally unexcited gas. The kinetic scheme includes processes of excitation of electronic states, destruction and ionization of heavy particles by electron impact, associative ionization, electron attachment and detachment, electron–ion and ion–ion recombination, chemical transformations of neutral particles (in ground and excited electronic states) and ion conversion. On the basis of kinetic modelling in the framework of the kinetic scheme proposed, the influence of the electronic excitations of nitrogen molecules and atoms on air composition dynamics is analysed.

1. Introduction

When solving numerous problems of electrophysics, gaseous lasers physics, plasma chemistry, atmospheric physics and so on, a detailed consideration of the kinetics of non-equilibrium plasma-chemical processes in nitrogen–oxygen mixtures is essential. Channels determining the balance of neutral and charged particles in a non-equilibrium plasma are highly diverse. A vast collection of experimental and theoretical information on separate elementary reactions has been accumulated, but in spite of this, the question of whether the description of physicochemical phenomena on the basis of available kinetic schemes is satisfactory has still not been clarified in a number of cases.

The desire to formulate the most complete variants of kinetic schemes of processes in non-equilibrium nitrogen–oxygen plasma is stimulated in many respects by the need to examine the ecological consequences of realization of expensive projects involving a prolonged influence of powerful electromagnetic radiation on the atmosphere. Specifically, the project of radiowave retranslation by artificial ionized regions [1, 2] should be counted among such projects. It envisages a prolonged maintenance by rather intensive microwave radiation of discharges in the stratosphere which can result in a local degradation of the ozone layer [3]. The requirement for a complete non-equilibrium microwave discharge kinetic scheme for atmospheric air is also important in connection with the problem of electrode-free long-distance energy transfer [4, 5].

The majority of authors have modelled plasma-

chemical processes proceeding in electrical discharges in airlike mixtures on the basis of reactions with nitrogen and oxygen atoms being formed by electron impact dissociation of the initial molecules. In some cases (see, for instance, [6]) the appearance of electronically excited atoms $\text{N}(^2\text{D})$ effectively being oxidized by oxygen molecules is taken into account in reactions of the type $\text{N}_2 + e \Rightarrow 2\text{N} + e$. The influence of oxygen molecules in the state $a^1\Delta_g$ on the process of ozone component formation in the afterglow period was also considered. The role of electronically excited nitrogen molecules in an atmosphere with changing chemical composition is hardly being discussed, but in [7] it was shown for non-equilibrium pulsed discharge in an airlike mixture $[\text{N}_2]:[\text{O}_2] = 4:1$ that, under conditions close to breakdown, reactions with such excited species radically affect the physicochemical kinetics.




This work undertakes an attempt to form a kinetic scheme of non-equilibrium discharge in nitrogen–oxygen mixtures, which could almost wholly describe chemical transformations of particles in the cold ($220\text{ K} \leq T \leq 500\text{ K}$) vibrationally unexcited gas. When selecting experimental and theoretical data on various reactions' rate constants a preference was made, if possible, in favour of fundamental review articles containing a critical analysis of accumulated information and of the most recent publications, taking into account the pre-history of the question under consideration and performed on the basis of the latest achievements of modern knowledge. Results of calculations taking into account the influence of electronic excitations of nitrogen molecules and atoms on air composition dynamics under

pulsed discharge in the framework of the kinetic scheme proposed are also presented.









2. The kinetic scheme of the non-equilibrium discharge

2.1. Excitation of electronic states, destruction and ionization of neutral particles by electron impact

The inclusion of all the electron impact electronic-vibrational excitation reactions of the nitrogen-oxygen mixture particles in the discharge scheme is an original and very laborious task. Its full solution, considering chemical and ion kinetics, is believed to be unexpedient in many cases. Indeed, for the greater part, electronic states of particles are short-lived with respect to quenching processes, radiative transitions and predissociation processes. This circumstance permits to simplify the scheme of molecules and atoms excitation by electron impact, in not too rarefied media, without any noticeable damage. In the framework of such a scheme, we propose considering only states sufficiently stable with respect to stepwise deactivation processes that they can play a noticeable role in the gas particles' chemical transformations balance. As the analysis has shown, the states $A^3\Sigma_u^+$, $B^3\Pi_g$, $a'^1\Sigma_u^-$ and $C^3\Pi_u$ for N_2 molecules, the states $a^1\Delta_g$ and $b^1\Sigma_g^+$ for O_2 molecules, the states 2D and 2P for N atoms and the states 1D and 1S for O atoms should be attributed to such states. Their population occurs not only by direct electron impact excitation but also due to quenching of near higher-lying states by unexcited molecules (see [8] for more details). According to the aforesaid, the kinetics of the electronic states' excitation and dissociation and ionization of the nitrogen-oxygen mixture neutral particles by electron impact can be described by the set of reactions



Higher resonant-radiative states of the N_2 molecule are strongly predissociated [9].



Here and after, for the purpose of shortening notation, the electronic states of molecules and atoms are designated by a single letter.

The rate constant values of reactions (1)–(24) are determined from kinetic calculations based on solving Boltzmann's equation for the electron energy distribution function. We will cite the results of such calculations for the airlike mixture $[N_2]:[O_2] = 4:1$, in a direct electric field. Conformably to the discharge scheme proposed, they can be represented according to [10] in the form of approximation formulae:

$$\log k_1 = -8.4 - 14/\vartheta$$

$$k_2 \simeq k_{e-B} + k_{e-W^3} + k_{e-B'}$$

$$\log k_{e-B} = -8.2 - 14.8/\vartheta$$

$$\log k_{e-W^3} = -8.3 - 15.4/\vartheta$$

$$\log k_{e-B'} = -8.7 - 16.8/\vartheta;$$

$$k_3 \simeq k_{e-a'} + k_{e-a} + k_{e-w^1}$$

$$\log k_{e-a'} = -8.8 - 16.7/\vartheta$$

$$\log k_{e-a} = -8.5 - 17.4/\vartheta$$

$$\log k_{e-w^1} = -8.7 - 17.5/\vartheta$$

$$k_4 \simeq k_{e-C} + k_{e-E} + k_{e-a''}$$

$$\log k_{e-C} = -8.2 - 21.1/\vartheta$$

$$\log k_{e-E} = -10.1 - 25.4/\vartheta$$

$$\log k_{e-a''} = -9.2 - 26.2/\vartheta$$

$$\log k_5 = \begin{cases} -9.5 - 5.2/\vartheta & \text{at } \vartheta < 4 \\ -10.2 - 0.35/\vartheta & \text{at } \vartheta > 4 \end{cases}$$

$$\log k_6 = \begin{cases} -9.5 - 6/\vartheta & \text{at } \vartheta < 3 \\ -11.2 - 0.72/\vartheta & \text{at } \vartheta > 3 \end{cases}$$

$$\log k_{13} \simeq \log k_{(X \rightarrow A + c^1 + C^3)} = -7.9 - 13.4/\vartheta$$

$$\log k_{14} \simeq \log k_{(X \rightarrow B^3 + ^3\Pi_g)} = -8 - 16.9/\vartheta$$

$$\log k_{15} \simeq \log k_{(X \rightarrow 9.7-12.1 \text{ eV})} = -8.8 - 11.9/\vartheta$$

$$\log k_{20} = -8.3-36.5/\vartheta$$

$$\log k_{21} = -8.8-28.1/\vartheta$$

$$\log k_{23} = \begin{cases} -9.3-12.3/\vartheta & \text{at } \vartheta > 8 \\ -10.2-5.7/\vartheta & \text{at } \vartheta > 8. \end{cases}$$

The values of a parameter $\vartheta = E/n$ in these formulae are taken in units of 10^{-16} V cm^2 (E is the electric field strength in V cm^{-1} , n is the concentration of N_2 and O_2 molecules in cm^{-3}); $k_{e-\alpha}$ is a rate constant of electron impact excitation of the α th electronic state of the N_2 molecule; $k_{(X \rightarrow A + c^1 + C^3)}$, $k_{(X \rightarrow B^3 + ^3\Pi_g)}$ and $k_{(X \rightarrow 9.7-12.1 \text{ eV})}$ are rate constants of O_2 molecule dissociation through electronically excited levels higher than the $b^1\Sigma_g^+$ level. The approximations given in [10] are correct in the region $3 \leq \vartheta < 30$.

Rate constants values of the reactions (7)–(9) and (10)–(12) can be obtained by using data on these reactions' cross sections adduced correspondingly in [11] and [12]. This allows a correct description of discharge formation of the excited nitrogen atoms playing a significant role in the overall balance of the mixture particles' chemical transformations.

It should be noted that dissociation of the N_2 molecule by electron impact takes place in two stages: first electrons excite corresponding electronic-vibrational levels of the molecule, and then the particle may transfer from these levels to the decay electronic term. In the frame of the scheme indicated, the rate of the dissociation reaction is limited by the second stage of the process. Thus, if the gas is dense enough, then due to the quick quenching of electronically excited molecules, the efficiency of the N-atom production may be much lower than in the hypothetical case of instantaneously proceeding to the final stage of the predissociation. According to [9] this should be ascribed, on the whole, to reaction (7).

The rate constant of ozone dissociation [13] by electron impact (19) is approximately ten times the rate constant of molecular oxygen dissociation:

$$k_{19} \simeq 10(k_{13} + k_{14} + k_{15})$$



Rate constants of the oxygen atom excitation reactions (16)–(18) may be determined with the aid of the corresponding cross sections calculated in [12]. Cross sections of electron impact ionization of NO and electron dissociative attachment to N_2O are given in [14] and [15].

It should be taken into account that, in the case under consideration, threshold kinetic processes of electron interaction with neutral particles of the nitrogen-oxygen mixture are not exhausted by reactions (1)–(24). Conditions may be realized when this set of reactions must be supplemented with stepwise excitation and ionization processes, and also with processes of electron interaction with other components of the mixture.


2.2. Associative ionization

Investigations devoted to the stability of the non-self-consistent discharge in pure nitrogen and in nitrogen


with oxygen admixtures under high pressure [16, 17] have revealed the possibility of a determining influence of associative ionization processes of electronically excited metastable species on the discharge development (see [8]). In the proposed kinetic scheme, such processes are represented by the reactions [18]



Additional channels of the associative ionization are




and










The rate constant of reaction (28) can be set equal in order of magnitude to the rate constant of the high-threshold reaction $\text{O} + \text{N}(\text{D}) \Rightarrow \text{NO}^+ + e$ [20], on condition that the activation barrier is absent. Then we get $k_{28} \sim 10^{-12} \text{ cm}^3 \text{ s}^{-1}$.

2.3. Recombination of electrons and positive ions



The highest rate in this group of processes is usually characteristic of the dissociative recombination reactions






Here and after the electron temperature T_e means the quantity $(2\varepsilon_e)/(3k)$ (ε_e is the mean electron energy); T_e is in kelvin





(according to [23] there should be fulfilled an inequality $k_{37} \ll 3 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$);





Electron-ion recombination is also possible in three-particle processes:





where $\text{A}^+ = \text{N}_2^+, \text{O}_2^+, \text{NO}^+, \text{N}^+, \text{O}^+$; $\text{M} = \text{N}_2, \text{O}_2$;

$$k_{43} = 10^{-19} \times (300/T_e)^{4.5} \text{ cm}^6 \text{ s}^{-1} \quad [25]$$

$$k_{44} = 6 \times 10^{-27} \times (300/T_e)^{1.5} \text{ cm}^6 \text{ s}^{-1} \quad [26]$$

2.4. Electron attachment and detachment processes

Under electric field the electron-loss processes with formation of negative ions are determined basically by the high-threshold reaction of electron dissociative attachment to oxygen (23) and reactions



An approximation of experimental data adduced in [27] gives the following expressions for k_{45} and k_{46} :

$$k_{45} = 1.4 \times 10^{-29} \times \frac{300}{T_e} \times \exp\left(-\frac{600}{T}\right) \times \exp\left(\frac{700 \times (T_e - T)}{T_e \times T}\right) \text{ cm}^6 \text{ s}^{-1}$$







$$k_{46} = 1.07 \times 10^{-31} \times \left(\frac{300}{T_e}\right)^2 \times \exp\left(-\frac{70}{T}\right) \times \exp\left(\frac{1500 \times (T_e - T)}{T_e \times T}\right) \text{ cm}^6 \text{ s}^{-1}.$$

Since cross sections of reactions (45) and (46) have the greatest values in the low-energy region, the employment of the approximative dependences of their rate constants on the electron temperature holds under an electric field. From another point of view, three-body attachment rate constants can be determined by averaging these processes' cross sections over the electron energy distribution function. On the basis of this approach [10], an approximation formula for the nitrogen-oxygen mixture $[\text{N}_2]:[\text{O}_2] = 4:1$ has been derived:


$$k_{45} = (4.7 - 0.259) \times 10^{-31} \text{ cm}^6 \text{ s}^{-1}$$

correct in the area $1 \leq \beta \leq 10$.



With the appearance of oxygen atoms, ozone and nitrogen oxides, an influence upon the negative-ion balance can be exerted by the following reactions:





Electron attachment to the NO_2 molecule actually has a three-particle nature, but the intermediate state $(\text{NO}_2^-)^*$ has a long lifetime, so reaction (52) may be treated as a two-particle one over a wide range of gas pressure [27, 31].



The rate constant of this reaction for $\text{M} = \text{N}_2, \text{O}_2$ is unknown. On the basis of data for other gases [27] one may accept the approximation $k_{53} \approx 10^{-30} \text{ cm}^6 \text{ s}^{-1}$. From [33],











The detachment of electrons from the O^- ions by collisions with unexcited particles N_2 and O_2 under conditions interesting for us (when the gas temperature is not too high) practically does not occur (see [34]). As for the O_2^- ion, on account of lesser electron- O_2 affinity energy in comparison with the O atom reactions [24],









became noticeable already at $T \approx 400 \text{ K}$.

The electron detachment process effectively proceeds, when electronically excited molecules of nitrogen and oxygen are accumulated in the gas. In that case, the detachment mechanism is determined by the reactions







With formation of atomic oxygen and nitrogen in the discharge, associative detachment processes become a rather important channel of destruction of negative ions






















Reactions (66)–(71) can be supplemented by two additional reactions










2.5. Chemical transformations of mixtures of neutral components

2.5.1. Processes with participation of ground electronic state particles. The most detailed and reliable information on chemical reactions between unexcited neutral particles of the nitrogen-oxygen mixture is contained in [39]. In reference to our scheme it looks thus:



†When writing formulae (90)–(92), it was taken into account that the recombination of particles N and O should result in formation of molecules in electronic states with a dissociative limit corresponding to the taken combination of atoms. Values of partial rate constants of specified reactions may be approximately determined proceeding from the condition of the proportionality of the recombination stream to the statistical weight of the forming molecules' electronic states (see [40], Chapter 8, Section 1).



Rate constants of reactions (96)–(99) are determined with the help of the expressions

$$k_\alpha = \frac{k_\alpha^{(a)}(F_{(a)})^{y_\alpha}}{[\text{M}] + k_\alpha^{(a)}/(k_0^{(a)} \sum_i R_i^{(a)} X_i)} \text{ cm}^6 \text{ s}^{-1}$$

$$F_{(a)} = \exp(-T/T_{(a)}) + \exp(-\tilde{T}_{(a)}/T)$$

$$y_\alpha = [1 + \{[\log(k_0^{(a)}[\text{M}] \sum_i R_i^{(a)} X_i / k_\alpha^{(a)})] / v_\alpha\}^2]^{-1}$$

where $\alpha = 96, 97, 98, 99$; $[\text{M}]$ is the total concentration of gas particles in cm^{-3} ; X_i is the relative concentration of the i th component of the mixture.

For $\alpha = 96$

$$k_\alpha^{(96)} = 3 \times 10^{-11} \times (T/300)^{0.3} \text{ cm}^3 \text{ s}^{-1}$$

$$k_0^{(96)} = 6.34 \times 10^{-32} \times (300/T)^{1.8} \text{ cm}^6 \text{ s}^{-1}$$

$$R_{\text{N}_2} = 1.6 \quad R_{\text{O}_2} = 1.3 \quad R_{\text{NO}} = 2.8$$

$$R_{\text{NO}_2} = 10 \quad R_{\text{N}_2\text{O}} = 7$$

$$T_{(96)} = 1800 \text{ K} \quad \tilde{T}_{(96)} = 7200 \text{ K} \quad v_{96} = 1.$$

For $\alpha = 97$

$$k_\alpha^{(97)} = 2.19 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$$

$$k_0^{(97)} = 8.08 \times 10^{-33} \times (1000/T)^2 \text{ cm}^6 \text{ s}^{-1}$$

$$R_{\text{N}_2} = R_{\text{O}_2} = 1 \quad T_{(97)} = 1300 \text{ K} \quad \tilde{T}_{(97)} = 5200 \text{ K}$$

$$v_{97} = 1.$$

For $\alpha = 98$

$$k_\alpha^{(98)} = 8.3 \times 10^{-13} \times (T/300)^{1.7} \text{ cm}^3 \text{ s}^{-1}$$

$$k_0^{(98)} = 6.62 \times 10^{-34} \times (300/T)^{2.5} \text{ cm}^6 \text{ s}^{-1}$$

$$R_{\text{N}_2} = R_{\text{O}_2} = 0.5 R_{\text{NO}_2} = 0.5 R_{\text{N}_2\text{O}_4} = 1 \quad T_{(98)} = 250 \text{ K}$$

$$\tilde{T}_{(98)} = 1050 \text{ K} \quad v_{98} = 0.75 - \log F_{(98)}.$$

For $\alpha = 99$

$$k_\alpha^{(99)} = 1.59 \times 10^{-12} \times (T/300)^{0.2} \text{ cm}^3 \text{ s}^{-1}$$

$$k_0^{(99)} = 3.69 \times 10^{-30} \times (300/T)^5 \text{ cm}^6 \text{ s}^{-1}$$

$$R_{\text{N}_2} = R_{\text{O}_2} = 1 \quad R_{\text{NO}} = 1.3 \quad R_{\text{N}_2\text{O}_5} = 4.4$$


$$T_{(99)} = 250 \text{ K} \quad \tilde{T}_{(99)} = 1050 \text{ K}$$

$$v_{99} = 0.75 - \log F_{(99)}.$$


2.5.2. Processes with participation of electronically excited particles.

It was said above that a correct description of chemical transformations of the mixture's neutral components requires a consideration of the electronic kinetics of gas particles. Such consideration is especially necessary under conditions when $E_{\text{eff}}/n \geq 10^{-15} \text{ V cm}^2$ (E_{eff} is the effective electric field strength). It is known [41] that in this case excitation of the N_2 and O_2 molecules' electronic states takes about half of the energy deposited in the discharge.


A comparatively complete analysis of the chemical kinetics stipulated by the presence of electronically excited particles in the gas can be realized on the basis of the following system of reactions




$$k_{100} = 2.54 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1} \quad [42] \quad (100)$$




$$k_{101} = 7.8 \times 10^{-14} \text{ cm}^3 \text{ s}^{-1} \quad [42] \quad (101)$$



$$k_{102} \approx 7 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1} \quad [43] \quad (102)$$




$$k_{103} = 10^{-11} \text{ cm}^3 \text{ s}^{-1} \quad [44, 45] \quad (103)$$




$$k_{104} \approx 2 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1} \quad [46]$$

$$k_{104} \approx 1.6 \times 10^{-10} \times (300/T)^{2.64} \text{ cm}^3 \text{ s}^{-1} \quad [47].$$


The former value of the quantity k_{104} was determined in [46] on the basis of the experimental data where ultra-violet radiation caused the $\text{N}_2(\text{A})$ state excitation and did not affect the vibrational degrees of freedom. The higher value for k_{104} was obtained in [47] under conditions of strong vibrational non-equilibrium of the gas, when an intensive mixing of the molecules' excited electronic states could occur by interaction with the vibrational reservoir of the ground state.



$$k_{105} = 3 \times 10^{-18} \text{ cm}^3 \text{ s}^{-1} \quad [48] \quad (105)$$




$$k_{106} = 1.29 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1} \quad [42] \quad (106)$$








$$k_{107} = 5 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1} \quad [40] \quad (107)$$







[†] It was shown in [46] that, in vibrationally unexcited gas, excitation of the nitrogen molecules' electronic states in reactions of the type $\text{N}_2(\text{A}) + \text{N}_2(\text{A}) \Rightarrow \text{N}_2(\text{Y}) + \text{N}_2(\text{X})$ ($\text{Y} = \text{B}, \text{W}, \text{B}'$ and so on) must occur in the afterglow period, mainly by channel (104).



† The analysis of the N_2 and O_2 molecules' electronic terms scheme shows that the channel of the dissociative quenching of the $N_2(B)$, $N_2(a')$ and $N_2(C)$ states corresponds to low values of the Messiah parameter. So reactions (113), (115) and (119) can be treated as the main channel of quenching of the $N_2(B)$, $N_2(a')$ and $N_2(C)$ states by unexcited oxygen molecules. The present suggestion seems to be quite probable because, in reference to the $N_2(A)$ state, it agrees with the known result (see [42]) on the principal role of channel (100) in the processes of quenching of the $N_2(A)$ molecules by unexcited oxygen molecules.





$$k_{147} = k_{148} = 1.2 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1} \quad [52]$$





$$k_{153} + k_{154} = 4.3 \times 10^{-12} \times \exp(-850/T) \text{ cm}^3 \text{ s}^{-1} \quad [20]$$

$$k_{153}:k_{154} = 0.69:0.31 \quad [66]$$





$$k_{155} + k_{156} = 5.8 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1} \quad [61]$$

$$k_{155} \approx k_{156}$$







$$k_{157} + k_{158} = 5 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1} \quad [61]$$

$$k_{157}:k_{158} = 0.36:0.64 \quad [67]$$



$$k_{159} + k_{160} = 9.4 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1} \quad [61]$$

$$k_{159}:k_{160} = 0.67:0.33 \quad [67]$$


















2.6. Ion conversion

The process of non-equilibrium discharge in nitrogen–oxygen mixtures is accompanied by the appearance of a wide spectrum of simple and complex ions. Incidentally, concentration of the complex ions increases as the total concentration of particles grows or the gas temperature decreases [21, 40]. The ionic composition depends also upon the electric discharge power.












In our conditions, the conversion of positive and negative ions is described by a system of reactions.

2.6.1. Positive ions




































$$k_{222} = 2.5 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1} \quad [36] \quad (222)$$



2.6.2. Negative ions






Figure 2. Time variation for the concentration of electron-excited nitrogen molecules and atoms in a pulsed microwave discharge. Nitrogen–oxygen mixture $[N_2]:[O_2] = 4:1$; $n = 10^{19} \text{ cm}^{-3}$; $T = 300 \text{ K}$; $E_{\text{eff}}/n = 10^{-15} \text{ V cm}^2$.




Figure 3. Time variation for the concentration of electron-excited oxygen molecules and atoms in a pulsed microwave discharge. Nitrogen–oxygen mixture $[N_2]:[O_2] = 4:1$; $n = 10^{19} \text{ cm}^{-3}$; $T = 300 \text{ K}$; $E_{\text{eff}}/n = 10^{-15} \text{ V cm}^2$.

Excited atoms $N(^2D)$ and $N(^2P)$ are obtained in reactions (8) and (9) of the N_2 molecule dissociation by electron impact and are converted almost entirely to NO. Because the effectiveness of the process (7) of N_2 dissociation with production of two ground state nitrogen atoms is low, with process (102) switched off, concen-





Figure 4. Time variation for the concentration of chemically active particles and chemical reaction products in a pulsed microwave discharge (calculations without taking into account reactions with electron-excited molecules of nitrogen).


trations of $N(^4S)$ and NO components should be equal by the end of the discharge, which is confirmed by the calculations. Further dynamics of the quantities $[N(^4S)]$ and $[NO]$ is conditioned by the reactions




which lead to the complete disappearance of N atoms and to partial decrease of the NO concentration.

The interaction of the excited molecules $N_2(A)$ with oxygen atoms ensures a doubling of the concentration of NO produced during the discharge. Incidentally, the final level of the concentration of all NO_x oxides increases by a factor of five. It should be noted that every interaction leads to the formation of two NO molecules, because the second product of the reaction, namely $N(^2D)$, quickly interacts with molecular oxygen, yielding another NO molecule.

Electronically excited nitrogen molecules determine also the main mechanism of N_2O production:



In the absence of $N_2(A)$ particles, the formation of N_2O proceeds only by the reaction



whose effectiveness under the conditions considered is 25 times lower.

Thus, neglecting the role of electronically excited nitrogen molecules and atoms in formation of chemical products in non-equilibrium nitrogen–oxygen plasmas may lead to crude errors.

4. Conclusion

A kinetic scheme for non-equilibrium discharge in nitrogen-oxygen mixtures is proposed. It is formed from an impressive set of reactions (about 450), with the help of which it is possible to describe the main part of the available experimental data on the chemical and ion composition of cold ($T \leq 500$ K) vibrationally unexcited gas in the discharge and after-discharge periods. In spite of this, however, the question of the role of various mechanisms of excitation, ionization and chemical (ion) transformation in the nitrogen-oxygen plasma cannot be considered to be completely clarified (see [19, 39]). At the same time, the formulated kinetic model, even with all its imperfection, will allow, for instance, more realistic characterization of a wide range of physicochemical phenomena arising from the impact upon the atmosphere of powerful microwave radiation.

Acknowledgments

The authors wish to express their appreciation to N L Aleksandrov and A P Napartovich for helpful discussions.

References

- [1] Gurevich A V 1980 *Sov. Proc. Phys. Sci.* **132** 685-9
- [2] Borisov N D, Gurevich A V and Milih G M 1985 *An Artificial Ionized Area in the Atmosphere* (Moscow: Izmir AN SSSR).
- [3] Askarian G A, Batanov G M, Kossyi I A and Kostinsky A Yu 1988 *Sov. Proc. Acad. Sci. USSR* **302** 566-70
- [4] Ocress E (ed) 1971 *Microwave Energetics* (Moscow: Mir) vol 1
- [5] Askarian G A, Batanov G M and Kossyi I A 1989 *Sov. J. Techn. Phys. Lett.* **15**(8) 18-23
- [6] Larin V F and Rumiantsev S A 1989 *Sov. J. Tech. Phys. Lett.* **15**(6) 87-90
- [7] Kossyi I A, Kostinsky A Yu, Matveyev A A and Silakov V P 1990 *Sov. J. Tech. Phys. Lett.* **16**(12) 57-61
- [8] Silakov V P 1988 *Sov. Plasma Phys.* **14** 1209-13
- [9] Polak L S, Slovetskii D I and Sokolov A S 1972 *Sov. High Energy Chem.* **6** 395-402
- [10] Aleksandrov N L, Vysikailo F I, Islamov R Sh, Kochetov I V, Napartovich A P and Pevgov V G 1981 *Sov. High Temp. Thermophys.* **19** 22-27; 485-9
- [11] Kirillov A S, Konahina A I, Ivanov G S and Ivanov V U 1984 Effective cross sections of electron scattering by atmospheric gases. 2. Molecular nitrogen. PGI AN SSSR. Preprint No 84-05-33 (Apatity)
- [12] Henry R J W, Burke P G and Sinafalaim A L 1969 *Phys. Rev.* **178** 218-24
- [13] Eliasson B, Hirth M and Kogelshatz V 1986 Brown Boweri Research Center, Research Report KLR 86-14OC (Baden).
- [14] Rapp D and Englander-Golden P 1965 *J. Chem. Phys.* **43** 1464-79
- [15] Krishnakumar E and Srivastava S K 1990 *Phys. Rev. A* **41** 2445-52
- [16] Gritsinin S I, Kossyi I A, Silakov V P, Tarasova N M and Terehin V E 1987 *Sov. J. Tech. Phys.* **57** 681-6
- [17] Baizadze K V, Vetsko V M, Lopantseva G V, Napartovich A P, Pal' A F, Perevoznov A F, Popova T E, Starostin A N and Filippov A V 1985 *Sov. Plasma Phys.* **11** 352-60
- [18] Brunet H and Rocca-Serra J 1985 *J. Appl. Phys.* **57** 1574-81
- [19] Mnatsakanyan A H and Naidis G V 1987 *Plasma Chemistry* No 14, ed B M Smirnov (Moscow: Energoatomizdat) pp 227-55
- [20] Bortner M H and Baurer T (eds) 1978 *Defence Nuclear Agency Reaction Rate Handbook* 2nd edn, DNA 1948H (General Electric, TEMPO) ch 24
- [21] Smirnov B M 1983 *Complex ions* (Moscow: Nauka)
- [22] Smith K and Tomson R 1978 *Computer Modeling of Gas Lasers* (New York: Plenum)
- [23] Polak L S, Sergeev P A and Slovetskii D I 1977 *Plasmachemical Reactions and Processes* (Moscow: Nauka) pp 81-107
- [24] Kozlov S I, Vlaskov V A and Smirnova N V 1988 *Sov. Space Res.* **26** 738-45
- [25] Niles F E 1974 Report No. 1702, US Army, Aberdeen Proving Grounds, Maryland 21005 pp 1-107.
- [26] Aleksandrov N L, Konchakov A M, Shachkin L V and Shashkov V M 1986 *Sov. Plasma Phys.* **12** 1218-24
- [27] Aleksandrov N L 1988 *Sov. Proc. Phys. Sci.* **154** 177-206
- [28] Bastien F and Haug R 1975 *J. Chim. Phys.* **72** 105-12
- [29] Pace and Lacombe 1978 *IEEE J. Quantum Electron QE-14* 263-74
- [30] Dettmer J W 1978 *Discharge Processes in the Oxygen Plasma Thesis* (Air Force Institute of Technology, USA)
- [31] Smirnov B M 1978 *Negative Ions* (Moscow: Atomizdat)
- [32] Kajita S, Ushiroda S and Kondo Y 1989 *Proc. 19th Int. Conf. Phenomena in Ionized Gases (Beograd)* vol 4 pp 662-3
- [33] Lopantseva G V, Pal' A F, Persiantsev I G and Popova T E 1982 Influence of oxygen admixtures on the current of the non-selfconsistent discharge in nitrogen Dep. in VINITI no 4392-82 Dep.
- [34] Blank C A, Bortner M H, Baurer T and Feryck A A 1974 *A Pocket Manual of the Physical and Chemical Characteristics of the Earth's Atmosphere* DNA 346711 (General Electric, TEMPO)
- [35] Aleksandrov N L 1978 *Sov. J. Tech. Phys.* **48** 1428-31
- [36] Albritton D L 1978 *Atmos. Nucl. Data Tables* **22** 1-101
- [37] Sabadil H, Bachmann P and Kastelewicz 1980 *Beiträge Plasmaphysik* **20** 283-95
- [38] MacIven M and Phillips L 1978 *Atmosphere Chemistry* (Moscow: Mir)
- [39] Krivonosova O E, Losev S A, Nalivaiko V P, Mukoseev Yu K and Shatalov O P 1987 *Plasma Chemistry* no 14, ed B M Smirnov (Moscow: Energoatomizdat) pp 3-31
- [40] Slovetskii D I 1980 *Mechanisms of Chemical Reactions in the Non-equilibrium Plasma* (Moscow: Nauka)
- [41] Rusanov V D and Fridman A A 1984 *Physics of Chemically Active Plasma* (Moscow: Nauka)
- [42] Jannuzzi M P, Jeffries J B and Kaufman F 1982 *Chem. Phys. Lett.* **87** 570-4
- [43] Piper L G 1982 *J. Chem. Phys.* **77** 2373-7
- [44] Black G, Slanger T G, St John G A and Young R A 1969 *J. Chem. Phys.* **51** 116-21
Young R A, Black G and Slanger T G 1969 *J. Chem. Phys.* **50** 303-8
- [45] Taylor G W and Setser D W 1971 *J. Am. Chem. Soc.* **93** 4930-2
- [46] Silakov V P 1990 The mechanism of molecular nitrogen long-living plasma maintenance under high pressure. Preprint MIFI no 010-90
- [47] Golubovsky Yu B, Telezhko V M and Stoyanov D G 1990 *Sov. Opt. Spectrosc.* **69** 322-7

- [48] Clark W G and Setser D W 1980 *J. Phys. Chem.* **84** 2225–33
- [49] Radsig A A and Smirnov B M 1980 *Reference Book on Atomic and Molecular Physics* (Moscow: Atomizdat)
- [50] Dreyer J W and Perner D 1972 *Chem. Phys. Lett.* **16** 169–73
- [51] Piper L G 1987 *J. Chem. Phys.* **87** 1625–9
- [52] Arnold J and Comes F J 1980 *Chem. Phys.* **47** 125–30
- [53] O'Keefe A, Mauclore G, Parent D and Bowers M T 1986 *J. Chem. Phys.* **84** 215–9
- [54] Dvoryankin A N, Ibragimov L B, Kulagin Yu A and Shelepin L A 1987 *Plasma Chemistry* no 14 ed B M Smirnov (Moscow: Energoatomizdat) pp 102–27
- [55] Yaron M, von Engel A and Vidaud P H 1976 *Chem. Phys. Lett.* **37** 159–61
- [56] Didyukov A I, Kulagin Yu A, Shelepin L A and Yarygina V N 1989 *Sov. Quantum Electron.* **16** 892–904
- [57] Zinn J, Sutherland C D, Stone S N and Duncan L M 1982 *J. Atmos. Terr. Phys.* **44** 1143–71
- [58] O'Brien R J and Myers G H 1970 *J. Chem. Phys.* **53** 3832–5
- [59] Kenner R D and Ogryzlo E A 1980 *Int. J. Chem. Kin.* **12** 501–8
- [60] Yau A W and Shepherd G G 1979 *Planet. Space Sci.* **27** 481–90
- [61] Smirnov B M 1982 *Excited atoms* (Moscow: Energoatomizdat)
- [62] Husain D, Mitra S K and Young A N 1974 *J. Chem. Soc. Faraday Trans. II* **70** 1721–31
- [63] Delcroix J L, Ferreira C M and Ricard A 1973 *Proc. 11th Int. Conf. Phenomena in Ionized Gases (Prague)* pp 301–412
- [64] Donovan R J and Husain D 1970 *Chem. Rev.* **70** 489–516
- [65] Baulch D L, Cox R A, Crutzen P J, Hampson R F, Kerr J A, Troe J A and Watson R T 1982 *J. Phys. Chem. Ref. Data* **11** 327–496
- [66] Slanger T G and Black G 1978 *J. Chem. Phys.* **68** 998–1000
- [67] Slanger T G and Black G 1978 *J. Chem. Phys.* **68** 989–97
- [68] Slanger T G and Black G 1976 *J. Chem. Phys.* **64** 3763–6
- [69] Niles F E 1970 *J. Chem. Phys.* **52** 408–24
- [70] Raizer Yu P 1987 *Gaseous Discharge Physics* (Moscow: Nauka)
- [71] Batanov G M, Gritsinin S I, Kossyi I A, Magunov A N, Silakov V P and Tarasova N M 1985 *Proc. Phys. Inst. Acad. Sci. USSR* **160** 174–203