

Homework 1

Michael Brodskiy

Professor: D. Wood

September 9, 2023

1. Calculate:

(a) $\vec{\nabla} \left(\frac{1}{r} \right)$

We know $r = \sqrt{x^2 + y^2 + z^2}$. Converting and applying the chain rule, we get:

$$\begin{aligned} \frac{\partial}{\partial x} (x^2 + y^2 + z^2)^{\frac{1}{2}} \hat{x} + \frac{\partial}{\partial y} (x^2 + y^2 + z^2)^{\frac{1}{2}} \hat{y} + \frac{\partial}{\partial z} (x^2 + y^2 + z^2)^{\frac{1}{2}} \hat{z} \\ (x^2 + y^2 + z^2)^{-\frac{3}{2}} (x\hat{x} + y\hat{y} + z\hat{z}) \\ \frac{x\hat{x} + y\hat{y} + z\hat{z}}{r^3} \end{aligned}$$

We also know that $\vec{r} = x\hat{x} + y\hat{y} + z\hat{z}$, and that $r\hat{r} = \vec{r}$. Thus, we get:

$$\boxed{\vec{\nabla} \left(\frac{1}{r} \right) = \frac{\vec{r}}{r^3} \rightarrow \frac{\hat{r}}{r^2}}$$

(b) $\vec{\nabla} \cdot \hat{x}$

This implies the following:

$$v_x = 1\hat{x}, \quad v_y = 0\hat{y}, \quad v_z = 0\hat{z}$$

Thus, we find:

$$\boxed{\vec{\nabla} \cdot \hat{x} = \frac{\partial}{\partial x} (1\hat{x}) = 0}$$

(c) $\vec{\nabla} \cdot \hat{r}$

This could be written as:

$$\vec{\nabla} \cdot \left(\frac{x\hat{x} + y\hat{y} + z\hat{z}}{\sqrt{x^2 + y^2 + z^2}} \right)$$

Thus, we need to compute:

$$\frac{\partial}{\partial x} \left(\frac{x}{\sqrt{x^2 + y^2 + z^2}} \right)$$

with respect to each variable. By the quotient rule, we find

$$\frac{\partial}{\partial x} \left(\frac{x}{\sqrt{x^2 + y^2 + z^2}} \right) \rightarrow \frac{\sqrt{x^2 + y^2 + z^2} - x^2(x^2 + y^2 + z^2)^{-0.5}}{x^2 + y^2 + z^2}$$

Converting back to r , we obtain:

$$\frac{1}{r} - \frac{x^2}{r^3}$$

By symmetry, we know that the corresponding y and z variables become:

$$\frac{1}{r} - \frac{y^2}{r^3} \quad \text{and} \quad \frac{1}{r} - \frac{z^2}{r^3}$$

Summing the results, we get:

$$\begin{aligned} & \left(\frac{1}{r} - \frac{x^2}{r^3} \right) + \left(\frac{1}{r} - \frac{y^2}{r^3} \right) + \left(\frac{1}{r} - \frac{z^2}{r^3} \right) \\ & \frac{3}{r} - \left(\frac{x^2 + y^2 + z^2}{r^3} \right) \\ & \boxed{\vec{\nabla} \cdot \hat{r} = \frac{3}{r} - \frac{1}{r} = \frac{2}{r}} \end{aligned}$$

(d) $\vec{\nabla} r^n \ (n > 0)$

$$\begin{aligned} \vec{\nabla}((x^2 + y^2 + z^2)^{\frac{n}{2}}) & \Rightarrow \frac{n}{2}(x^2 + y^2 + z^2)^{\frac{n}{2}-1} (2x\hat{x} + 2y\hat{y} + 2z\hat{z}) \\ n\vec{r}r^{n-2} & \rightarrow \frac{n\vec{r}}{r^{2-n}} \end{aligned}$$

Thus, we get:

$$\boxed{\frac{n\hat{r}}{r^{1-n}} \quad \text{or} \quad n\hat{r}r^{n-1}}$$

2. Calculate the divergence and curls of the following functions:

(a) $\vec{v}_a = xy\hat{x} + yz\hat{y} + zy\hat{z}$

- Divergence:

$$\vec{\nabla} \cdot \vec{v}_a = \frac{\partial}{\partial x}(xy) + \frac{\partial}{\partial y}(yz) + \frac{\partial}{\partial z}(zy)$$

$$\boxed{\operatorname{div}(\vec{v}_a) = y + z + y = 2y + z}$$

- Curl:

$$\vec{\nabla} \times \vec{v}_a = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy & yz & zy \end{vmatrix} = (z - y)\hat{x} - (0 - 0)\hat{y} + (0 - x)\hat{z}$$

$$\boxed{\operatorname{curl}(\vec{v}_a) = \langle z - y, 0, -x \rangle}$$

(b) $\vec{v}_b = y^2\hat{x} + (2xy + z^2)\hat{y} + 2xz\hat{z}$

- Divergence:

$$\vec{\nabla} \cdot \vec{v}_b = \frac{\partial}{\partial x}(y^2) + \frac{\partial}{\partial y}(2xy + z^2) + \frac{\partial}{\partial z}(2xz)$$

$$\boxed{\operatorname{div}(\vec{v}_b) = 0 + 2x + 2x = 4x}$$

- Curl:

$$\vec{\nabla} \times \vec{v}_b = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 & 2xy + z^2 & 2xz \end{vmatrix} = (0 - 2z)\hat{x} - (2z - 0)\hat{y} + (2y - 2y)\hat{z}$$

$$\boxed{\operatorname{curl}(\vec{v}_b) = \langle -2z, -2z, 0 \rangle}$$

(c) $\vec{v}_c = yz\hat{x} + xz\hat{y} + xy\hat{z}$

- Divergence:

$$\vec{\nabla} \cdot \vec{v}_c = \frac{\partial}{\partial x}(yz) + \frac{\partial}{\partial y}(xz) + \frac{\partial}{\partial z}(xy)$$

$$\boxed{\operatorname{div}(\vec{v}_c) = 0 + 0 + 0 = 0}$$

- Curl:

$$\vec{\nabla} \times \vec{v}_c = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ yz & xz & xy \end{vmatrix} = (x - x)\hat{x} - (y - y)\hat{y} + (z - z)\hat{z}$$

$$\boxed{\operatorname{curl}(\vec{v}_c) = \langle 0, 0, 0 \rangle}$$

3. Calculate the following. Note that some results should be vectors. It is helpful to check the dimensionality of your answers:

(a) $(\vec{r} \cdot \vec{\nabla})\vec{r}$

$$(\vec{r} \cdot \vec{\nabla}) = \frac{\partial}{\partial x}(x\hat{x}) + \frac{\partial}{\partial y}(y\hat{y}) + \frac{\partial}{\partial z}(z\hat{z}) = 1 + 1 + 1 = 3$$

$$\boxed{(\vec{r} \cdot \vec{\nabla})\vec{r} = 3\vec{r} = 3x\hat{x} + 3y\hat{y} + 3z\hat{z}}$$

(b) $(\hat{r} \cdot \vec{\nabla})r$

From Problem **1c**, we know the value of $(\hat{r} \cdot \vec{\nabla})$, which gives us:

$$\boxed{\left(\frac{2}{r}\right)r = 2}$$

(c) $(\hat{r} \cdot \vec{\nabla})\hat{r}$

Again, employing what we know from Problem **1c**, we get:

$$\boxed{\left(\frac{2}{r}\right)\hat{r} = \frac{2\hat{r}}{r}}$$

4. (a)

(b)

5. (a)

(b)

(c)

(d)

6. (a)

(b)

7. (a)

(b)

(c)