

Lecture 5 — Einstein's Equations and Schwarzschild

Michael Brodskiy

Professor: J. Blazek

October 7, 2024

- We may begin with Einstein's equations

$$G_{\mu\nu} = 8\pi G T_{\mu\nu}$$

- Note that the tensors form second order non-linear differential equations

- Symmetric Tensor ($n = 4$ dimensions)

- $\frac{n^2-n}{2} + n = \frac{n^2+n}{2} = 10$ degrees of freedom
- General relativity sees diffeomorphism invariance, so four of the degrees of freedom are removed (since it isn't $x^\mu \rightarrow x^{\mu'}$)
- Solving these differential equations is extremely complex, so we will make some assumptions to simplify analysis:
 - * Boundary conditions and initial conditions
 - * Limits
 - * Simplify through symmetry

- Symmetric General Relativity

- Spherical symmetry and static
- Homogenous and isotropic: FLRW universe/cosmology
- $T_{\mu\nu} = 0$, small perturbations are gravitational waves
- In Newtonian mechanics, with three masses M_1 , M_2 , and much smaller M_3 , we may write: $\Phi_{M_3} = \Phi_{M_1} + \Phi_{M_2}$
 - * In General Relativity, $g_{\mu\nu}$ depends on M_1 and M_2 , as well as the binding energy between

- Schwarzschild

- Only vacuum solution with spherical symmetry
- We assume a spherical system that is static
- We use Minkowski space, with spherical coordinates:

$$ds^2 = -dt^2 + dx^2 + dy^2 + dz^2 = -dt^2 + dr^2 + r^2 \underbrace{(d\theta^2 + \sin^2(\theta) d\phi^2)}_{d\Omega^2}$$

- * We may rescale this with functions of r :

$$ds^2 = -A(r) dt^2 + B(r) dr^2 + C(r) r^2 d\Omega^2$$

- * Furthermore, we define $r \rightarrow \sqrt{C(r)}$

$$ds^2 = -A(r) dt^2 + B(r) dr^2 + r^2 d\Omega^2$$

$$ds^2 = -e^{2\alpha(r)} dt^2 + e^{\beta(r)} dr^2 + r^2 d\Omega^2$$

- * For the diagonal metric, we may find $\Gamma_{r\phi}^\phi = (1/r)$, then continuing Christoffel calculations, using Riemann, and then contracting to Ricci, we find:

$$R_{tt} = e^{2(\alpha-\beta)} \left[\partial_r^2 \alpha + (\partial_r \alpha)^2 - \partial_r \alpha \partial_r \beta + \frac{2}{r} \partial_r \alpha \right]$$

$$R_{rr} = -\partial_r^2 \alpha - (\partial_r \alpha)^2 + \partial_r \alpha \partial_r \beta + \frac{2}{r} \partial_r \beta$$

$$R_{\theta\theta} = e^{-2\beta} [r(\partial_r \beta - \partial_r \alpha) - 1] + 1$$

$$R_{\phi\phi} = \sin^2(\theta) R_{\theta\theta}$$

- * We want a $T_{\mu\nu} = 0$ solution, which implies $G_{\mu\nu} = 0$, which then implies $R_{\mu\nu} = 0$
 - This is known as “Ricci Flat” (not really flat)
- * We move terms around to find:

$$e^{2(\beta-\alpha)} R_{tt} + R_{rr} = 0$$

$$\frac{2}{r} (\partial_r \alpha + \partial_r \beta) = 0$$

$$\alpha = -\beta$$

- * Taking:

$$R_{\theta\theta} = 0 \quad \text{and} \quad -e^{2\alpha} [2r \partial_r \alpha + 1] + 1 = 0$$

- * We get:

$$e^{2\alpha} [2r \partial_r \alpha + 1] = \partial_r (r e^{2\alpha})$$

* We define $A(r) = e^{2\alpha}$ and $y(r) = rA(r)$, which gives:

$$y = r + C \Rightarrow A(r) = 1 + \frac{C}{r}$$

$$A(r) = 1 - \frac{R_s}{r}$$

* Where R_s is the Schwarzschild radius, which allows us to write:

$$ds^2 = - \left(1 - \frac{R_s}{r}\right) dt^2 + \left(1 - \frac{R_s}{r}\right)^{-1} dr^2 + r^2 d\Omega$$

- Black Holes

- Using Newtonian $g_{tt} = -(1 + 2\Phi)$, we get:

$$R_s = -2\Phi$$

- With a point mass:

$$\Phi = -\frac{GM}{r}$$

- Thus, $R_s = 2GM$

- Schwarzschild Properties:

1. $M \rightarrow 0, \quad g_{MV} \rightarrow \eta_{M\nu}$

2. $r \rightarrow \infty, \quad g_{M\nu} \rightarrow \eta_{M\nu}$

3. $r = 0, \quad \frac{R_s}{r} \rightarrow \infty$

4. $r = R_s, \quad \left(1 - \frac{R_s}{r}\right)^{-1} \rightarrow \infty$

- For Black Holes, light cones are deformed by null geodesics. We may derive:

$$\frac{dt}{dr} = \pm \left(1 - \frac{2GM}{r}\right)^{-1}$$

* As $r \rightarrow \infty$ back to 45°

- Anthropic Principle