

Lecture 5 — Linear Time Invariant Systems

Michael Brodskiy

Professor: I. Salama

October 2, 2024

- The Impulse Response

- Linear algebra concepts apply to LTI system analysis; if the response to basis input signals are known, any general input can be expressed as a linear combination of these basis signals, allowing the output to be determined
- The system is fully characterized by its responses to the basis signals
- In the time domain representation, we use delayed unit impulse functions as basis signals
- The impulse response, $h(t)/h[n]$, of a linear time invariant system is defined as the response of the system to a unit impulse input, $x(t) = \delta(t)/x[n] = \delta[n]$
- The response of a linear time invariant system with impulse response $h[n]$ to a general input $x[n]$ can be obtained using the convolution sum; the convolution sum sifts through the sequence values, $x[k]$. represented by an impulse located at $n = k$, and finds the corresponding response given by $h[n - k]$

$$x[n] = \sum_{k=-\infty}^{\infty} x[k]\delta[n - k] \implies y[n] = \sum_{k=-\infty}^{\infty} x[k]h[n - k]$$

- The Convolution Sum

- The convolution sum may also be expressed as:

$$y[n] = x[n] * h[n] = h[n] * x[n] = \sum_{k=-\infty}^{\infty} x[n - k]h[k]$$

- Geometric Series

- Sums of geometric series are given by:

$$\sum_{k=0}^n r^k = \frac{(1 - r^{n+1})}{1 - r}$$

$$\sum_{k=n_1}^{n_2} r^k = \frac{(r^{n_1} - r^{n_2+1})}{1 - r}$$

$$\sum_{k=n_1}^{\infty} r^k = \frac{r^{n_1}}{1 - r}$$

* For $|r| < 1$

- Convolution with an Impulse:

$$- x[n] * \delta[n - n_o] = x[n - n_o]$$

- Continuous Time Impulse Response

- The impulse response, $h(t)$, of a linear time invariant system is defined as the response of the system to a unit impulse