

Lecture 6

Michael Brodskiy

Professor: G. Fiete

April 7, 2025

- Harmonic Oscillator

- Classical

$$F = -kx$$
$$V(x) = \frac{1}{2}kx^2$$
$$F = -\frac{dV}{dx}$$

- Quantum

$$E = \frac{p^2}{2m} + \frac{1}{2}kx^2$$
$$H = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2$$
$$\omega = \sqrt{\frac{k}{m}}$$

- * Energy eigenvalues can be found as:

$$E_n = \hbar\omega(n + 1/2)$$

- * We may find:

$$H = \hbar\omega(aa^\dagger + 1/2)$$

- * And from here, we find:

$$[a, a^\dagger] = 1$$

- This indicates that the operators a and a^\dagger raise and lower the energy eigenstates

- We can write this as:

$$a |E\rangle \propto |E - \hbar\omega\rangle$$

$$a^\dagger |E\rangle \propto |E + \hbar\omega\rangle$$

- These are called “ladder operators”
- Note that there is an asymmetry in the ladder, since $aa^\dagger \neq a^\dagger a$
- Since there is a lowest energy state in the harmonic oscillator well, states can not be lowered in energy indefinitely, such that:

$$a |E_{lowest}\rangle = 0$$

- This is called the ladder termination condition

$$H |E_{lowest}\rangle = \hbar\omega(aa^\dagger + 1/2) |E_{lowest}\rangle = \frac{\hbar\omega}{2} |E_{lowest}\rangle$$

- Thus, we may conclude that the lowest energy is $\hbar\omega/2$
- Since this is finite, we say the quantum mechanical ground state has a zero-point energy of $\hbar\omega/2$

- Excited States

- We may obtain:

$$|n\rangle = \frac{(a^\dagger)^n}{\sqrt{n!}} |0\rangle$$

$$\phi_n(x) = \frac{1}{\sqrt{n!}} \left[\sqrt{\frac{m\omega}{2\hbar}} \left(x - \frac{\hbar}{m\omega} \frac{d}{dx} \right)^n \right] \phi_0(x)$$

- Dirac Notation

$$|n\rangle = |\phi_n\rangle = |E_n\rangle = |(n + 1/2)\hbar\omega\rangle$$

- With $\phi_n(x) = \langle x|n\rangle$
- Since $\langle n|n\rangle = 1$, and $\int_{-\infty}^{\infty} |x\rangle \langle x| dx = \mathbb{1}$

$$1 = \langle n| \int_{-\infty}^{\infty} |x\rangle \langle x| dx |n\rangle = \int_{-\infty}^{\infty} \phi_n^*(x) \phi_n(x) dx$$

- By orthonormality, we have:

$$\delta_{mn} = \langle m|n\rangle = \int_{-\infty}^{\infty} \phi_m^*(x) \phi_n(x) dx$$

- Since the Hermitian operator states are eigenstates of the Hamiltonian, they form a complete set of states, such that:

$$\sum_{n=0}^{\infty} |n\rangle \langle n| = \mathbb{1}$$

- A general state $|\psi\rangle$ can be written as:

$$|\psi\rangle = \sum_{n=0}^{\infty} \underbrace{(\langle n|\psi\rangle)}_{c_n} |n\rangle$$

- We know:

$$c_n = \int_{-\infty}^{\infty} \phi_n^*(x) \psi(x) dx$$

- The probability of the state $|\psi\rangle$ having energy E_n is:

$$P_{E_n} = |\langle n|\psi\rangle|^2 = |c_n|^2$$