

Lecture 4

Michael Brodskiy

Professor: G. Fiete

February 19, 2025

- Wave Equation for Unidimensional Particle

- In classical mechanics, we have the energy problem, with E as the total energy, T as the kinetic energy, and V as the potential energy:

$$E = T + V = \frac{p^2}{2m} + V(x)$$

- In quantum mechanics, we may define the Hamiltonian as:

$$\hat{H} = \frac{\hat{p}}{2m} + V(\hat{x})$$

- * Where \hat{p} is the momentum operator and \hat{x} is the position

- From here, the time-independent Schrödinger equation may be written as:

$$\hat{H}\phi_E(x) = E\phi_E(x)$$

- * Where $\phi_E(x)$ represents the wave function/eigenfunction

- We may continue to get:

$$\hat{p} = -i\hbar \frac{d}{dx}, \quad \hat{x} = x$$

- * This can be used to obtain:

$$\hat{p}\phi(x) = -i\hbar \frac{d}{dx}\phi(x), \quad \hat{x}\phi(x) = x\phi(x)$$

- From here, we get:

$$\hat{H}\psi(x) = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2}\psi(x) + V(x)\psi(x) = E\psi(x)$$

- The above is the wave equation we need to solve. The wave function can generically be written as:

$$\psi(x) = \sum_{n=0}^d \psi_n \phi_{E_n}(x)$$

- * Note that ψ_n is a scalar coefficient (projections of ψ along the n -th direction) and $\phi_{E_n}(x)$ represents basis functions
- * Also, note that, from previous lessons, we may recall that the probability of finding the system in a particular eigenstate is:

$$P_{E_n} = |\psi_n|^2$$

- In Dirac notation:

$$|\psi\rangle = \begin{pmatrix} \langle E_1 | \psi \rangle \\ \langle E_2 | \psi \rangle \\ \vdots \\ \langle E_n | \psi \rangle \end{pmatrix}$$

- * And also:

$$\langle \psi | = (\langle E_1 | \psi \rangle^* \quad \langle E_2 | \psi \rangle^* \quad \cdots \quad \langle E_n | \psi \rangle^*)$$

- Change of Basis

- * Changing basis to a position representation allows us to obtain the probability of finding the particle at x as:

$$P_x = |\psi(x)|^2$$

- * This means that $|\psi(x)|^2$ is now a probability density such that:

$$\int_{-\infty}^{\infty} P(x) dx = \int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1$$

- * Furthermore, we may find the probability that the particle is in a certain range as:

$$P[a \leq x \leq b] = \int_a^b P(x) dx = \int_a^b |\psi(x)|^2 dx$$

- * In summary, we determine:

$$\langle x | \psi \rangle = \psi(x)$$

$$\langle \psi | x \rangle = \psi^*(x)$$

$$\hat{A} = \hat{A}(x)$$