## Lecture 2 — Random Variables

Michael Brodskiy

Professor: I. Salama

January 30, 2025

- A random variable is a function that maps the outcomes of a random experiment into a set of real numbers
- The Probability Mass Function (PMF)
  - The PMF is a probability measure that gives us probabilities of the possible values for a random variable
  - The PMF may be defined as:

$$P_x(x) = \begin{cases} P(X=x), & \text{if } x \in S_x \\ 0, & \text{Otherwise} \end{cases}$$

- The probability mass function can be obtained using the probabilities of the corresponding sample space outcomes
- The Bernoulli Random Variable
  - We may write this as X = Bernoulli(p)
  - This can be expressed as:

$$P_x(x) = \begin{cases} 1 - p, & x = 0 \\ p, & x = 1 \\ 0, & \text{otherwise} \end{cases}$$

- The Binomial Random Variable
  - We may write this as X = Binomial(n, p)
  - This can be expressed as:

$$P_x(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k}, & 0 \le k \le n \\ 0, & \text{otherwise} \end{cases}$$

- The Geometric Random Variable
  - We may write this as: X = Geometric(p)
  - This can be expressed as:

$$P_x(x) = \begin{cases} p(1-p)^{x-1}, & x = 1, 2, 3 \dots \\ 0, & \text{otherwise} \end{cases}$$

- The Poisson Random Variable
  - We may write this as: X = Poisson(a)
  - This can be expressed as:

$$P_K(k) = \begin{cases} \frac{a^k}{k!} e^{-a}, & k = 0, 1, 2, 3 \dots \\ 0, & \text{otherwise} \end{cases}$$

- The Poisson random variable can be described using the average arrival rate  $a=\lambda T$
- The Pascal or Negative Binomial Random Variable
  - We may write this as:  $X = \operatorname{Pascal}(k, p)$
  - This can be expressed as:

$$P_X(x) = \begin{cases} \binom{x-1}{k-1} p^k (1-p)^{x-k}, & x = k, k+1, \dots \\ 0, & \text{otherwise} \end{cases}$$

- The Discrete Uniform Random Variable
  - We may write this as: N = Uniform(k, l)
  - This can be expressed as:

$$P_N(n) = \begin{cases} \frac{1}{l-k-1}, & l-k=2,3,4,\cdots \\ 0, & \text{otherwise} \end{cases}$$

- The Cumulative Distribution Function (CDF)
  - The CDF of a discrete random variable is defined for all values of the random variable as:

$$F_x(x) = P[X \le x] = \sum_{i: x_i \le x} P_x(x_i)$$

- For any value x,  $F_x(x)$  is the probability that the observed value of X will be at most x