

Lecture XVI Notes

Michael Brodskiy

Professor: V. Cherkassky

July 13, 2020

1 Mass and Density – 15.4

The mass may be expressed as

$$m = \iint_D \rho(x, y) dA$$

Total charge is expressed as:

$$Q = \iint_D \sigma(x, y) dA$$

The (first) moments on the x and y axis, respectively, may be found by using:

$$M_x = \iint_D y\rho(x, y) dA \text{ and } M_y = \iint_D x\rho(x, y) dA$$

The center of mass, (\bar{x}, \bar{y}) may be found by rearranging the formula:

$$\begin{aligned} m\bar{x} &= M_y \text{ and } m\bar{y} = M_x \\ \bar{x} &= \frac{M_y}{m} = \frac{\iint_D x\rho(x, y) dA}{\iint_D \rho(x, y) dA} \\ \bar{y} &= \frac{M_x}{m} = \frac{\iint_D y\rho(x, y) dA}{\iint_D \rho(x, y) dA} \end{aligned}$$

The moment of inertia (or the second moment) is mr^2 , where r is the distance from a particle to the axis. About the x axis, the moment of inertia is equal to:

$$I_x = \iint_D y^2 \rho(x, y) dA$$

Across the y axis the moment of inertia is equal to:

$$I_y = \iint_D x^2 \rho(x, y) dA$$

In addition to this, the polar moment of inertia (about the origin) is equal to:

$$I_0 \iint_D (x^2 + y^2) \rho(x, y) dA$$

This means that $I_0 = I_x + I_y$

2 Surface Area – 15.5

The surface area, $A(S)$ is defined by:

$$A(S) = \lim_{(m,n) \rightarrow \infty} \sum_{i=1}^m \sum_{j=1}^n \Delta T_{ij}$$

T_{ij} is defined by $|a \times b|$, where:

$$a = \Delta x \hat{\mathbf{i}} + f_x(x_i, y_i) \Delta x \hat{\mathbf{k}}$$

$$b = \Delta y \hat{\mathbf{j}} + f_y(x_i, y_j) \Delta y \hat{\mathbf{k}}$$

Therefore, as an iterated integral, the surface area may be expressed as:

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dA$$