
Matrix multiplication

Matrix multiplication is an operation between two matrices that creates a new matrix such that given two matrices **A** and **B**, each column of the product **AB** is formed by multiplying **A** by each column of **B** (Definition 1).

Note that this definition requires that if we multiply an $m \times n$ matrix by a $n \times p$ matrix, the result will be an $m \times p$ matrix where the number of rows is determined by the first matrix and the number of columns is determined by the second matrix. More succinctly:

$$\mathbf{A}_{m \times n} \mathbf{B}_{n \times p} = (\mathbf{AB})_{m \times p}$$

Definition 1 *Matrix multiplication between an $m \times n$ matrix **A** and a $n \times p$ matrix **B** is given by*

$$\mathbf{AB} := \begin{bmatrix} \mathbf{Ab}_{*,1} & \mathbf{Ab}_{*,2} & \dots & \mathbf{Ab}_{*,n} \end{bmatrix}$$

Intuition

There are several perspectives for which one can view matrix multiplication each depending on the perspective taken on each of the matrix factors. Recall we can view a matrix via a number of perspectives including

1. As an ordered list of column vectors
2. As an ordered list of row vectors
3. As a linear transformation

These views of matrices lead to the following views of matrix multiplication:

1. *Matrix multiplication as the linear transformation of a set of vectors:* This perspective follows from viewing **A** as a linear transformation and **B** as an ordered list of column-vectors.
2. *Matrix multiplication as a composition of linear transformations:* This perspective follows from viewing both **A** and **B** as linear transformations.
3. *Matrix multiplication as the computation of all pair-wise dot products between two lists of vectors:* This perspective follows from viewing **A** as an ordered list of row-vectors and viewing **B** as an ordered list of column-vectors.

These perspectives are discussed in more depth in the following sections.

1. Matrix multiplication as the linear transformation of a set of vectors

The most obvious way to view matrix multiplication is as simply performing a linear transformation on a set of vectors. That is, we view \mathbf{A} as a linear transformation and we view the matrix \mathbf{B} as an ordered list of column vectors

$$\mathbf{B} := [\mathbf{b}_{*,1}, \mathbf{b}_{*,2}, \dots, \mathbf{b}_{*,n}]$$

Then, we form the matrix \mathbf{AB} by taking the linear transformation characterized by \mathbf{A} of each $\mathbf{b}_{*,i}$.

2. Matrix multiplication as a composition of linear transformations

Alternatively if we view *both* \mathbf{A} and \mathbf{B} as linear transformations, then the matrix \mathbf{AB} is the matrix that characterizes the composition of the linear transformations characterized by \mathbf{A} and \mathbf{B} (Theorem 1). That is, given two linear transformations

$$\begin{aligned} T_{\mathbf{A}}(\mathbf{x}) &:= \mathbf{Ax} \\ T_{\mathbf{B}}(\mathbf{x}) &:= \mathbf{Bx} \end{aligned}$$

the matrix resulting from multiplying \mathbf{A} with \mathbf{B} , denoted \mathbf{AB} , is the matrix that characterizes the composition $T_{\mathbf{A}} \circ T_{\mathbf{B}}$. That is, \mathbf{AB} characterizes the linear transformation:

$$T_{\mathbf{AB}}(\mathbf{x}) := \mathbf{A}(\mathbf{Bx})$$

This concept is illustrated in Figure 1.

Recall that a matrix's number of rows determines the dimensions of the vectors in its range and the number of columns correspond to the number of dimensions the domain. Since \mathbf{AB} characterizes the composition $T_{\mathbf{A}} \circ T_{\mathbf{B}}$, it follows that the matrix \mathbf{AB} will map from the domain of \mathbf{B} to the range of \mathbf{A} .

Theorem 1 *Matrix multiplication between an $m \times n$ matrix \mathbf{A} and a $n \times p$ matrix \mathbf{B} results in a matrix \mathbf{AB} such that given a vector $\mathbf{x} \in \mathbb{R}^p$, the following holds:*

$$(\mathbf{AB})\mathbf{x} = \mathbf{A}(\mathbf{Bx})$$

Proof:




Figure 1: The matrix \mathbf{AB} performs the mapping by \mathbf{B} followed by \mathbf{A} .

First, let's expand \mathbf{Bx} .

$$\begin{aligned}\mathbf{Bx} &= \mathbf{b}_{*,1}x_1 + \mathbf{b}_{*,2}x_2 + \cdots + \mathbf{b}_{*,n}x_n \\ &= \begin{bmatrix} b_{1,1}x_1 + b_{1,2}x_2 + \cdots + b_{1,n}x_n \\ b_{2,1}x_1 + b_{2,2}x_2 + \cdots + b_{2,n}x_n \\ \vdots \\ b_{n,1}x_1 + b_{n,2}x_2 + \cdots + b_{n,n}x_n \end{bmatrix}\end{aligned}$$

Now,

$$\begin{aligned}(\mathbf{AB})\mathbf{x} &= [\mathbf{Ab}_{*,1}, \mathbf{Ab}_{*,2}, \dots, \mathbf{Ab}_{*,n}] \mathbf{x} \\ &= \mathbf{Ab}_{*,1}x_1 + \mathbf{Ab}_{*,2}x_2 + \cdots + \mathbf{Ab}_{*,n}x_n \\ &= (\mathbf{a}_{*,1}b_{1,1} + \cdots \mathbf{a}_{*,n}b_{n,1})x_1 + (\mathbf{a}_{*,1}b_{1,2} + \cdots \mathbf{a}_{*,n}b_{n,2})x_2 + \cdots + (\mathbf{a}_{*,1}b_{1,n} + \cdots \mathbf{a}_{*,n}b_{n,n})x_n \\ &= (\mathbf{a}_{*,1}b_{1,1}x_1 + \cdots \mathbf{a}_{*,n}b_{n,1}x_1) + (\mathbf{a}_{*,1}b_{1,2}x_2 + \cdots \mathbf{a}_{*,n}b_{n,2}x_2) + \cdots + (\mathbf{a}_{*,1}b_{1,n}x_n + \cdots \mathbf{a}_{*,n}b_{n,n}x_n) \\ &= \mathbf{a}_{*,1}(b_{1,1}x_1 + \cdots + b_{1,n}x_n) + \mathbf{a}_{*,2}(b_{2,1}x_1 + \cdots + b_{2,n}x_n) + \cdots + \mathbf{a}_{*,n}(b_{n,1}x_1 + \cdots + b_{n,n}x_n) \\ &= \mathbf{a}_{*,1}(\mathbf{Bx})_1 + \mathbf{a}_{*,2}(\mathbf{Bx})_2 + \cdots + \mathbf{a}_{*,n}(\mathbf{Bx})_n \\ &= \mathbf{A}(\mathbf{Bx})\end{aligned}$$

□

3. Matrix multiplication as the computation of all pair-wise dot products between two lists of vectors

If we view the matrix **A** as an ordered list of row-vectors and the matrix **B** as an ordered list of column vectors, then the product **AB** is the matrix that stores all of the pair-wise dot products of the vectors in **A** and **B**. That is, the i, j th element of **AB** is the dot product of the i th row of **A** and the j th column of **B** (Theorem 2). This fact, known as **row-column rule**, can be used for computing each element of **AB**. Figure 2 demonstrates this view of matrix multiplication.




Figure 2: The i, j th element of **AB** is the dot product of the i th row of **A** and j th column of **B**.

Theorem 2 *The i, j th element of **AB** is computed by*

$$\mathbf{a}_{i,*} \cdot \mathbf{b}_{*,j}$$

Proof:

$$\begin{aligned}
 (\mathbf{AB}) &= [\mathbf{Ab}_{*,1}, \mathbf{Ab}_{*,2}, \dots, \mathbf{Ab}_{*,n}] \\
 &= \begin{bmatrix} \mathbf{a}_{*,1}b_{1,1} + \dots + \mathbf{a}_{*,n}b_{n,1}, & \dots & \mathbf{a}_{*,1}b_{1,n} + \dots + \mathbf{a}_{*,n}b_{n,n} \end{bmatrix} \\
 &= \begin{bmatrix} a_{1,1}b_{1,1} + \dots + a_{1,n}b_{n,1} & \dots & a_{1,1}b_{1,n} + \dots + a_{1,n}b_{n,n} \\ a_{2,1}b_{1,1} + \dots + a_{2,n}b_{n,1} & \dots & a_{2,1}b_{1,n} + \dots + a_{2,n}b_{n,n} \\ \vdots & \ddots & \vdots \\ a_{n,1}b_{1,1} + \dots + a_{n,n}b_{n,1} & \dots & a_{n,1}b_{1,n} + \dots + a_{n,n}b_{n,n} \end{bmatrix} \\
 &= \begin{bmatrix} \mathbf{a}_{1,*} \cdot \mathbf{b}_{*,1} & \dots & \mathbf{a}_{1,*} \cdot \mathbf{b}_{*,n} \\ \mathbf{a}_{2,*} \cdot \mathbf{b}_{*,1} & \dots & \mathbf{a}_{2,*} \cdot \mathbf{b}_{*,n} \\ \vdots & \ddots & \vdots \\ \mathbf{a}_{m,*} \cdot \mathbf{b}_{*,1} & \dots & \mathbf{a}_{m,*} \cdot \mathbf{b}_{*,n} \end{bmatrix}
 \end{aligned}$$

□

Properties of matrix multiplication

The following properties hold for matrix multiplication:

1. **Associative law for matrices** (Theorem 3)

$$\mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}$$

2. **Commutative property of scalars** (Theorem 4)

$$r(\mathbf{AB}) = (r\mathbf{A})\mathbf{B} = \mathbf{A}r\mathbf{B}$$

where r is a scalar.

3. **Left distributive law** (Theorem 5)

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$$

4. **Right distributive law** (Theorem 6)

$$(\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA}$$

5. **Identity** (Theorem 7)

$$\mathbf{I}_m\mathbf{A} = \mathbf{A} = \mathbf{A}\mathbf{I}_n$$

Theorem 3 Given matrices $\mathbf{A} \in \mathbb{R}^{m \times l}$, $\mathbf{B} \in \mathbb{R}^{l \times p}$, and $\mathbf{C} \in \mathbb{R}^{p \times n}$, the following holds:

$$\mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}$$

Proof:

Since matrix-multiplication can be understood as a composition of functions, and since compositions of functions are associative, it follows that matrix-multiplication is associative

Theorem 4 Given matrices $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\mathbf{B} \in \mathbb{R}^{n \times p}$, the following holds:

$$r(\mathbf{AB}) = (r\mathbf{A})\mathbf{B} = \mathbf{A}(r\mathbf{B})$$

Proof:

First we prove $r(\mathbf{AB}) = (r\mathbf{A})\mathbf{B}$:

$$\begin{aligned} r(\mathbf{AB}) &= r \begin{bmatrix} \mathbf{Ab}_{*,1} & \dots & \mathbf{Ab}_{*,p} \end{bmatrix} \\ &= \begin{bmatrix} r\mathbf{Ab}_{*,1} & \dots & r\mathbf{Ab}_{*,p} \end{bmatrix} \\ &= (r\mathbf{A})\mathbf{B} \end{aligned}$$

Next, we prove $r(\mathbf{AB}) = \mathbf{A}(r\mathbf{B})$:

$$\begin{aligned} r(\mathbf{AB}) &= r \begin{bmatrix} \mathbf{Ab}_{*,1} & \dots & \mathbf{Ab}_{*,p} \end{bmatrix} \\ &= \begin{bmatrix} r\mathbf{Ab}_{*,1} & \dots & r\mathbf{Ab}_{*,p} \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{A}(r\mathbf{b}_{*,1}) & \dots & \mathbf{A}(r\mathbf{b}_{*,p}) \end{bmatrix} \quad \text{linearity of matrix-vector multiplication} \\ &= \mathbf{A}(r\mathbf{B}) \end{aligned}$$

□

Theorem 5 Given matrices $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{B} \in \mathbb{R}^{n \times p}$, and $\mathbf{C} \in \mathbb{R}^{n \times p}$, the following holds:

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$$

Proof:

$$\begin{aligned}
\mathbf{A}(\mathbf{B} + \mathbf{C}) &= \begin{bmatrix} \mathbf{A}(\mathbf{b}_{*,1} + \mathbf{c}_{*,1}) & \dots & \mathbf{A}(\mathbf{b}_{*,p} + \mathbf{c}_{*,p}) \end{bmatrix} && \text{definition of matrix multiplication} \\
&= \begin{bmatrix} (\mathbf{A}\mathbf{b}_{*,1} + \mathbf{A}\mathbf{c}_{*,1}) & \dots & (\mathbf{A}\mathbf{b}_{*,p} + \mathbf{A}\mathbf{c}_{*,p}) \end{bmatrix} && \text{linearity of matrix-vector multiplication} \\
&= \begin{bmatrix} \mathbf{A}\mathbf{b}_{*,1} & \dots & \mathbf{A}\mathbf{b}_{*,p} \end{bmatrix} + \begin{bmatrix} \mathbf{A}\mathbf{c}_{*,1} & \dots & \mathbf{A}\mathbf{c}_{*,p} \end{bmatrix} && \text{definition of matrix-addition} \\
&= \mathbf{AB} + \mathbf{AC} && \text{definition of matrix multiplication}
\end{aligned}$$

□

Theorem 6 Given matrices $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{B} \in \mathbb{R}^{n \times p}$, and $\mathbf{C} \in \mathbb{R}^{n \times p}$, the following holds:

$$(\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA}$$

Proof:

$$\begin{aligned}
(\mathbf{B} + \mathbf{C})\mathbf{A} &= \begin{bmatrix} (\mathbf{B} + \mathbf{C})\mathbf{a}_{*,1} & \dots & (\mathbf{B} + \mathbf{C})\mathbf{a}_{*,p} \end{bmatrix} && \text{definition of matrix-matrix multiplication} \\
&= \begin{bmatrix} (\mathbf{B}\mathbf{a}_{*,1} + \mathbf{C}\mathbf{a}_{*,1}) & \dots & (\mathbf{B}\mathbf{a}_{*,p} + \mathbf{C}\mathbf{a}_{*,p}) \end{bmatrix} && \text{definition of matrix addition} \\
&= \begin{bmatrix} \mathbf{B}\mathbf{a}_{*,1} & \dots & \mathbf{B}\mathbf{a}_{*,p} \end{bmatrix} + \begin{bmatrix} \mathbf{C}\mathbf{a}_{*,1} & \dots & \mathbf{C}\mathbf{a}_{*,p} \end{bmatrix} \\
&= \mathbf{BA} + \mathbf{CA}
\end{aligned}$$

□

Theorem 7 Given an $m \times n$ matrix \mathbf{A} , the following holds:

$$\mathbf{I}_m \mathbf{A} = \mathbf{A} = \mathbf{A} \mathbf{I}_n$$

Proof:

By the fact that an identity function simply maps each element in its domain back to itself. It follows that the composition of a function f and identity function is simply the function f . Thus, it follows that any matrix multiplied on the left or right by the identity matrix returns the original matrix. Thus, $\mathbf{I}_m \mathbf{A} = \mathbf{A} = \mathbf{A} \mathbf{I}_n$.

□