

Main Manuscript for

Uncovering thousands of endosymbiont DNA transfer events within single cockroach genomes

Kyle M. Ewart^{1,2*†}, Maxim W. D. Adams^{1†}, Zhuzhi Zhang^{1†}, Louise Baker³, Kokuto Fujiwara⁴, Yoshinobu Hayashi⁵, Leo A. Featherstone⁶, Oscar Lo Lu¹, Jil Helbling¹, Maya Moral¹, Kiyoto Maekawa⁷, Harley Rose¹, Aaron Jex^{3,8}, Simon Y. W. Ho¹, Nathan Lo¹

¹ School of Life and Environmental Sciences, University of Sydney, Sydney, New South Wales, Australia

² Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh, Edinburgh, UK

³ Infection and Global Health Division, WEHI, Parkville, Victoria, Australia

⁴ Hokkaido Center, Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Sapporo, Japan

⁵ Department of Biology, Keio University, Yokohama, Japan

⁶ The Kirby Institute, University of New South Wales, Sydney, New South Wales, Australia

⁷ Faculty of Science, Academic Assembly, University of Toyama, Toyama, Japan

⁸ Melbourne Veterinary School, University of Melbourne, Parkville, Victoria, Australia

[†] These authors contributed equally to this work.

* Corresponding author: Kyle M. Ewart

Email: kyle.ewart@sydney.edu.au

Author Contributions: K.M.E., M.W.D.A., Z.Z. and N.L. designed research; K.M.E., M.W.D.A., Z.Z., L.A.F., O.L.L., J.H., M.M. and N.L. analyzed data; Z.Z., L. B. and A.J. carried out laboratory work; and K.M.E., M.W.D.A., Z.Z. and N.L. wrote the paper; all authors revised the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: *Blattabacterium cuenoti*; Blattodea; Endosymbiosis; Horizontal gene transfer; Insect genomics

This file includes:

Main Text

Figures 1 to 2

Abstract

Horizontal gene transfer (HGT) between organisms can be a valuable source of genetic variation and innovation. Research on HGT in eukaryotes has hitherto focused on transfers of coding sequences; insertions of non-coding DNA remain poorly understood. Here, we investigated HGT in cockroaches, which have a long-standing evolutionary relationship with the transovarially transmitted endosymbiont *Blattabacterium cuenoti*, making them a valuable system for assessing the potential scale of HGT. We aligned 150 bp genomic fragments of *Blattabacterium cuenoti* to 23 cockroach and termite genomes, including 8 genomes newly sequenced, and revealed pervasive endosymbiont DNA transfer events. Australian panesthiine and geoscapheine cockroaches were consistently found to harbor >3000 HGT inserts, more than an order of magnitude higher than the previous maximum estimate in other eukaryotes, excluding rotifers. Some inserts appear to have persisted for >28.7 million years in this group, which may reflect functional roles. We identified numerous chimeric inserts comprising up to nine short segments from different locations in the *Blattabacterium cuenoti* genome. Our findings indicate pervasive HGT in eukaryote genomes, with potentially far-reaching implications for adaptation and speciation.

Main Text

Introduction

The number of DNA inserts acquired via horizontal gene transfer (HGT) that become fixed in a species is expected to represent only a small fraction of the total number of transfer events that occur over evolutionary time [1]. While there is an abundant body of research on insertions with homology to protein-coding sequence, transfers that do not encode proteins, or with unknown function, have been largely ignored. Nevertheless, such insertions abruptly introduce genetic variation that can persist if fitness effects are minimal, if purifying selection is insufficient, or if the inserts acquire functional roles [2].

We explored the prevalence and characteristics of DNA inserts originating from the endosymbiont *Blattabacterium cuenoti* (hereafter *Blattabacterium*), in the genomes of 14 cockroaches and 4 termites, including 8 newly sequenced using long-read technology. *Blattabacterium* is found in specialized cells of cockroach fat bodies and is inherited transovarially [3]. Infecting the ancestors of cockroaches >150–200 Mya [4], *Blattabacterium* was lost early in termite evolution. Although long-term obligate endosymbionts have been reported to contribute no or few DNA transfers to their host genomes, previous studies focused only on protein-coding sequences [5–8].

Results and Discussion

Cockroach genomes contain thousands of Blattabacterium-derived inserts

By focusing on transferred DNA inserts ≥ 50 bp and applying stringent filters to remove artifactual inserts and inserts potentially derived from duplication events (see SI Appendix), we characterized 40,485 *Blattabacterium*-derived inserts across 18 cockroach genomes (93–4900 per genome; Fig. 1A), with a median size of 161 bp (Fig. 1B). With the exception of rotifers, which have frequent HGT associated with anhydrobiosis [9], the highest number of transfers reported in any eukaryotic species was previously ~ 282 [8]. Inserts were clustered on some contigs, but generally were broadly distributed across host genomes (Fig. 1C; Dataset S6), and were inferred to have been transferred across multiple time points (Dataset S5), consistent with many independent HGT events (see SI Appendix).

Explanations for the high numbers of inserts we identified in cockroaches are as follows. First, we focused on *Blattabacterium*, an obligate endosymbiont whose genome is likely to come in close contact with host nuclear DNA during oogenesis and development. Second, we analyzed both intra- and intergenic regions of the genome. A recent study did not detect any horizontal transfers from *Blattabacterium* to cockroaches [8]; however, this is likely due to a focus on gene regions. Third, while previous studies filtered for the completeness of inserted coding sequences, we identified inserts as short as 50 bp. Fourth, previous studies often relied on short-read sequencing, where inserts spanning most of a sequence read may have been misidentified as bacterial contamination and discarded. In contrast, we did not actively exclude *Blattabacterium*-like sequences during assembly, and analysed genomes generated using both short- and long-read data (Dataset S2). Long-read data enables detection of inserts flanked by host sequence that may have been missed when using short-read data. Using the same pipeline, we identified comparatively low numbers of inserts from *Wolbachia*, and few to no putative inserts from *Buchnera* (which infects aphids, and not cockroaches) (see SI Appendix).

Panesthiinae and Geoscapheinae were found to possess particularly high numbers of *Blattabacterium*-derived inserts (Fig. 1A). Previous work found that this group has experienced considerable genome-wide relaxed selection [10], possibly due to reduced effective population sizes, which may allow larger numbers of inserts to persist. The absence of inserts in 4 termite genomes likely reflect the long-term absence of *Blattabacterium*.

Evolution of Blattabacterium-derived inserts in cockroach genomes

Examination of insert position across the cockroach genomes revealed that 15.38–25.56% were found in intragenic regions, and 74.44–84.62% were found in intergenic regions. This differs somewhat from the proportion of genic regions in the genomes, which was >25% in all but four (range 13.79–33.71%). The average size of inserts (+/- standard deviation) in intragenic regions (199.13 +/- 138.75 bp) was similar to that in intergenic regions (191.70 +/- 126.24 bp), although significantly higher in the former (one-way nested analysis of variance: $F_{1,39424} = 17.96$, $p < 0.01$; Fig. 1B). Insert sequences had GC content (17.9–29.6%) similar to those of *Blattabacterium* genomes (21.3–28.2%), but considerably lower than those of cockroach and termite genomes (34.5–41.1%). There was no apparent bias in the origin of inserts with respect to their position in the *Blattabacterium* genome, which is highly conserved in size, gene content, and gene order across strains (Fig. 1D).

Closely related taxa carried similar numbers of inserts, suggesting that many had been inherited vertically (Fig. 1A). We identified putatively ancestral HGT inserts by calculating their similarity to inserts in other species and to the host's specific *Blattabacterium* strain, and subsequently manually inspecting homology (and common origin) of cockroach genome sequence upstream and downstream from selected candidates ($n = 22$) across multiple taxa (Fig. 2A; see SI Appendix for details). The oldest ancestral inserts identified were inferred to have been present in the most recent common ancestor of all Panesthiinae and Geoscapheinae, at least ~28.7 Mya [11].

Transcription and chimeric nature of inserts

We used RNAseq data to investigate the transcription of inserts in *Panesthia t. tegminifera*, *Macropanesthia lithgowae*, and *Panesthia sloanei*. By mapping insert sequences to RNA contigs using BLASTn, we found that 91.42–94.96% of inserts showed no evidence of transcription (although RNA from all life stages and tissues was not assessed). Of the remaining inserts, which were observed in assembled RNA contigs, 26.59–32.97% overlapped with annotated genic

regions of the cockroach genomes (1.66–2.28% of total inserts), while the remainder overlapped with intergenic regions (3.38–6.30% of total inserts). A small fraction (0.40–0.65%) of inserts mapped to exon sequences within RNA contigs (see Dataset S4 for gene details).

During BLAST inspections of the longest inserts, we identified 32 inserts displaying a chimeric nature, comprising *Blattabacterium*-like sequences from distant areas within the *Blattabacterium* genome placed directly adjacent to each other within host genome contigs (Fig. 2B). These chimeric inserts were present in all cockroach genomes, comprising up to nine *Blattabacterium*-derived segments. Similar to chimeric numts previously identified in *Arabidopsis thaliana* [12], they are likely to have been produced by non-homologous end-joining repair that incorporated segments of *Blattabacterium* DNA that possibly arose following symbiont cell replication or death.

Conclusion

Our findings reveal extensive horizontal transfer of DNA from prokaryote symbionts to eukaryotes. Furthermore, the persistence of numerous inserts over millions of years indicates that they may have assumed functional roles in both genes and intergenic regions, are effectively neutral, or are only slightly deleterious. Future research on cockroaches and other species harboring obligate endosymbiotic prokaryotes will help to uncover any functional effects of inserts, providing a more comprehensive understanding of how HGT shapes genome evolution.

Materials and Methods

Detailed methods are described in the SI Appendix and on GitHub (https://github.com/MEEP-projects/HGT_pipeline). Briefly, we generated 8 new genome assemblies from species of Blaberidae (Dataset S1), and analyzed these with 15 previously generated genomes from Blattodea and 77 *Blattabacterium* (Dataset S2). To locate DNA transfers from *Blattabacterium* to host genomes, we aligned 150-bp fragments of the *Blattabacterium* genomes to the cockroach and termite genomes. Aligned fragments were characterized as putative HGT inserts, with a minimum length of 50 bp. Several filtering steps and QC analyses were applied to confirm that the identified inserts were ‘true’ HGT inserts, rather than assembly artifacts or misidentified *Blattabacterium* DNA, and to identify unique insert events within a genome (i.e., excluding inserts arising from duplication or transposition). We also ran control analyses using *Buchnera* (negative control) and *Wolbachia* (known to infect hosts) using the same pipeline. After characterizing inserts, we investigated their minimum age, length, origin, insertion locality, and GC content.

Acknowledgments

We thank James Walker, Bruce Gray for assistance with sample provision for genome sequencing, Pawsey Supercomputing Research Centre, and Australian BioCommons Leadership Share (ABLeS). This study was supported by grants from the Australian Research Council (FT160100463; DP220103265).

References

1. J. Van Etten, D. Bhattacharya, Horizontal gene transfer in eukaryotes: not if, but how much? *Trends Genet.* **36**, 915–925 (2020)
2. F. Husnik, J. P. McCutcheon, Functional horizontal gene transfer from bacteria to eukaryotes. *Nat. Rev. Microbiol.* **16**, 67–79 (2018).
3. L. Sacchi *et al.*, Symbionts in the oocytes of *Blattella germanica* (Dictyoptera: Blattellidae): Their mode of transmission. *Int. J. Insect Morph. Embryol.* **17**, 437–446 (1988).


4. N. Lo, C. Bandi, H. Watanabe, C. Nalepa, T. Beninati, Evidence for cocladogenesis between diverse dictyopteran lineages and their intracellular endosymbionts. *Mol. Biol. Evol.* **20**, 907–913 (2003).
5. N. Nikoh *et al.*, Bacterial genes in the aphid genome: absence of functional gene transfer from *Buchnera* to its host. *PLoS Genet.* **6**, e1000827 (2010).
6. D. B. Sloan *et al.*, Parallel histories of horizontal gene transfer facilitated extreme reduction of endosymbiont genomes in sap-feeding insects. *Mol. Biol. Evol.* **31**, 857–871 (2014).
7. Y. Li *et al.*, HGT is widespread in insects and contributes to male courtship in lepidopterans. *Cell*, **185**, 2975–2987 (2022).
8. C. Liu *et al.*, Horizontal gene transfers are widespread across termite genomes but do not confer metabolic innovations. *bioRxiv* [Preprint] (2025). <https://doi.org/10.1101/2025.03.16.643567> (accessed 6 July 2025).
9. P. Simion *et al.*, Chromosome-level genome assembly reveals homologous chromosomes and recombination in asexual rotifer *Adineta vaga*. *Sci. Adv.* **7**, eabg4216 (2021).
10. K. M. Ewart *et al.*, Pervasive relaxed selection in termite genomes. *Proc. R. Soc. B.* **291**, 20232439 (2024).
11. P. G. Beasley-Hall *et al.*, Digging deep: a revised phylogeny of Australian burrowing cockroaches (Blaberidae: Panesthiinae, Geoscapheinae) confirms extensive nonmonophyly and provides insights into biogeography and evolution of burrowing. *Syst. Entomol.* **46**, 767–783 (2021).
12. P. D. Fields *et al.*, Complete sequence of a 641-kb insertion of mitochondrial DNA in the *Arabidopsis thaliana* nuclear genome. *Genome Biol. Evol.* **14**, evac059 (2022).
13. D. A. Evangelista *et al.*, An integrative phylogenomic approach illuminates the evolutionary history of cockroaches and termites (Blattodea). *Proc. R. Soc. B.* **286**, 20182076 (2019).
14. M. W. D. Adams *et al.*, Pleistocene sea-level fluctuation shapes archipelago-wide population structure in the Endangered Lord Howe Island cockroach *Panesthia lata*. *Ecol. Evol.* **16**, e72760.

Figure legends


Figure 1. Dated evolutionary tree of analyzed taxa alongside total number of HGT inserts and number of unique inserts (excluding potential cases of duplication or transposition) (A). Ranges are given for taxa with multiple analyzed genomes. Node ages (Mya) are based on previous studies [11,13,14]. Length distribution of inserts (B). 50 bp represents our minimum length threshold, and inserts >1000 bp are omitted for clarity (51 inserts omitted). Cumulative number of inserts plotted along host genomes (C). Individual lines correspond to individual contigs (coloured by genome), such that each line starts at position 0 and terminates at the length of the contig. Cumulative number of inserts plotted along the *Blattabacterium* genome (D), based on the start position of each insert. Linear trends indicating random origins from the *Blattabacterium* genome. Only species with a published, host-specific *Blattabacterium* strain are included (Dataset S2). For species with multiple genomes, the version with the most inserts was analyzed (for B-D).

Figure 2. Putative ancestral insert evaluated by examination of flanking region homology (A). Homology of an insert (green bar) and its 5-kb flanking regions (blue bars) across species is shown on right (species lacking the insertion are shown with dashed lines on left). Arrowhead denotes inferred ancestral insertion event. Example of a chimeric HGT insertion in the *Panesthia tryoni tegminifera* genome (B). The *P. t. tegminifera* contig (top) is divided into seven segments of different origin, and the circular *Blattabacterium cuenoti* genome (CP142624.1; bottom) indicates the source of these insertions. Numbers indicate corresponding segments in the cockroach and *Blattabacterium cuenoti* genomes (with yellow labels marking genomic coordinates).


A




B




C



D



A



B

