


Assessment Complexity vs Applicability: The Case Study of the Cannibal European Hake in the Atlantic Iberian Waters

Anxo Paz¹; Santiago Cerviño¹; Marta Cousido-Rocha¹; Gustav Delius²; Francisco Velasco¹; Izaskun Preciado¹; Maria Grazia Pennino¹



¹Instituto Español de Oceanografía (IEO-CSIC)
²University of York

 anxo.paz@ieo.csic.es

 @a_paz_

 <https://mervex-group.github.io/MERVEX/proof/FRESCO.html>



Fisheries management is increasingly adopting an Ecosystem-Based Fisheries Management (EBFM) approach, which integrates not only environmental and anthropogenic factors, but also biological and socioeconomic interactions to promote sustainability. In this context, the **F**rom single stock ass**ES**sment to **e**COsystem-based fishery **man**agement (**FRESCO**) project arises, where we are developing the first MIZER (*Multi-Species Dynamic Size Spectrum Modeling in R*) model in the Iberian Atlantic, highlighting its potential to improve the understanding of ecological dynamics and facilitate its adoption in fisheries management. MIZER offers a framework for integrating ecological processes, such as size-dependent predation or cannibalism, into fisheries management.



Test the effect of modelling the cannibalism explicitly vs considering cannibalism in the definition of natural mortality.

Evaluate different management strategies

Use of single-species model as an evaluation model: test with European hake (*Merluccius merluccius*)



Future work

Improving the hake evaluation models inputs (natural mortality) using Mizer outputs

Studying effect in multi-species model