

Hamiltonian Systems

Manuel Gijón Agudo

Contents

1	Hamiltonian Equations	3
1.1	Notation	3
2	Hamiltonian formalism	4
3	Celestial mechanics	5
4	Geometric theory and invariant objects of Hamiltonian systems	6
5	Integrable systems	7
6	Quasi-integrable Hamiltonian systems	8
7	Lagrangian systems and variational methods	9
8	Hamiltonian Partial Differential Equations	10
9	Interactions between Dynamical Systems and Partial Differential Equations	11
10	Exercises	12
10.1	Chapter 1: Introduction to Hamiltonian systems	12
10.2	Chapter 2: The N-body problem	14
10.3	Chapter 3: Linear Hamiltonian systems	15
10.4	Chapter 6: Symplectic Transformations	15
10.5	Chapter 8: Geometric Theory	15
10.6	Chapter 9: Continuation of solutions	15
10.7	Chapter 10: Normal forms	15
10.8	Chapter 13: Stability and KAM Theory	15
11	Apendix	17
11.1	Complete examples	17
11.1.1	Harmonic oscillator	17
11.1.2	The Pendulum	17
11.2	Needed resoultls and definitions	17

11.2.1 Linear Algebra	17
11.2.2 Calculus	17
11.3 Geometry	17
11.4 Funcional Analysis	17
11.4.1 Differential forms	17
11.4.2 Measure Theory	17
11.4.3 Ordinary Differential Equations	17

1 Hamiltonian Equations

1.1 Notation

2 Hamiltonian formalism

3 Celestial mechanics

4 Geometric theory and invariant objects of Hamiltonian systems

5 Integrable systems

6 Quasi-integrable Hamiltonian systems

7 Lagrangian systems and variational methods

8 Hamiltonian Partial Differential Equations

9 Interactions between Dynamical Systems and Partial Differential Equations

10 Exercises

10.1 Chapter 1: Introduction to Hamiltonian systems

Make the phase portrait of the Hamiltonian system

$$\begin{aligned}\dot{x} &= y \\ \dot{y} &= x - \frac{x^3}{3}\end{aligned}$$

and compute its Hamiltonian

[Solución](#)

Make the phase portrait of the Hamiltonian system

$$\begin{aligned}\dot{x} &= x \\ \dot{y} &= -y + x^2\end{aligned}$$

and compute its Hamiltonian

[Solución](#)

(Meyer-Hall-Offin) Let x, y, z be the usual coordinates in \mathbb{R}^3 , $r = xi + yj + zk$, $X = \dot{x}$, $Y = \dot{y}$, $Z = \dot{z}$, $R = \dot{r} = Xi + Yj + Zk$.

1. Compute the three components of angular momentum $mr \times R$.
2. Compute the Poisson bracket of any two of the components of angular momentum and show that it is $\pm m$ times the third component of angular momentum.
3. Show that if a system admits two components of angular momentum as integrals, then the system admits all three components of angular momentum as integrals.

1. [adea](#)

2. [dsa](#)

3. [dadsa](#)

(Meyer-Hall-Offin) **A Lie algebra** A is a vector space with a product: $A \times A \rightarrow A$ that satisfies:

- **Anticommutative:** $ab \neq ba$
- **Distributive:** $a(b + c) = ab + ac$
- **Scalar associative:** $(\alpha a)b = \alpha(ab)$
- **Jacobis identity:** $a(bc) + b(ca) + c(ab) = 0$, $a, b, c \in A$, $\alpha \in \{\mathbb{R}, \mathbb{C}\}$

1. Show that vectors in \mathbb{R}^3 form a Lie algebra where the product $*$ is the cross product.
2. Show that smooth functions on an open set in \mathbb{R}^{2n} form a Lie algebra, where $fg = \{f, g\}$, the Poisson bracket.
3. Show that the set of all $n \times n$ matrices, $gl(n, \mathbb{R})$, is a Lie algebra, where $AB = AB - BA$, the Lie product.

1. bla

2. bla

3. bla

(Meyer-Hall-Offin) The pendulum equation is $\ddot{\theta} + \sin \theta = 0$.

1. Show that $2I = \frac{1}{2}\dot{\theta}^2 + (1 - \cos \theta) = \frac{1}{2}\dot{\theta}^2 + 2\sin^2(\theta/2)$ is an integral.
2. Sketch the phase portrait.
3. Make the substitution $y = \sin(\theta/2)$ to get $\dot{y}^2 = (1 - y^2)(I - y^2)$. Show that when $0 < I < 1$, $y = ksn(t, k)$ solves this equation when $k^2 = I$ (Look at the definition of elliptic sine function of Section 1.6 of Meyer-Hall-Offin).

1. bla

2. bla

3. bla

(Meyer-Hall-Offin) Let $H : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ be a globally defined conservative Hamiltonian, and assume that $H(z) \rightarrow +\infty$ as $|z| \rightarrow +\infty$. Show that all solutions of $\dot{z} = J\nabla H(z)$ are bounded (Hint: Think like Dirichlet).

Solución

Consider a \mathcal{C}^2 Hamiltonian $H = H(q, p, t) : U \subset \mathbb{R}^{2n+1} \rightarrow \mathbb{R}$ such that $\det(\partial_p^2 H) \neq 0$ on U . Define $v = \partial_p H(q, p, t)$. Prove:

1.

$$\begin{aligned}\partial_{q_i} L(q, v, t) &= -\partial_{q_i} H(q, p, t) \\ \partial_{v_i} L(q, v, t) &= p_i \\ \partial_t L(q, v, t) &= -\partial_t H(q, p, t)\end{aligned}$$

2. The Lagrangian L is \mathcal{C}^2 and $\det(\partial_v^2 L) \neq 0$.

3. The Euler-Lagrange equations associated to L and the Hamiltonian equations $\dot{q}_i = \partial_{p_i} H$, $\dot{p}_i = -\partial_{q_i} H$ are equivalent.

1. bla

2. bla

3. bla

10.2 Chapter 2: The N-body problem

Prove that the linear momentum is a first integral and that the center of mass moves with constant velocity for the 3 body problem.

[Solución](#)

Prove that if (a_1, a_2, \dots, a_N) is a central configuration with value λ :

1. For any $\tau \in \mathbb{R}$ then $(\tau a_1, \tau a_2, \dots, \tau a_N)$ is also a central configuration with value $\frac{\lambda}{\tau^3}$.
2. If A is an orthogonal matrix, then $Aa = (Aa_1, Aa_2, \dots, Aa_N)$ is also a central configuration with the same value λ .

1. bal bla

2. bla bla

(Meyer-Hall-Offin) Draw the complete phase portrait of the collinear Kepler problem. Integrate the collinear Kepler problem.

[Solución](#)

(Meyer-Hall-Offin) Show that $\varpi^2(\epsilon^2 - 1) = 2hc^2$ for the Kepler problem. (Attention: Meyer-Hall-Offin has a typo)

[Solución](#)

(Meyer-Hall-Offin) The area of an ellipse is $\pi a^2(1 - \epsilon^2)^{1/2}$, where a is the semi-major axis. We have seen in Keplers problem that area is swept out at a constant rate of $c/2$. Prove Keplers third law: The period p of a particle in a circular or elliptic orbit ($\epsilon < 1$) of the Kepler problem is $p = (\frac{2\pi}{\sqrt{\mu}})a^{3/2}$.

[Solución](#)

10.3 Chapter 3: Linear Hamiltonian systems

10.4 Chapter 6: Symplectic Transformations

10.5 Chapter 8: Geometric Theory

10.6 Chapter 9: Continuation of solutions

(Meyer-Hall-Offin) Show that the scaling used in Section 9.4 of Meyer-Hall-Offin to obtain Hills orbits for the restricted problem works for Hills lunar problem (see previous problem) also. Why does not the scaling for comets work?

[Solución](#)

Prove Lemma 9.7.1 in Meyer-Hall-Offin. Verify that formula (9.11) is the condition for an orthogonal crossing of the line of syzygy in Delaunay elements.

[Solución](#)

10.7 Chapter 10: Normal forms

10.8 Chapter 13: Stability and KAM Theory

(Meyer-Hall-Offin) Using Poincaré elements show that the continuation of the circular orbits established in Section 6.2 (Poincaré orbits) are of twist type and hence stable.

[Solución](#)

11 Apendix

11.1 Complete examples

11.1.1 Harmonic oscillator

11.1.2 The Pendulum

This is a case of a one DEGREE OF FREEDOM of second order, bla bla bla bla

11.2 Needed resoults and definitions

11.2.1 Linear Algebra

matriz ortogonal

bla bla bla bla

DEVOLVER A SU SITIO LAS FOTOS TAMAÑO CARNET

11.2.2 Calculus

teorema punto fijo de bla bla bla bla

Chain Rule: bla bla bla bla

proof:

reason why:

11.3 Geometry

11.4 Funcional Analysis

11.4.1 Differential forms

11.4.2 Measure Theory

11.4.3 Ordinary Differential Equations