

GAME Documentation

GAME Development Team

1 Overview of schemes

- Time stepping: Runge-Kutta third order scheme (RK3). In the vertical, at every substep, implicit methods are used.
- Corolis: [1] modified by [4]
- kinetic energy: [3]

2 Code structure

The code of the model resides in the directory `core/src`. It contains:

- The file `coordinator.c`. This is the file containing the main method. It allocates and deallocates memory and manages the calls to the time stepping scheme as well as the output function.
- The file `coordinator.h` contains the include statements of the file `coordinator.c`.
- The file `enum_and_typedefs.h` contains physical constants as well as typedefs.
- The directory `time_stepping` contains an implementation of the RK3 scheme as well as vertical column solvers.
- The directory `spatial_operators` contains the implementations of the spatial operators needed for the prognostic equations.
- The directory `diagnostics` contains functions computing diagnostic quantities needed for determining the values of the spatial operators.
- The directory `io` contains anything related to input and output.

3 Performance

The *speed* s of a model is defined by

$$s := \frac{\Delta t}{\Delta \tilde{t}}, \quad (1)$$

where Δt is the time step of the model and $\Delta \tilde{t}$ is the duration it takes to integrate from one time step to the next.

4 NWP mode

In NWP mode, one wants to be able to integrate one day within 8.5 minutes, which corresponds to $s = \frac{24 \cdot 60}{8.5} = 169$. As a general rule, one can say that the computation time is in equal parts needed for data assimilation, the dynamical core and the processes involving moisture and radiation. Thus, the dynamical core has a minimum speed of

$$s \geq 509. \quad (2)$$

4.1 Grid optimization

Hexagonal spherical grids need to be optimized for numerical modeling. Therefore, the Lloyd algorithm is used, which yields a *spherical centroidal Voronoi tessellation (SCVT)* after convergence [2]. [6] gives an overview of optimization alternatives and it seems to be that the SCVT is the most suitable for modeling. The procedure employed for executing the Lloyd algorithm is the one described in [5].

4.2 Parallelization

5 Radiation scheme

GAME employs the so-called RTE+RRTMGP (Radiative Transfer for Energetics + Rapid and Accurate Radiative Transfer Model for General Circulation Model Applications-Parallel) [7], [8] scheme. It is bound to the C code through the API RTE-RRTMGP-C [9].

References

- [1] J. Thuburn et al. Numerical representation of geostrophic modes on arbitrarily structured C-grids. In: *Journal of Computational Physics* 228 (22 2009), pp. 8321–8335.
- [2] Qiang Du, Max D. Gunzburger, and Lili Ju. Constrained Centroidal Voronoi Tessellations for Surfaces. In: *SIAM J. Sci. Comput.* 24.5 (Apr. 2003), pp. 1488–1506.
- [3] Almut Gassmann. A global hexagonal C-grid non-hydrostatic dynamical core (ICON-IAP) designed for energetic consistency. In: *Quarterly Journal of the Royal Meteorological Society* 139.670 (2013), pp. 152–175. DOI: 10.1002/qj.1960. eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/qj.1960>. URL: <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.1960>.
- [4] Almut Gassmann. Discretization of generalized Coriolis and friction terms on the deformed hexagonal C-grid. In: *Quarterly Journal of the Royal Meteorological Society* 144.716 (2018), pp. 2038–2053. DOI: 10.1002/qj.3294. eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3294>. URL: <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3294>.
- [5] Hiroaki Miura and Masahide Kimoto. A Comparison of Grid Quality of Optimized Spherical Hexagonal–Pentagonal Geodesic Grids. In: *Monthly Weather Review* 133.10 (Oct. 2005), pp. 2817–2833. ISSN: 0027-0644. DOI: 10.1175/MWR2991.1. eprint: https://journals.ametsoc.org/mwr/article-pdf/133/10/2817/4213997/mwr2991_1.pdf. URL: <https://doi.org/10.1175/MWR2991.1>.
- [6] Pedro S. Peixoto and Saulo R.M. Barros. Analysis of grid imprinting on geodesic spherical icosahedral grids. In: *Journal of Computational Physics* 237 (2013), pp. 61–78. ISSN: 0021-9991. DOI: <https://doi.org/10.1016/j.jcp.2012.11.041>. URL: <http://www.sciencedirect.com/science/article/pii/S0021999112007218>.
- [7] Robert Pincus, Eli J. Mlawer, and Jennifer S. Delamere. Balancing Accuracy, Efficiency, and Flexibility in Radiation Calculations for Dynamical Models. In: *Journal of Advances in Modeling Earth Systems* 11.10 (2019), pp. 3074–3089. DOI: 10.1029/2019MS001621. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019MS001621>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019MS001621>.
- [8] *RTE-RRTMGP github repository*. June 22, 2020. URL: <https://github.com/earth-system-radiation/rte-rrtmgp>.
- [9] *RTE-RRTMGP-C github repository*. June 22, 2020. URL: <https://github.com/MHBalsmeier/rte-rrtmgp-c>.