

# Core Temperature Responses to Cold-Water Immersion Recovery: A Pooled-Data Analysis

Jessica M. Stephens, Ken Sharpe, Christopher Gore, Joanna Miller, Gary J. Slater, Nathan Versey, Jeremiah Peiffer, Rob Duffield, Geoffrey M. Minett, David Crampton, Alan Dunne, Christopher D. Askew, and Shona L. Halson

**Purpose:** To examine the effect of postexercise cold-water immersion (CWI) protocols, compared with control (CON), on the magnitude and time course of core temperature ( $T_c$ ) responses. **Methods:** Pooled-data analyses were used to examine the  $T_c$  responses of 157 subjects from previous postexercise CWI trials in the authors' laboratories. CWI protocols varied with different combinations of temperature, duration, immersion depth, and mode (continuous vs intermittent).  $T_c$  was examined as a double difference ( $\Delta\Delta T_c$ ), calculated as the change in  $T_c$  in CWI condition minus the corresponding change in CON. The effect of CWI on  $\Delta\Delta T_c$  was assessed using separate linear mixed models across 2 time components (component 1, immersion; component 2, postintervention). **Results:** Intermittent CWI resulted in a mean decrease in  $\Delta\Delta T_c$  that was  $0.25^\circ\text{C}$  ( $0.10^\circ\text{C}$ ) (estimate [SE]) greater than continuous CWI during the immersion component ( $P = .02$ ). There was a significant effect of CWI temperature during the immersion component ( $P = .05$ ), where reductions in water temperature of  $1^\circ\text{C}$  resulted in decreases in  $\Delta\Delta T_c$  of  $0.03^\circ\text{C}$  ( $0.01^\circ\text{C}$ ). Similarly, the effect of CWI duration was significant during the immersion component ( $P = .01$ ), where every 1 min of immersion resulted in a decrease in  $\Delta\Delta T_c$  of  $0.02^\circ\text{C}$  ( $0.01^\circ\text{C}$ ). The peak difference in  $T_c$  between the CWI and CON interventions during the postimmersion component occurred at 60 min postintervention. **Conclusions:** Variations in CWI mode, duration, and temperature may have a significant effect on the extent of change in  $T_c$ . Careful consideration should be given to determine the optimal amount of core cooling before deciding which combination of protocol factors to prescribe.

**Keywords:** hydrotherapy, performance, exercise, ice bath, protocol variance

Cold-water immersion (CWI) is a widely practiced recovery modality aiming to reduce fatigue and facilitate postexercise recovery.<sup>1</sup> It is thought that the combination of cold temperature and hydrostatic pressure promotes reductions in tissue temperatures and blood flow, facilitating subsequent reductions in thermal and cardiovascular strain, edema, inflammation, and pain.<sup>1,2</sup> The dominant mechanism by which CWI is believed to be effective for acute recovery is its ability to ameliorate hyperthermia and the subsequent central nervous system mediated fatigue.<sup>1</sup> Indeed, previous research has attributed the enhanced recovery of maximal voluntary contraction force to faster return of central activation, which is the result of larger CWI-induced reductions in core temperature ( $T_c$ ).<sup>3</sup> With hyperthermia-mediated fatigue being a key fatiguing factor for many forms of exercise,<sup>1</sup> a greater understanding of the impact of CWI on  $T_c$  (as an indicator of hyperthermia) will enhance the effectiveness of CWI.

There is increasing evidence to support the notion that CWI enhances both short- and long-term recovery of performance, particularly for endurance and team sports. Several studies also have shown CWI to have either a negligible or detrimental effect on performance.<sup>1,4,5</sup> With consideration of this variance in findings, it may be that CWI is not suitable for all postexercise contexts, and the exercise mode performed prior to immersion, in addition to the time frame available and the environment in which exercise is performed (eg, hot vs cool vs thermoneutral) are key factors influencing the effectiveness of CWI.<sup>4,6</sup> Endurance-based performance has been shown to be most responsive to CWI; however, there is still considerable variability across studies assessing endurance performance.<sup>1,4</sup> For example, while a number of studies have found CWI to be effective for maintaining cycling time-trial performance in a subsequent exercise bout performed 40 minutes to 3 days post-CWI,<sup>7-10</sup> others observed a decrease in time-trial performance over the same time frame.<sup>11,12</sup> The factors responsible for this large variation in findings across the current literature are unclear, and as such, there is substantial debate as to the true efficacy of CWI as a recovery strategy.<sup>1,4</sup>

Variation in the physiological and performance recovery responses to CWI is likely to depend on the degree of cooling that can be achieved, which is a result of the initial interaction between the protocol utilized and the characteristics of the individual (eg, body composition, age, sex, ethnicity).<sup>4</sup> Understanding the optimal degree of cooling is important, as too little cooling may cause CWI to be less effective due to limited reductions in muscle temperature ( $T_m$ ) and  $T_c$ .<sup>13</sup> Conversely, too much cooling may lead to a reduction in muscle contractile force.<sup>14</sup> Current CWI protocols administered in practice vary in terms of the water temperature,

Stephens, Gore, Miller, Versey, and Halson are with the Dept of Physiology, Australian Inst of Sport, Canberra, ACT, Australia. Stephens, Slater, and Askew are with the School of Health and Sport Sciences, University of the Sunshine Coast, Maroochydore, QLD, Australia. Sharpe is with the Statistical Consulting Centre, School of Mathematics and Statistics, University of Melbourne, Melbourne, VIC, Australia. Gore is also with the Research Inst for Sport and Exercise, University of Canberra, Bruce, ACT, Australia. Peiffer is with the School of Psychology and Exercise Science, Murdoch University, Perth, WA, Australia. Duffield is with the Sport and Exercise Discipline Group, Health, University of Technology Sydney (UTS), Lindfield, NSW, Australia. Minett is with the School of Exercise and Nutrition Sciences, Queensland University of Technology, Kelvin Grove, QLD, Australia. Crampton and Dunne are with the Dept of Physiology, Trinity College Dublin, Ireland. Stephens ([jessica.stephens@act.gov.au](mailto:jessica.stephens@act.gov.au)) is corresponding author.

duration, depth, and mode of immersion, and the optimal combination of these factors remains unknown.<sup>4,14,15</sup> The interaction between each of these protocol factors is complex, and previous research has shown the same degree of  $T_c$  change (0.4°C) in response to different CWI protocols (eg, 5 min, 14°C, whole-body immersion<sup>16</sup> vs 5 min, 10°C, leg-only immersion<sup>17</sup>). However, it remains unknown whether the thermal stress applied by the temperature stimulus, the duration of exposure to the cold stimulus, the depth of immersion and body surface area exposed to the cold stimulus, or the change in temperature gradient by moving in and out of the water during intermittent immersion has the greatest impact on  $T_c$  responses.

Recently, it has been suggested that continuous immersion in water temperatures between 11°C and 15°C for 11 to 15 minutes is optimal for reducing muscle soreness.<sup>15</sup> However, the most effective approach for reducing  $T_c$  and exercise-induced hyperthermia remains unknown, and further research is required to understand how each factor contributes to  $T_c$  change.<sup>4</sup> With previous research showing that the change in  $T_c$  is related to a change in performance,<sup>3,9,18</sup> it is important to gain a greater understanding of  $T_c$  responses, as this will enable CWI protocols to be optimized and ultimately improve the restoration of performance for individual athletes. Therefore, the aims of this study were 2-fold: (1) to conduct a pooled analysis across a large data set to examine the impact of variability in different CWI protocol factors on  $T_c$  change relative to a control condition and (2) to characterize the time course of  $T_c$  responses to postexercise CWI both during immersion and postimmersion.

## Methods

### Study Design

This study adopted a pooled analysis approach using data from 157 male subjects from 13 previous investigations of postexercise CWI in our laboratories. Data were assessed using 2 respective linear mixed models based on different time components. The first component examined the change in  $T_c$  between the end of exercise and the end of the CWI/Control (CON) recovery intervention (component 1: immersion). The second component examined the postrecovery change only and is defined as the difference in  $T_c$  between the end of the CWI/CON recovery intervention and each of the available postintervention time points (component 2: postintervention).

### Data Sources

Individual deidentified raw data were collated from 13 previous studies by our groups for inclusion in this pooled analysis (Table 1). Criteria for inclusion were: (1) use of a cross-over controlled design, (2) included seated passive CON condition, (3) CWI performed postexercise, (4) measured  $T_c$  by rectal thermistor or telemetric pill, and (5) exercise resulted in a significant increase from baseline in mean  $T_c$  ( $\geq 38^\circ\text{C}$ ). Studies with missing data (where raw data could not be accessed) or without  $T_c$  measures immediately postexercise and/or postrecovery were excluded (Figure 1). There were no specific criteria for type of exercise utilized; however, 11 studies examined cycling<sup>8,10,11,20–27</sup> and 2 examined sprint running.<sup>3,19</sup> Of the 13 studies included, 10 are published in academic journals<sup>3,8,10,11,18–20,23,24,27</sup> and 3 in PhD theses.<sup>21,22,25</sup>

### Subjects

Deidentified raw data were extracted from 13 studies, providing data on 157 trained male subjects (Table 2). Subjects across all

studies were classified as well trained, with 94 identifying as predominantly participating in cycling or triathlon and 29 in team sports, leaving 34 with an unspecified sporting background.

### CWI Protocol Combinations

CWI protocols varied across studies, with 7 different temperatures, 8 immersion durations, 3 depths, and 2 modes of immersion utilized (Table 1), making a total of 336 possible combinations, of which 16 were utilized. Of the 13 studies included, 9 studies used just 1 CWI protocol,<sup>3,8,10,18,19,21,23,24,27</sup> 2 studies used 2 protocols,<sup>20,22</sup> 1 study included 3 protocols,<sup>11</sup> and another study used 4<sup>25</sup> different protocols, giving a total of 20 within-study-protocol combinations. Of these protocols, 4 were used in 2 studies so that there were only 16 of the 336 possible CWI protocols represented across the 13 studies. Further, there were only 15 (out of a possible 56) combinations of duration and temperature used, with just 1 combination used at more than 1 immersion depth. In addition, all 15 of these combinations were associated with just one of the 2 modes, continuous or intermittent, resulting in partial confounding between the 4 components of the CWI protocols so that it is not possible to completely separate the effects of the various (protocol) factors. For the pooled-data analysis and to allow comparisons between studies, immersion depth was converted into a predicted body-surface-water-contact area of 1.3 m<sup>2</sup> for waist depth, 1.6 m<sup>2</sup> for chest depth, and 1.8 m<sup>2</sup> for neck depth based on normative measurements of an average, and therefore comparable, male.<sup>28</sup> The offset time between the end of exercise and the commencement of CWI also varied, and 7 different offset times were used across the 13 studies (Table 1).

### Calculation of the Change in $T_c$

$T_c$  was either measured by rectal thermistor<sup>8,10,11,18,21,23,24,27</sup> or by sensor telemetry.<sup>3,19,20,22,25</sup>  $T_c$  was measured at different time points across the 13 studies (Table 1), including immediately postexercise, immediately postrecovery (0 min), and at 13 postrecovery time points (5, 10, 15, 20, 30, 40, 60, 90, 120, 150, 180, 210, and 240 min postintervention). Two of the studies<sup>3,23</sup> recorded just 2  $T_c$  values for each participant—one at the end of exercise and the other at the end of CWI—and were therefore only included in the immersion component analysis. The other 11 studies recorded  $T_c$  values at additional times following the completion of CWI and were therefore included in the postintervention component analysis. One study controlled postexercise  $T_c$  to ensure it was equal across subjects and trials,<sup>27</sup> whereas the remaining 12 studies did not attempt to control postexercise  $T_c$ . Regardless, there was no significant difference between trials (CWI vs CON) for each participant, as determined by initial *t*-test analysis. The  $T_c$  response was calculated in each of the models as a double difference ( $\Delta\Delta T_c$ ): the change in  $T_c$  in the CWI condition minus the corresponding difference under the control condition relative to postexercise in component 1 and immediately postrecovery in component 2 (eg,  $\Delta\Delta T_c = [\text{CWI postexercise } T_c - \text{CWI postrecovery } T_c] - [\text{CON postexercise } T_c - \text{CON postrecovery } T_c]$ ). A negative  $\Delta\Delta T_c$  indicates that the change in  $T_c$  is greater in the CWI condition compared with the control.

### Statistical Analysis

The statistical analysis consisted of 2 distinct components. The first component (immersion) considered the  $\Delta\Delta T_c$  changes from the end of exercise to the end of the recovery treatment, whereas the second component (postintervention) considered the  $\Delta\Delta T_c$  changes

Table 1 Data Sources

Study No.	Reference	Number of participants	CWI Conditions					$T_c$ method	$T_c$ measurement time points	Offset (EndEx to Rec0)	Ambient temp	EndEx $T_c$
			Duration	Temp	Depth	Mode	CON condition					
1	Peiffer et al <sup>11</sup>	12	<i>Condition 1:</i> 5 min	14°C	Chest	C	20 min, seated, room temperature 24°C	R	EndEx EndRec	25 min	40°C	38.9°C
			<i>Condition 2:</i> 10 min	14°C	Chest	C			PostRec: 5, 10, 20, 30, and 40 min			
			<i>Condition 3:</i> 20 min	14°C	Chest	C						
2	Peiffer et al <sup>29</sup>	8	20 min	14°C	Chest	C	20 min, seated, room temperature 24°C	R	EndEx EndRec	7.5 min	32°C	38.9°C
3	Peiffer et al <sup>10</sup>	10	5 min	14°C	Chest	C	15 min, seated, room temperature 35°C	R	EndEx EndRec	5 min	35°C	38.6°C
4	Stephens et al <sup>4</sup>	20	15 min	15°C	Neck	C	15 min, seated, room temperature 25°C	R	PostRec: 5 min EndEx EndRec	15 min	22°C	38.1°C
5	Minett et al <sup>3</sup>	9	20 min	10°C	Chest	C	20 min, seated, room temperature 32°C	G	EndEx EndRec	10 min	32°C	38.9°C
6	Vaile et al <sup>7</sup>	12	<i>Condition 1:</i> 5 min (5 × 1 min in; 2 min out) <i>Condition 2:</i> 5 min (5 × 1 min in; 2 min out) <i>Condition 3:</i> 5 min (5 × 1 min in; 2 min out) <i>Condition 4:</i> 15 min	10°C	Neck	I	14 min, seated, room temperature unknown	R	EndEx EndRec PostRec: 40 min	0 min	n/a	38.5°C
7	Pointon et al <sup>19</sup>	8	18 min (2 × 9 min in; 1 min out)	20°C	Neck	C	20 min, seated, room temperature 32°C		EndEx EndRec	10 min	32°C	39.0°C
8	Vaile et al <sup>8</sup>	12	14 min	15°C	Neck	C	14 min, seated, room temperature not reported	R	120 min Post EndEx EndRec	0 min	n/a	38.5°C

(continued)

Table 1 (continued)

Study No.	Reference	Number of participants	CWI Conditions					T <sub>c</sub> measurement method	T <sub>c</sub> measurement time points	Offset (EndEx to Rec0)	Ambient temp	EndEx T <sub>c</sub>
			Duration	Temp	Depth	Mode	CON condition					
9	Dunne et al <sup>20</sup>	9	Condition 1: 15 min	15°C	Waist	C	15 min, seated, room temperature 18°C	G	EndEx EndRec PostRec: 5 min	5 min	22°C	38.6°C
10	Stephens et al <sup>18</sup>	27	Condition 2: 15 min 15 min	8°C 15°C	Waist Neck	C C	15 min, seated, room temperature 25°C	R	EndEx EndRec PostRec: 5, 30, 60, 90, 120, 150, 180, 210, and 240 min	15 min	23°C	38.5°C
11	Versey <sup>21</sup>	9	14 min	15°C	Neck	C	14 min, seated, room temperature 21°C	R	EndEx EndRec PostRec: 5, 30, 60, and 90 min	15 min	22°C	38.8°C
12	Crampton <sup>22</sup>	10	Condition 1: 30 min Condition 2: 30 min	15°C	Waist	C	30 min, seated, room temperature 20°C	G	EndEx EndRec PostRec: 5 min	5 min	20°C	38.0°C
13	Halsen et al <sup>23</sup>	11	3 min (3 × 1 min in; 2 min out)	8°C 11°C	Waist Neck	C I	9 min, seated, room temperature 24°C	R	EndEx EndRec	20 min	24°C	39.6°C

Abbreviations: C, continuous; CON, control; CWI, cold-water immersion; EndEx, immediately postexercise; EndRec, immediately postrecovery; G, gastrointestinal temperature; I, intermittent; n/a, information not available; PostRec, postrecovery intervention; R, rectal temperature; Rec0, start of recovery intervention; T<sub>c</sub>, core temperature; temp, temperature.

following the recovery intervention. For each component, a linear mixed model was used with CWI protocols (combination of duration, temperature, depth, and mode), and the offset from the end of exercise to the start of the CWI treatment was treated as a fixed effects and either study-protocol (ie, the different protocols within a study were essentially treated as being different studies) or subject as random effects for components 1 and 2, respectively. Five of the 11 studies with data following the CWI treatment period included more than 1 post-CWI observation, and the models fitted to these data made allowance for possible autocorrelation within subjects. To fit these models, it was necessary to treat the subjects that used more than 1 protocol (within a study) as though they were different subjects. In addition to the effect of CWI treatment, it was also of interest to evaluate how the  $\Delta\Delta T_c$  varied with time postrecovery. When time postintervention was fitted as a (fixed effect) factor (only 13 time points were used in the studies), the relationship was deemed appropriate to then subsequently model using regression splines. All models were fitted using the lme or gamm components of the mgcv package<sup>30</sup> available in R.<sup>31</sup> The significance level was  $P \leq .05$  and data are reported as mean (SD) or estimate (SE).

## Results

### Component 1—Immersion

Across all subjects average postexercise  $T_c$  was  $38.56^\circ\text{C}$  ( $0.60^\circ\text{C}$ ) and immediately postintervention was  $37.72^\circ\text{C}$  ( $0.53^\circ\text{C}$ ). The



**Figure 1** — Flow chart on all relevant cold-water immersion studies performed in our laboratories and the reason for exclusion.

effects of CWI time, temperature, and mode are illustrated in Figure 2, which gives the estimated overall responses for each of the 20 study-protocol combinations used in the 13 studies. Intermittent CWI results in a significantly ( $P = .02$ ) greater decrease in  $\Delta\Delta T_c$   $0.25^\circ\text{C}$  ( $0.10^\circ\text{C}$ ) (estimate [SE]) than that obtained with continuous CWI. The effect of CWI temperature can be described by a significant ( $P = .05$ ) linear regression with a coefficient of  $0.03^\circ\text{C}$  ( $0.01^\circ\text{C}$ ). That is, for each reduction in CWI temperature of  $1^\circ\text{C}$ ,  $\Delta\Delta T_c$  is estimated to decrease on average by  $0.03^\circ\text{C}$ . The effect of CWI duration was significant ( $P = .01$ ), with a decrease of  $0.02^\circ\text{C}$  ( $0.01^\circ\text{C}$ )  $\Delta\Delta T_c$  for each additional minute of CWI immersion. Neither depth ( $P = .19$ ) nor offset time ( $P = .90$ ) had a significant effect on  $\Delta\Delta T_c$ .

The inclusion of the study-protocol in the model had a minimal effect on the parameter estimates, though it resulted in slight increases in the SEs and hence, slight increases in the  $P$  values. The residual SD, which includes the between-subject variation not accounted for by the fitted model and indicates the variation that was observed between the changes in individual subjects, was estimated to be  $0.44^\circ\text{C}$ .

### Component 2—Postintervention

The effect of offset time was significant ( $P = .00$ ), with an increase of  $0.01^\circ\text{C}$   $\Delta\Delta T_c$  for each minute increase in offset time. Further, the effect of postrecovery time was also significant ( $P < .001$ ) and was adequately described by a cubic regression spline. Specifically, peak difference between CWI and CON occurred at  $\sim 60$  minutes postintervention; following this,  $\Delta\Delta T_c$  slowly increased until the intervention had no impact (Figure 3). Also displayed in Figure 3 are estimates of the effect of postrecovery time when it was treated as a factor (with 13 levels, the number of different times used in the studies). Other effects such as CWI type (intermittent or continuous), duration, temperature, and depth were also evaluated, but none of them made a (statistically) significant contribution.


The inclusion of within-subject autocorrelation in the model had an appreciable effect on the parameter estimates, with the autocorrelation being highly significant ( $P < .001$ ). The residual SD, which includes within-subject variation not accounted for by the fitted model, was estimated to be  $0.36^\circ\text{C}$ .

**Table 2** Participant Characteristics in Each Study


Study No.	Reference	Height, cm	Body mass, kg	Age, y	VO <sub>2</sub> max, mL·kg <sup>-1</sup> ·min <sup>-1</sup>
1	Peiffer et al <sup>11</sup>	181.0 (6.0)	77.9 (6.6)	27.0 (7.0)	61.7 (5.0)
2	Peiffer et al <sup>29</sup>	178.8 (5.4)	77.1 (6.5)	29.3 (3.0)	64.0 (5.7)
3	Peiffer et al <sup>10</sup>	182.6 (7.0)	80.3 (9.7)	n/a	n/a
4	Stephens et al <sup>4</sup>	181.9 (7.9)	78.7 (9.6)	32.1 (7.5)	59.7 (6.2)
5	Minett et al <sup>3</sup>	183.0 (7.0)	78.7 (8.1)	21.0 (2.0)	n/a
6	Vaile et al <sup>7</sup>	181.3 (4.6)	76.4 (7.1)	32.8 (3.8)	69.9 (4.8)
7	Pointon et al <sup>19</sup>	179.6 (3.8)	78.9 (6.3)	19.9 (1.1)	n/a
8	Vaile et al <sup>8</sup>	176.6 (4.5)	68.8 (7.2)	32.2 (4.3)	68.8 (3.6)
9	Dunne et al <sup>20</sup>	177.0 (5.0)	68.0 (5.0)	29.0 (7.0)	62.1 (5.0)
10	Stephens et al <sup>18</sup>	181.7 (7.5)	83.2 (11.9)	32.7 (7.9)	55.8 (7.9)
11	Versey <sup>21</sup>	177.2 (5.3)	74.3 (8.4)	29.9 (5.6)	62.0 (5.2)
12	Crampton <sup>22</sup>	184.0 (5.0)	86.0 (86.0)	26.0 (5.0)	54.6 (7.4)
13	Halsen et al <sup>23</sup>	182.2 (4.2)	72.1 (4.0)	23.8 (1.6)	71.3 (1.2)

Abbreviations: n/a, information not available; VO<sub>2</sub>max, volume of oxygen consumption at maximum exertion. Note: Data are presented as mean (SD).





**Figure 2** — Estimated responses for each of the 20 study-protocol combinations used in the 13 studies. Numbers next to data points indicate water temperature.  $\Delta\Delta T_c$  = change in  $T_c$  in CWI condition minus change in  $T_c$  in CON condition. CON indicates control;  $T_c$ , core temperature; CWI, cold-water immersion.



**Figure 3** — Parameter estimates and fitted spline with 95% confidence limits for the change in  $T_c$  from end of intervention to each of the postintervention time points.  $\Delta\Delta T_c$  = change in  $T_c$  in CWI condition minus change in  $T_c$  in CON condition. CON indicates control;  $T_c$ , core temperature; CWI, cold-water immersion.

## Discussion

This study aimed to understand the implications of varying the temperature, duration, depth, and mode of CWI protocols on  $T_c$ ,

and to identify the ensuing time course of  $T_c$  responses based on these postexercise CWI protocol variations. The main findings were that (1) intermittent protocols resulted in a significantly greater decrement in  $T_c$  compared with continuous protocols for the  $T_c$  change during immersion, (2) decreasing water temperature and increasing duration of CWI resulted in a significant decrease in  $\Delta\Delta T_c$  during immersion, (3) the longer the offset time (end of exercise to immersion commencement), the smaller the change in  $T_c$  postrecovery, and (4) the peak difference in  $T_c$  between CON and CWI protocols occurred at ~60 minutes postrecovery, irrespective of protocol mode.

Reported postexercise CWI protocols vary substantially,<sup>4,15</sup> and while CWI is widely utilized by athletes, a lack of consensus as to the best protocols for different sport/athlete scenarios remains.<sup>14</sup> Accordingly, this study combined the data from a range of studies representing the variety of protocols currently utilized to determine the impact of different combinations and interaction of these factors on the change in  $T_c$ . One of the major findings of this study was that intermittent CWI protocols appear to be more effective in lowering  $T_c$  compared with continuous CWI. It may be postulated that the lower  $T_c$  observed, on average, in response to intermittent CWI might be related to the frequent change in thermal gradient occurring each time the participant moves between the cold water and the warmer air. This frequent change may have led to repeated reactive hyperemia responses where both skin and muscle blood flow increases when the participant moves out of the pool after a period of cold-induced vasoconstriction and ischemia which occurs during immersion.<sup>32</sup> This theory is supported by the findings of Romet<sup>33</sup> and Seo et al<sup>34</sup> who found that, following removal from CWI, vasodilation occurred in the extremities and greater conductive heat transfer occurred due to the return of cooler blood to the central circulation. Nevertheless, as only 3 studies utilized intermittent protocols, the conclusions which can be drawn from these data need to be confirmed by future research.

Often in practical settings, the **duration** and **depth** of CWI are determined by the water temperature based on athlete tolerance; thus, these variables were also examined in this study given their ecological interactions in many protocols. Although it has been suggested that the physiological changes in response to postexercise CWI are temperature dependent,<sup>14</sup> the way these factors interact with each other and which factor has the greatest impact on  $T_c$  responses remains **unknown**.<sup>4</sup> Both **temperature** and **duration** were found to have a **highly significant** impact on  $\Delta\Delta T_c$ . This study found that CWI temperature led to a decrease in  $\Delta\Delta T_c$  of **0.025°C** for every **1°C** reduction in water temperature, and that CWI duration led to a reduction in  $\Delta\Delta T_c$  of **0.018°C** for every additional minute of **immersion** time. Collectively, colder water temperatures and greater immersion durations lead to a **greater reduction** in  $T_c$  compared with an equivalent duration CON. However, such an effect was only observed for **continuous immersion** protocols, as no evidence of a duration effect was apparent for **intermittent** protocols given the small range of intermittent protocols included in the analyses. The **depth** of immersion was **not significant** and highly confounded with the other protocol factors. Increasing immersion **depth** is believed to enhance responses to CWI by increasing **hydrostatic** pressure as well as exposing a greater **body surface** area for thermal exchange through convection to occur.<sup>4</sup> The impact of hydrostatic pressure was recently examined by comparing seated versus standing CWI, with no significant difference reported between the 2 conditions, suggesting water temperature may be of greater importance.<sup>35</sup> Given the absence of studies examining the effect of different immersion depths on  $T_c$  responses to postexercise CWI, further research is required to fully determine the impact of varying CWI depth.

**Postexercise CWI** has been shown to **significantly reduce**  $T_c$ ; however, the extent of this reduction is highly **variable**, and the time course of change remains to be fully elucidated.<sup>4</sup> This study examined the change in  $T_c$  during and postimmersion as 2 separate components, as it was recognized that the rate of  $T_c$  change would be vastly different depending on the **thermal environment** the body is placed in. This study found that the sooner CWI is commenced postexercise, the greater the **reduction** in postimmersion  $T_c$  will be. This may be due to  $T_c$  and blood flow being elevated at the end of exercise, therefore increasing the **thermal gradient** between the body and the water and thermal exchange between blood and body tissues. It was also found that when examining  $T_c$  change post-recovery, the greatest difference between CWI and CON occurred at **60 minutes** postrecovery (Figure 2). This **novel** finding highlights the importance of this **time period postimmersion**, and it highlights the potentially **negative effect** of a **hot shower** post-immersion. A **hot shower** immediately postimmersion is a common practice of some athletes, which may prevent the afterdrop in  $T_c$ , therefore potentially limiting the effectiveness of CWI on core cooling. However, with only 3 studies examining  $T_c$  change for  $\geq 60$  minutes postimmersion, the estimates of  $\Delta\Delta T_c$  become weaker as time increases, potentially limiting the strength of conclusions which can be drawn.

This prolonged decrease in  $T_c$  after CWI may have practical implications for repeat performance and should be considered when prescribing protocols. It is hypothesized that the **optimal** protocol **parameters** will **vary** depending on recovery needs of the athlete, which will be determined by the specific **type of fatigue** (eg, central nervous system fatigue, cardiovascular fatigue), time frame available, and type of performance (eg, **endurance** vs **sprint**) required.<sup>1,4</sup> It is also important to consider the environmental conditions.<sup>1,36</sup> For example, performing CWI during a **short** time frame between endurance tasks may provide precooling

benefits for subsequent exercise, particularly when environmental conditions are **warm** or **hot**. However, when performance requires **maximal contractions** and the time frame between repeat performances is short, CWI-induced changes in body temperature will likely **reduce muscular performance**.<sup>4,15</sup>

Future studies should focus on determining the exact degree of change in  $T_c$  that leads to an **optimal cooling** effect for subsequent performance and how different CWI protocol factors work toward inducing this  $T_c$  change. Future research should also look to establish the **optimal cooling effect for other physiological variables** such as **muscle temperature** and blood flow, as these also have the potential to impact performance recovery. The residual SDs were estimated to be 0.44°C and 0.36°C for components 1 and 2, respectively. Compared with the estimated effects of CWI, these values are relatively large, which means that, while various effects have been found, on average, to be statistically significant, there is a lot of **additional variation between subjects** (for component 1) and within subjects (for component 2), so it is not yet possible to deduce how individual athletes will respond to CWI.

The present **pooled-data analysis** study builds on a previous meta-analysis<sup>37</sup> by examining individual responses to a range of CWI protocols, and highlights the fact that responses to postexercise CWI are **highly variable** and are impacted by myriad factors. It is not solely the dose of cooling provided by the combination of CWI temperature, duration, depth, and mode that impacts these responses. Other factors such as laboratory/**environmental conditions**, differences in exercise-induced **thermoregulatory stress**, **offset differences** (ie, time between end of exercise and start of CWI), and individual participant differences (eg, body composition, age, sex, and ethnicity) also impact responses and may **explain much of the variation** in the current literature. The relatively **homogenous cohort** examined in this pooled analysis acts to delimit several of these potentially confounding factors (eg, sex, age, body composition), yet this may also limit the applicability of findings to other populations. The large number of factors impacting the cooling response makes attempting to predict the optimal “dose” of CWI quite difficult, especially when many combinations of factors have not been tested. Nevertheless, this study has drawn on a large data set to provide some clarity around the influence of CWI protocol mode, temperature, duration, and offset differences on  $T_c$  response. An understanding of how variations in these factors impact temperature change will enable future researchers to better prescribe CWI protocols, and will ultimately facilitate the optimization of performance recovery.

## Practical Applications

1. Before prescribing a CWI protocol, it is important to determine **how much core cooling** needs to be **induced**. For situations where more **intense** cooling is required, **longer duration** and colder water temperatures may be more **effective**.
2. When greater reductions in  $T_c$  are required, CWI should be performed **as soon as possible** after **exercise**.
3. Intermittent CWI protocols are effective in reducing  $T_c$  and can be used when there are a large number of athletes who need to complete CWI with limited resources (eg, 1 ice bath) or when an athlete is uncomfortable with long-duration CWI.
4. Consideration should be given to what activities the athletes have in the **60-minute postimmersion** time frame, as  $T_c$  continues to decrease during this period.

## Acknowledgments

The authors would like to thank all of the researchers who contributed to each of the studies utilized in this analysis. The authors would like to acknowledge Dr Andrew Govus for his assistance in the early stages of planning this study. This investigation was supported by funding from a University of the Sunshine Coast Faculty Higher Degree by Research grant. The authors declare that there are no conflicts of interest in undertaking this study.

## References

- Ihsan M, Watson G, Abbiss CR. What are the physiological mechanisms for post-exercise cold water immersion in the recovery from prolonged endurance and intermittent exercise? *Sports Med.* 2016;46(8):1095–1109. PubMed ID: [26888646](#) doi:[10.1007/s40279-016-0483-3](#)
- Bleakley CM, Davison GW. What is the biochemical and physiological rationale for using cold-water immersion in sports recovery? A systematic review. *Br J Sports Med.* 2010;44(3):179–187. PubMed ID: [19945970](#) doi:[10.1136/bjism.2009.065565](#)
- Minett GM, Duffield R, Billaut F, Cannon J, Portus MR, Marino FE. Cold-water immersion decreases cerebral oxygenation but improves recovery after intermittent-sprint exercise in the heat. *Scand J Med Sci Sports.* 2014;24(4):656–666. PubMed ID: [23458430](#) doi:[10.1111/sms.12060](#)
- Stephens JM, Halson S, Miller J, Slater GJ, Askew CD. Cold-water immersion for athletic recovery: one size does not fit all. *Int J Sports Physiol Perform.* 2017;12(1):2–9. PubMed ID: [27294485](#) doi:[10.1123/ijssp.2016-0095](#)
- Bieuzen F, Bleakley CM, Costello JT. Contrast water therapy and exercise induced muscle damage: a systematic review and meta-analysis. *PLoS ONE.* 2013;8(4):e62356. PubMed ID: [23626806](#) doi:[10.1371/journal.pone.0062356](#)
- Minett GM, Costello JT. Specificity and context in post-exercise recovery: it is not a one-size-fits-all approach. *Front Physiol.* 2015;6:130. PubMed ID: [25964762](#) doi:[10.3389/fphys.2015.00130](#)
- Vaile J, Halson S, Gill N, Dawson B. Effect of cold water immersion on repeat cycling performance and thermoregulation in the heat. *J Sports Sci.* 2008;26(5):431–440. PubMed ID: [18274940](#) doi:[10.1080/02640410701567425](#)
- Vaile J, Halson S, Gill N, Dawson B. Effect of hydrotherapy on recovery from fatigue. *Int J Sports Med.* 2008;29(7):539–544. PubMed ID: [18058595](#) doi:[10.1055/s-2007-989267](#)
- Vaile J, O'Hagan C, Stefanovic B, Walker M, Gill N, Askew CD. Effect of cold water immersion on repeated cycling performance and limb blood flow. *Br J Sports Med.* 2011;45(10):825–829. PubMed ID: [20233843](#) doi:[10.1136/bjism.2009.067272](#)
- Peiffer JJ, Abbiss CR, Watson G, Nosaka K, Laursen PB. Effect of a 5-min cold-water immersion recovery on exercise performance in the heat. *Br J Sports Med.* 2010;44(6):461–465. PubMed ID: [18539654](#) doi:[10.1136/bjism.2008.048173](#)
- Peiffer JJ, Abbiss CR, Watson G, Nosaka K, Laursen PB. Effect of cold-water immersion duration on body temperature and muscle function. *J Sports Sci.* 2009;27(10):987–993. PubMed ID: [19847682](#) doi:[10.1080/02640410903207424](#)
- Buchheit M, Peiffer JJ, Abbiss CR, Laursen PB. Effect of cold water immersion on postexercise parasympathetic reactivation. *Am J Physiol Heart Circ Physiol.* 2009;296(2):H421–H427. PubMed ID: [19074671](#) doi:[10.1152/ajpheart.01017.2008](#)
- Getto CN, Golden G. Comparison of active recovery in water and cold-water immersion after exhaustive exercise. *Athl Train Sports Health Care.* 2013;5(4):169–176.
- Machado AF, Ferreira PH, Micheletti JK, et al. Can water temperature and immersion time influence the effect of cold water immersion on muscle soreness? A systematic review and meta-analysis. *Sports Med.* 2016;46(4):503–514. PubMed ID: [26581833](#) doi:[10.1007/s40279-015-0431-7](#)
- Machado AF, Almeida AC, Micheletti JK, et al. Dosages of cold-water immersion post exercise on functional and clinical responses: a randomized controlled trial. *Scand J Med Sci Sports.* 2017;27(11):1356–1363. PubMed ID: [27430594](#) doi:[10.1111/sms.12734](#)
- Peiffer JJ, Abbiss CR, Watson G, Nosaka K, Laursen PB. Effect of cold water immersion on repeated 1-km cycling performance in the heat. *J Sci Med Sport.* 2010;13(1):112–116. PubMed ID: [18948061](#) doi:[10.1016/j.jsams.2008.08.003](#)
- Stanley J, Peake JM, Coombes JS, Buchheit M. Central and peripheral adjustments during high-intensity exercise following cold water immersion. *Eur J Appl Physiol.* 2014;114(1):147–163. PubMed ID: [24158407](#) doi:[10.1007/s00421-013-2755-z](#)
- Stephens JM, Halson S, Miller J, Slater G, Chapman D, Askew C. Effect of body composition on physiological responses to cold-water immersion and the recovery of exercise performance. *Int J Sports Physiol Perform.* 2018;13(3):382–389. doi:[10.1123/ijssp.2017-0083](#)
- Pointon M, Duffield R, Cannon J, Marino FE. Cold water immersion recovery following intermittent-sprint exercise in the heat. *Eur J Appl Physiol.* 2012;112(7):2483–2494. PubMed ID: [22057508](#) doi:[10.1007/s00421-011-2218-3](#)
- Dunne A, Crampton D, Egana M. Effect of post-exercise hydrotherapy water temperature on subsequent exhaustive running performance in normothermic conditions. *J Sci Med Sport.* 2013;16(5):466–471. PubMed ID: [23246445](#) doi:[10.1016/j.jsams.2012.11.884](#)
- Versey NG. *Hydrotherapy and Recovery From Exercise Induced Fatigue: Performance Effects and Mechanisms Involved* [dissertation]. Perth, WA, Australia: University of Western Australia Research Repository: School of Sport Science, University of Western Australia; 2012.
- Crampton D. *The Use of Water Immersion as a Recovery Intervention Following High-Intensity Exercise: An Investigation of the Physiological and Performance Effects.* [dissertation]. Dublin, Ireland: Trinity's Access to Research Archive: Department of Physiology, Trinity College Dublin; 2012.
- Halson SL, Quod MJ, Martin DT, Gardner AS, Ebert TR, Laursen PB. Physiological responses to cold water immersion following cycling in the heat. *Int J Sports Physiol Perform.* 2008;3(3):331–346. PubMed ID: [19211945](#) doi:[10.1123/ijssp.3.3.331](#)
- Peiffer JJ, Abbiss CR, Nosaka K, Peake JM, Laursen PB. Effect of cold water immersion after exercise in the heat on muscle function, body temperatures, and vessel diameter. *J Sci Med Sport.* 2009;12(1):91–96. PubMed ID: [18083634](#) doi:[10.1016/j.jsams.2007.10.011](#)
- Vaile J. *Effect of Hydrotherapy on Recovery of Muscle-Damage and Exercise-Induced Fatigue: School of Sports Science* [dissertation]. Perth, WA, Australia: University of Western Australia; 2008.
- Stephens J, Halson S, Vaile J, Slater G, Askew C. Effect of body composition on core temperature responses to post-exercise cold water immersion. *J Sci Med Sport.* 2015;19:e4. doi:[10.1016/j.jsams.2015.12.390](#)
- Stephens JM, Halson SL, Miller J, Slater GJ, Askew CD. Influence of body composition on physiological responses to post-exercise hydrotherapy. *J Sports Sci.* 2018;36(9):1044–1053. doi:[10.1080/02640414.2017.1355062](#)
- de Leva P. Adjustments to Zatsiorsky-Seluyanov's segment inertia parameters. *J Biomech.* 1996;29(9):1223–1230. PubMed ID: [8872282](#) doi:[10.1016/0021-9290\(95\)00178-6](#)



29. Peiffer JJ, Abbiss CR, Nosaka K, Peake JM, Laursen PB. Effect of cold water immersion after exercise in the heat on muscle function, body temperatures, and vessel diameter. *J Sci Med Sport*. 2009;12(1):91–96.
30. Wood SN. Stable and efficient multiple smoothing parameter estimation for generalized additive models. *J Am Stat Assoc*. 2004;99:673–686. doi:10.1198/016214504000000980
31. Team RC. R: a language and environment for statistical computing. 2014. <http://www.R-project.org/>. Accessed April 4, 2016.
32. White GE, Rhind SG, Wells GD. The effect of various cold-water immersion protocols on exercise-induced inflammatory response and functional recovery from high-intensity sprint exercise. *Eur J Appl Physiol*. 2014;114(11):2353–2367. PubMed ID: 25074283 doi:10.1007/s00421-014-2954-2
33. Romet TT. Mechanism of afterdrop after cold water immersion. *J Appl Physiol*. 1988;65(4):1535–1538. PubMed ID: 3182516 doi:10.1152/jappl.1988.65.4.1535
34. Seo Y, Kim CH, Ryan EJ, Gunstad J, Glickman EL, Muller MD. Cognitive function during lower body water immersion and post-immersion afterdrop. *Aviat Space Environ Med*. 2013;84(9):921–926. PubMed ID: 24024303 doi:10.3357/ASEM.3571.2013
35. Leeder JD, van Someren KA, Bell PG, et al. Effects of seated and standing cold water immersion on recovery from repeated sprinting. *J Sports Sci*. 2015;33(15):1544–1552. PubMed ID: 25573221 doi:10.1080/02640414.2014.996914
36. Halson S. Does the time frame between exercise influence the effectiveness of hydrotherapy for recovery? *Int J Sports Physiol Perform*. 2011;6:147–159. PubMed ID: 21725101 doi:10.1123/ijsp.6.2.147
37. Zhang Y, Davis JK, Casa DJ, Bishop PA. Optimizing cold water immersion for exercise-induced hyperthermia: a meta-analysis. *Med Sci Sports Exerc*. 2015;47(11):2464–2472. PubMed ID: 25910052 doi:10.1249/MSS.0000000000000693

Copyright of International Journal of Sports Physiology & Performance is the property of Human Kinetics Publishers, Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.